

별첨 2

배제문헌

문헌배제사유

1. 사전에 정의한 연구대상자에 대한 연구가 아닌 문헌
2. 사전에 정의한 중재법에 대해 연구가 아닌 문헌
3. 사전에 정의한 연구결과가 하나 이상 보고되지 않은 문헌
4. 사전에 정의한 연구설계에 해당하지 않은 문헌
5. 원저가 아닌 연구(총설, letter, comment 등)
6. 동물실험 또는 전임상시험
7. 한국어나 영어로 출판되지 않은 문헌

| 연번 | 서지정보 | 배제 사유 |
|----|---|-------|
| 1 | Actrn. BRAIN Training Trial: balance, Resistance, or INterval Training Trial: a Randomised Controlled Trial of Three Exercise Modalities in Mild Cognitive Impairment. 2017. | 1 |
| 2 | Actrn. Evaluation of the effects of the Good Cholesterol on Wound Healing. http://www.who.int/trialssearch/Trial2.aspx?TrialID=ACTRN12619000269134 . 2019. | 2 |
| 3 | Amin N, Doupis J. Diabetic foot disease: From the evaluation of the "foot at risk" to the novel diabetic ulcer treatment modalities. World Journal of Diabetes. 2016;7(7):153-64. | 5 |
| 4 | Ammendola M, Sacco R, Butrico L, Sammarco G, de Franciscis S, Serra R. The care of transmetatarsal amputation in diabetic foot gangrene. International Wound Journal. 2017;14(1):9-15. | 4 |
| 5 | Andersen CA, Ponticello M, Byerley N, Marmolejo V. Insulated Offloading Provides Offloading Protection and Enhanced Skin Perfusion. Wounds-A Compendium of Clinical Research & Practice. 2018;30(10):300-5. | 2 |
| 6 | Anderson TJ, Phillips SA. Assessment and prognosis of peripheral artery measures of vascular function. Progress in Cardiovascular Diseases. 2015;57(5):497-509. | 1 |
| 7 | Anonymous. Abstracts from Issue 42 of EJVES Short Reports: Editor: Jean-Baptiste Ricco. European Journal of Vascular and Endovascular Surgery. 2019;57 (4):607-9. | 5 |
| 8 | Antonopoulos CN, Lazaris A, Venermo M, Geroulakos G. Predictors of Wound Healing Following Revascularization for Chronic Limb-Threatening Ischemia. Vascular & Endovascular Surgery. 2019;53(8):649-57. | 5 |
| 9 | Aref Z, de Vries MR, Quax PHA. Variations in surgical procedures for inducing hind limb ischemia in mice and the impact of these variations on neovascularization assessment. International Journal of Molecular Sciences. 2019;20 (15) (3704). | 5 |
| 10 | Arenas IA, Ergui I, Lamas GA. Safety of chelation therapy with EDTA in patients with critical limb ischemia: A pilot trial of limb preservation in diabetic patients. Arteriosclerosis, Thrombosis, and Vascular Biology Conference: American Heart Association's Arteriosclerosis, Thrombosis and Vascular Biology/Peripheral Vascular Disease. 2017;37(Supplement 1). | 5 |

| 연번 | 서지정보 | 배제 사유 |
|----|--|-------|
| 11 | Arima Y, Hokimoto S, Tabata N, Nakagawa O, Oshima A, Matsumoto Y, et al. Evaluation of Collateral Source Characteristics With 3-Dimensional Analysis Using Micro-X-Ray Computed Tomography. <i>Journal of the American Heart Association</i> . 2018;7(6):23. | 2 |
| 12 | Abejon D, Krames ES. Peripheral nerve stimulation or is it peripheral subcutaneous field stimulation; What is in a moniker? <i>Neuromodulation</i> . 2009;12(1):1-4. | 5 |
| 13 | Abejon D, Rueda P, Del Saz J, Arango S, Monzon E, Gilsanz F. Is the introduction of another variable to the strength-duration curve necessary in neurostimulation? <i>Neuromodulation</i> . 2015;18(3):182-90. | 5 |
| 14 | Abhinav K, Park ND, Prakash SK, Love-Jones S, Patel NK. Novel use of narrow paddle electrodes for occipital nerve stimulation--technical note. <i>Neuromodulation</i> ;16(6):607-9. | 5 |
| 15 | Abou-Zeid Ahmed H, Davis JRE, Kearney T, Gnanalingham KK. Transient asystole during endoscopic transsphenoidal surgery for Acromegaly: An example of trigeminocardiac reflex. <i>Pituitary</i> . 2009;12(4):373-4. | 7 |
| 16 | Abraham ME, Potdar A, Ward M, Herschman Y, Mammis A. Risk Factors Associated with Lead Migration Requiring Revision in Dorsal Root Ganglion Stimulation. <i>World Neurosurgery</i> ;128:e649-e52. | 5 |
| 17 | Abraham WT, Zile MR, Weaver FA, Butter C, Ducharme A, Halbach M, et al. Baroreflex Activation Therapy for the Treatment of Heart Failure With a Reduced Ejection Fraction. <i>JACC Heart failure</i> . 2015;3(6):487-96. | 4 |
| 18 | Barwick A, Lanting S, Chuter V. Intra-tester and inter-tester reliability of post-occlusive reactive hyperaemia measurement at the hallux. <i>Microvascular Research</i> . 2015;99:67-71. | 3 |
| 19 | Becker AB, Chen L, Hossack JA, Klibanov AL, French BA. Contrast enhanced ultrasound detects sustained reductions in perfusion and differential patterns of muscle perfusion in surgical mouse models of hindlimb ischemia. <i>Circulation Conference</i> . 2018;138(Supplement 1). | 5 |
| 20 | Benitez E, Sumpio BJ, Chin J, Sumpio BE. Contemporary assessment of foot perfusion in patients with critical limb ischemia. <i>Seminars in Vascular Surgery</i> . 2014;27(1):3-15. | 5 |
| 21 | bhbrp RBR. Changes in oxygenation of skin and muscles after the use of Functional Bandage in women with Varicose Veins. http://www.whoint/trialsearch/Trial2.aspx?TrialID=RBR-9bhbrp . 2018. | 2 |
| 22 | Blaise S, Boulon C, Decamps J, Senet P, Lazareth I, Tribout L, et al. Correlation between capillaroscopic classifications, digital arterial pressure and disease stage in systemic sclerosis: Baseline results from sclerocapstudy. <i>Journal of Scleroderma and Related Disorders</i> . 2018;3 (Supplement 1):25. | 5 |
| 23 | Brownrigg JR, Hinchliffe RJ, Apelqvist J, Boyko EJ, Fitridge R, Mills JL, et al. International Working Group on the Diabetic, Foot. Performance of prognostic markers in the prediction of wound healing or amputation among patients with foot ulcers in diabetes: a systematic review. <i>Diabetes/Metabolism Research Reviews</i> . 2016;32 Suppl 1:128-35. | 4 |
| 24 | Bunte MC, Shishehbor MH. Resolving the high stakes of limb salvage with skin perfusion pressure. <i>Vascular Medicine</i> . 2018;23(3):250-2. | 5 |
| 25 | Campo A, Heuten H, Goovaerts I, Ennekens G, Vrints C, Dirckx J. A non-contact approach for PWV detection: application in a clinical setting. <i>Physiological Measurement</i> . 2016;37(7):990-1003. | 2 |
| 26 | Caolo V, Vries M, Zupancich J, Houben M, Mihov G, Wagenaar A, et al. CXCL1 microspheres: a novel tool to stimulate arteriogenesis. <i>Drug Delivery</i> . 2016;23(8):2919-26. | 6 |
| 27 | Chen M, Fan H, Ledford BT, Farah Z, Barron C, Liu Z, et al. Impacts of femoral artery and vein excision versus femoral artery excision on the hindlimb ischemic model in CD-1 mice. <i>Microvascular Research</i> . 2017;110:48-55. | 6 |
| 28 | Chen Q, Rosenson RS. Systematic Review of Methods Used for the Microvascular Assessment of Peripheral Arterial Disease. <i>Cardiovascular Drugs & Therapy</i> . 2018;32(3):301-10. | 4 |

| 연번 | 서지정보 | 배제 사유 |
|----|--|-------|
| 29 | Chen Y, Kuang X, Zhou J, Zhen P, Zeng Z, Lin Z, et al. Proximal Tibial Cortex Transverse Distraction Facilitating Healing and Limb Salvage in Severe and Recalcitrant Diabetic Foot Ulcers. <i>Clinical orthopaedics and related research</i> . 2019. | 2 |
| 30 | Cooper KJ, Pena C, Benenati J. Determining End Points for Critical Limb Ischemia Interventions. <i>Techniques in Vascular & Interventional Radiology</i> . 2016;19(2):104-12. | 5 |
| 31 | Coppelli A, Abbruzzese L, Goretti C, Iacopi E, Riitano N, Piaggese A. Does Microangiopathy Contribute to the Pathogenesis of the Diabetic Foot Syndrome? <i>Frontiers in Diabetes</i> . 2017;26:70-82. | 5 |
| 32 | De Wert LA, Geerts M, Van Der Brug S, Adriaansen L, Poeze M, Schaper N, et al. The Effect of Shear Force on Skin Viability in Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> . 2019;2019 (1973704). | 1 |
| 33 | Drks. Multi-center, Randomized Controlled Clinical Investigation Evaluating a Unique Advance Wound Care Matrix versus Standard of Care in the Treatment of Diabetic Foot Ulcers. http://www.who.int/trialsearch/Trial2.aspx?TrialID=DRKS00016754 . 2019. | 5 |
| 34 | Dubin A, Henriquez E, Hernandez G. Monitoring peripheral perfusion and microcirculation. <i>Current Opinion in Critical Care</i> . 2018;24(3):173-80. | 5 |
| 35 | Ellis-McConnell W, Taylor A, Kelly P, Meyr AJ. Quantitative Assessment of Peroneal Artery Pressure at the Ankle With Noninvasive Vascular Testing. <i>Journal of Foot & Ankle Surgery</i> . 2017;56(3):551-4. | 1 |
| 36 | England CG, Im HJ, Feng L, Chen F, Graves SA, Hernandez R, et al. Re-assessing the enhanced permeability and retention effect in peripheral arterial disease using radiolabeled long circulating nanoparticles. <i>Biomaterials</i> . 2016;100:101-9. | 6 |
| 37 | Fan X, Li K, Zhu L, Deng X, Feng Z, Xu C, et al. Prolonged therapeutic effects of photoactivated adipose-derived stem cells following ischaemic injury. <i>Acta Physiologica</i> . 2020;230 (1) (e13475). | 6 |
| 38 | Federman DG, Lzdiznski B, Dardik A, Kelly M, Shapshak D, Ueno C, et al. Wound Healing Society 2014 update on guidelines for arterial ulcers. <i>Wound Repair & Regeneration</i> . 2016;24(1):127-35. | 5 |
| 39 | Fedorovich AA, Bagdasarian AG, Uchkin IG, Soboleva GN, Boitsov SA. [Present-day possibilities of non-invasive control over microcirculation and metabolism in man]. <i>Angiologija i Sosudistaia Khirurgiia/Angiology & Vascular Surgery</i> . 2018;24(1):7-18. | 7 |
| 40 | Ferreira CA, Hernandez R, Yang Y, Valdovinos HF, Engle JW, Cai W. ImmunoPET of CD146 in a Murine Hindlimb Ischemia Model. <i>Molecular Pharmaceutics</i> . 2018;15(8):3434-41. | 6 |
| 41 | Forsythe RO, Apelqvist J, Boyko EJ, Fitrige R, Hong JP, Katsanos K, et al. Performance of prognostic markers in the prediction of wound healing or amputation among patients with foot ulcers in diabetes: A systematic review. <i>Diabetes/Metabolism Research Reviews</i> . 2020;36 Suppl 1:e3278. | 4 |
| 42 | Forsythe RO, Hinchliffe RJ. Assessment of foot perfusion in patients with a diabetic foot ulcer. <i>Diabetes/Metabolism Research and Reviews</i> . 2016;32(Supplement 1):232-8. | 5 |
| 43 | Fronek A, Allison M. Noninvasive evaluation of endothelial activity in healthy and diseased individuals. <i>Vascular & Endovascular Surgery</i> . 2014;48(2):134-8. | 2 |
| 44 | Fu J, Zou J, Chen C, Li H, Wang L, Zhou Y. Hydrogen molecules (H2) improve perfusion recovery via antioxidant effects in experimental peripheral arterial disease. <i>Molecular Medicine Reports</i> . 2018;18(6):5009-15. | 6 |
| 45 | Fujihara M, Yazu Y, Takahara M. Intravascular Ultrasound-Guided Interventions for Below-the-Knee Disease in Patients With Chronic Limb-Threatening Ischemia. <i>Journal of Endovascular Therapy</i> . 2020;27(4):565-74. | 2 |
| 46 | Fujii M, Terashi H. Treatment of diabetic foot ulcers with osteomyelitis. <i>Wound Repair and Regeneration</i> . 2019;27 (5):A3. | 5 |
| 47 | Fukunaga M, Kawasaki D, Nishimura M, Yamagami M, Fujiwara R, Nakata T. Clinical Effects of Planned Endovascular Therapy for Critical Limb Ischemia Patients with Tissue Loss. <i>Journal of Atherosclerosis & Thrombosis</i> . 2019;26(3):294-301. | 2 |
| 48 | Funabashi S, Kawarada O, Hirano T, Ayabe S, Yagyu T, Noguchi T, et al. Awareness of scleroderma peripheral artery disease presenting with critical limb ischemia in elderly patients. <i>Circulation Journal</i> . 2019;83(5):1081. | 4 |

| 연번 | 서지정보 | 배제 사유 |
|----|---|-------|
| 49 | Jensen GS, Carter SG, Reeves SG, Robinson LE, Benson KF. Anti-Inflammatory Properties of a Dried Fermentate In Vitro and In Vivo. <i>Journal of medicinal food</i> . 2015;18(3):378-84. | 1 |
| 50 | Hammer A, Koppensteiner R, Steiner S, Niessner A, Goliasch G, Gschwandtner M, et al. Dark chocolate and vascular function in patients with peripheral artery disease: A randomized, controlled cross-over trial. <i>Clinical Hemorheology and Microcirculation</i> . 2015;59(2):145-53. | 2 |
| 51 | Hinchliffe RJ, Brownrigg JRW, Apelqvist J, Boyko EJ, Fitridge R, Mills JL, et al. IWGDF guidance on the diagnosis, prognosis and management of peripheral artery disease in patients with foot ulcers in diabetes. <i>Diabetes/Metabolism Research and Reviews</i> . 2016;32(Supplement 1):37-44. | 5 |
| 52 | Hiratsuka M, Koyama K, Takahashi H, Kasugai T, Hagita J, Kondo A, et al. Low plantar skin perfusion pressure predicts long-term atherosclerotic vascular events and mortality in maintenance haemodialysis patients. <i>Atherosclerosis</i> . 2020;312:66-71. | 1 |
| 53 | Hodges GJ, Nawaz S, Tew GA. Evidence that reduced nitric oxide signal contributes to cutaneous microvascular dysfunction in peripheral arterial disease. <i>Clinical Hemorheology & Microcirculation</i> . 2015;59(1):83-95. | 1 |
| 54 | Hodges G, Tew G, Nawaz S. Evidence that cutaneous microvascular dysfunction in peripheral arterial disease is due to reduced nitric oxide. <i>FASEB Journal Conference: Experimental Biology</i> . 2014;28(1 SUPPL. 1). | 5 |
| 55 | Hoel H, Hoiseth LO, Sandbaek G, Sundhagen JO, Mathiesen I, Hisdal J. The acute effects of different levels of intermittent negative pressure on peripheral circulation in patients with peripheral artery disease. <i>Physiological Reports</i> . 2019;7 (20) (e14241). | 2 |
| 56 | Hoinoiu B, Jiga LP, Nistor A, Dornean V, Barac S, Miclaus G, et al. Chronic hindlimb ischemia assessment; Quantitative evaluation using laser doppler in a rodent model of surgically induced peripheral arterial occlusion. <i>Diagnostics</i> . 2019;9 (4) (39). | 6 |
| 57 | Hong G, Lee JC, Jha A, Diao S, Nakayama KH, Hou L, et al. Near-infrared II fluorescence for imaging hindlimb vessel regeneration with dynamic tissue perfusion measurement. <i>Circulation: Cardiovascular Imaging</i> . 2014;7(3):517-25. | 6 |
| 58 | Hoyer C, Paludan JP, Pavar S, Biurun Manresa JA, Petersen LJ. Reliability of laser Doppler flowmetry curve reading for measurement of toe and ankle pressures: intra- and inter-observer variation. <i>European Journal of Vascular & Endovascular Surgery</i> . 2014;47(3):311-8. | 3 |
| 59 | Ikeoka K, Hoshida S, Watanabe T, Shinoda Y, Minamisaka T, Fukuoka H, et al. Pathophysiological Significance of Velocity-Based Microvascular Resistance at Maximal Hyperemia in Peripheral Artery Disease. <i>Journal of Atherosclerosis & Thrombosis</i> . 2018;25(11):1128-36. | 2 |
| 60 | Ikeoka K, Watanabe T, Shinoda Y, Minamisaka T, Fukuoka H, Inui H, et al. Below-the-Ankle Arrival Time as a Novel Limb Tissue Perfusion Index: Two-dimensional Perfusion Angiography Evaluation. <i>Journal of Endovascular Therapy</i> . 2020;27(2):198-204. | 2 |
| 61 | Ishii T, Takabe S, Yanagawa Y, Ohshima Y, Kagawa Y, Shibata A, et al. Laser Doppler blood flowmeter as a useful instrument for the early detection of lower extremity peripheral arterial disease in hemodialysis patients: an observational study. <i>BMC Nephrology</i> . 2019;20(1):470. | 1 |
| 62 | Iwagami M, Mochida Y, Ishioka K, Oka M, Moriya H, Ohtake T, et al. LDL-apheresis dramatically improves generalized calciphylaxis in a patient undergoing hemodialysis. <i>Clinical Nephrology</i> . 2014;81(3):198-202. | 4 |
| 63 | Jan YK, Liao F, Cheing GLY, Pu F, Ren W, Choi HMC. Differences in skin blood flow oscillations between the plantar and dorsal foot in people with diabetes mellitus and peripheral neuropathy. <i>Microvascular Research</i> . 2019;122:45-51. | 2 |
| 64 | Jprn U. Bone marrow mononuclear cell implantation for patients with critical limb ischemia. http://www.who.int/trialssearch/Trial2.aspx?TrialID=JPRN-UMIN000021422 . 2016. | 5 |
| 65 | Jprn U. Clinical trial to evaluate efficacy and safety of apheresis therapy on peripheral arterial disease with critical limb ischemia. http://www.who.int/trialssearch/Trial2.aspx?TrialID=JPRN-UMIN000020336 . 2015. | 5 |

| 연번 | 서지정보 | 배제 사유 |
|----|---|-------|
| 66 | Jprn U. Effect of belt electrode skeletal muscle electrical stimulation (B-SES) on leg function and ambulatory state during the bed rest period after surgery for diabetic foot patients. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000036558 . 2019. | 2 |
| 67 | Jprn U. Evaluation of tissue blood flow for limb ischemia by finger-mounted tissue oximetry. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000025021 . 2016. | 2 |
| 68 | Jprn U. Ex Vivo Expanded Erythroblast-Transplantation for patient with critical limb ischemia. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000021374 . 2016. | 5 |
| 69 | Jprn U. Far Infrared Therapy on Peripheral Arterial Disease in Hemodialysis Patients. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000016680 . 2015. | 5 |
| 70 | Jprn U. LDLapheresis-mediated Endothelial activation Therapy to Severe-Peripheral Artery Disease study. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000021684 . 2016. | 5 |
| 71 | Jprn U. Revascularization by Peripheral Blood Mononuclear Cell Transplantation. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000029597 . 2017. | 5 |
| 72 | Jprn U. Therapeutic efficacy of magnesium oxide for peripheral artery disease among patients undergoing hemodialysis - Open label randomized controlled trial. http://www.who.int/trialsearch/Trial2.aspx?TrialID=JPRN-UMIN000024275 . 2016. | 5 |
| 73 | Jumar A, Ott C, Kistner I, Friedrich S, Michelson G, Harazny JM, et al. Early Signs of End-Organ Damage in Retinal Arterioles in Patients with Type 2 Diabetes Compared to Hypertensive Patients. <i>Microcirculation (new york, NY : 1994)</i> . 2016;23(6):447-55. | 1 |
| 74 | Kamenskaia OV, Klinkova AS, Karpenko AA, Karas'kov AM, Zeidlits GA. [Peripheral microcirculation in patients with lower-limb atherosclerosis on the background of metabolic syndrome]. <i>Angiologiia i Sosudistaia Khirurgiia/Angiology & Vascular Surgery</i> . 2014;20(4):22-6. | 7 |
| 75 | Katare R, Stroemer P, Hicks C, Stevanato L, Patel S, Corteling R, et al. Clinical-grade human neural stem cells promote reparative neovascularization in mouse models of hindlimb ischemia. <i>Arteriosclerosis, Thrombosis & Vascular Biology</i> . 2014;34(2):408-18. | 6 |
| 76 | Kawarada O, Yasuda S, Nishimura K, Sakamoto S, Noguchi M, Takahi Y, et al. Effect of single tibial artery revascularization on microcirculation in the setting of critical limb ischemia. <i>Circulation: Cardiovascular Interventions</i> . 2014;7(5):684-91. | 2 |
| 77 | Keiji S. Assessment of Circulatory Status of the Newborn in the Neonatal Intensive Care Unit. <i>Neonatal medicine</i> . 2016;23(2):67-73. | 1 |
| 78 | Kikuchi S, Miyake K, Tada Y, Uchida D, Koya A, Saito Y, et al. Laser speckle flowgraphy can also be used to show dynamic changes in the blood flow of the skin of the foot after surgical revascularization. <i>Vascular</i> . 2019;27(3):242-51. | 2 |
| 79 | Kimura T, Watanabe Y, Tokuoka S, Nagashima F, Ebisudani S, Inagawa K. Utility of skin perfusion pressure values with the Society for Vascular Surgery Wound, Ischemia, and foot Infection classification system. <i>Journal of Vascular Surgery</i> . 2019;70(4):1308-17. | 3 |
| 80 | Klonizakis M, Manning G, Lingam K, Donnelly R, Yeung JMC. Effect of diabetes on the cutaneous microcirculation of the feet in patients with intermittent claudication. <i>Clinical Hemorheology and Microcirculation</i> . 2016;61(3):439-44. | 2 |
| 81 | Kolluri R, Jaff MR. Peripheral Artery Disease: What Hospitalists Need to Know for Effective Management. <i>Hospital Medicine Clinics</i> . 2017;6(3):397-412. | 5 |
| 82 | Kovacs D, Csiszar B, Biro K, Koltai K, Endrei D, Juricskay I, et al. Toe-brachial index and exercise test can improve the exploration of peripheral artery disease. <i>Atherosclerosis</i> . 2018;269:151-8. | 3 |
| 83 | Kumar VA, Liu Q, Wickremasinghe NC, Shi S, Cornwright TT, Deng Y, et al. Treatment of hind limb ischemia using angiogenic peptide nanofibers. <i>Biomaterials</i> . 2016;98:113-9. | 6 |
| 84 | Lamova SN. The Place of Nailfold Capillaroscopy Among Instrumental Methods for Assessment of Some Peripheral Ischaemic Syndromes in Rheumatology. <i>Folia Medica (Plovdiv)</i> . 2016;58(2):77-88. | 5 |
| 85 | Laneelle D, Stivalet O, Mahe G, Jaquinandi V, Omarjee L. Exercise Oximetry and Laser Speckle Contrast Imaging to Assess Microvascular Function in Lower Extremity Peripheral Artery Disease. <i>Cardiovascular Drugs & Therapy</i> . 2018;32(4):405-7. | 5 |

| 연번 | 서지정보 | 배제 사유 |
|-----|--|-------|
| 86 | Langston JW, Toombs CF. Defining the minimally effective dose and schedule for parenteral hydrogen sulfide: long-term benefits in a rat model of hindlimb ischemia. <i>Medical Gas Research</i> . 2015;5:5. | 6 |
| 87 | Lanting SM, Twigg SM, Johnson NA, Baker MK, Caterson ID, Chuter VH. Non-invasive lower limb small arterial measures co-segregate strongly with foot complications in people with diabetes. <i>Journal of Diabetes & its Complications</i> . 2017;31(3):589-93. | 2 |
| 88 | Lebas B, Galley J, Renaud-Gabardos E, Pujol F, Lenfant F, Garmy-Susini B, et al. Therapeutic Benefits and Adverse Effects of Combined Proangiogenic Gene Therapy in Mouse Critical Leg Ischemia. <i>Annals of Vascular Surgery</i> . 2017;40:252-61. | 6 |
| 89 | Leung OM, Li J, Li X, Chan VW, Yang KY, Ku M, et al. Regulatory T Cells Promote Apelin-Mediated Sprouting Angiogenesis in Type 2 Diabetes. <i>Cell Reports</i> . 2018;24(6):1610-26. | 6 |
| 90 | Liao F, Jan YK. Nonlinear dynamics of skin blood flow response to mechanical and thermal stresses in the plantar foot of diabetics with peripheral neuropathy. <i>Clinical Hemorheology & Microcirculation</i> . 2017;66(3):197-210. | 1 |
| 91 | Lowry D, Saeed M, Narendran P, Tiwari A. The Difference between the Healing and the Nonhealing Diabetic Foot Ulcer: A Review of the Role of the Microcirculation. <i>Journal of Diabetes Science and Technology</i> . 2017;11(5):914-23. | 5 |
| 92 | Lung CW, Wu FL, Liao F, Pu F, Fan Y, Jan YK. Emerging technologies for the prevention and management of diabetic foot ulcers. <i>Journal of Tissue Viability</i> . 2020;29(2):61-8. | 4 |
| 93 | Ma KF, Kleiss SF, Schuurmann RCL, Bokkers RPH, Unlu C, De Vries J. A systematic review of diagnostic techniques to determine tissue perfusion in patients with peripheral arterial disease. <i>Expert Review of Medical Devices</i> . 2019;16(8):697-710. | 5 |
| 94 | Manu CA, Casey J, Morris V, Vas P, Rashid H, Edmonds ME. Better tissue oxygenation in patients with peripheral arterial disease post bypass surgery compared to angioplasty only: Detailed assessment of microvascular and macrovascular circulation. <i>Diabetic Medicine</i> . 2015;1):150. | 5 |
| 95 | Manu CA, Morris V, Vas P, Simmggen M, Edmonds ME. Measurement of TcPO ₂ , toe pressures and laser Doppler flow reveals hitherto undiagnosed subclinical ischaemia in patients with palpable pedal pulses and foot ulceration. <i>Diabetologia</i> . 2014;1):S471. | 5 |
| 96 | Matsui A, Murakami M, Hata S, Terabe Y, Nakabayashi K, Kaneko N, et al. The Efficacy of the Transdermal Isosorbide Dinitrate Patch in Patients With Chronic Limb-Threatening Ischemia. <i>International Journal of Lower Extremity Wounds</i> . 2020;1534734620959515. | 2 |
| 97 | McCormick KG, Scorletti ES, Bhatia L, Calder PC, Griffin MJ, Clough GF, et al. Peripheral sensory nerve function is independently associated with microvascular function, but neither are improved by n-3 fatty acids. <i>Diabetic medicine</i> . 2015;32:99. | 5 |
| 98 | Meisner JK, Annex BH, Price RJ. Despite normal arteriogenic and angiogenic responses, hind limb perfusion recovery and necrotic and fibroadipose tissue clearance are impaired in matrix metalloproteinase 9-deficient mice. <i>Journal of Vascular Surgery</i> . 2015;61(6):1583-94.e10. | 6 |
| 99 | Midttun M, Azad BB, Broholm R, Jensen LT, Svarer C, Jensen PE. Heat-washout measurements compared to distal blood pressure and perfusion in orthopaedic patients with foot ulcers. <i>Clinical Physiology & Functional Imaging</i> . 2017;37(1):79-83. | 2 |
| 100 | Mills JL. Lower limb ischaemia in patients with diabetic foot ulcers and gangrene: recognition, anatomic patterns and revascularization strategies. <i>Diabetes/Metabolism Research Reviews</i> . 2016;32 Suppl 1:239-45. | 5 |
| 101 | Ministro A, De Oliveira P, Nunes RJ, Dos Santos Rocha A, Correia A, Carvalho T, et al. Low-dose ionizing radiation induces therapeutic neovascularization in a pre-clinical model of hindlimb ischemia. <i>Cardiovascular Research</i> . 2017;113(7):783-94. | 6 |
| 102 | Misra S, Shishehbor MH, Takahashi EA, Aronow HD, Brewster LP, Bunte MC, et al. Perfusion Assessment in Critical Limb Ischemia: Principles for Understanding and the Development of Evidence and Evaluation of Devices: A Scientific Statement from the American Heart Association. <i>Circulation</i> . 2019;140(12):E657-E72. | 5 |

| 연번 | 서지정보 | 배제 사유 |
|-----|--|-------|
| 103 | Montero M, Najafi B, Hinko V, Hoeglner S, Rahemi H, Enriquez A, et al. Using frailty and cognitive assessment to predict adverse events after major vascular intervention: Application of wearable technologies. <i>Journal of Vascular Surgery</i> . 2017;66 (3):e56. | 5 |
| 104 | Mora-Rodríguez R, Ramirez-Jimenez M, Fernandez-Elias VE, Guio de Prada MV, Morales-Palomo F, Pallares JG, et al. Effects of aerobic interval training on arterial stiffness and microvascular function in patients with metabolic syndrome. <i>Journal of clinical hypertension</i> . 2018;20(1):11-8. | 1 |
| 105 | Morimoto N, Kakudo N, Valentin Notodihardjo P, Suzuki S, Kusumoto K. Comparison of neovascularization in dermal substitutes seeded with autologous fibroblasts or impregnated with bFGF applied to diabetic foot ulcers using laser Doppler imaging. <i>Journal of Artificial Organs</i> . 2014;17(4):352-7. | 2 |
| 106 | Nakada H, Kashiwagi T, Iino Y, Katayama Y. Therapeutic effects of the long-term use of PAN membrane dialyzer in hemodialysis patients: efficacy in old dialysis patients with mild PAD. <i>Journal of Nippon Medical School = Nihon Ika Daigaku Zasshi</i> . 2014;81(4):221-35. | 1 |
| 107 | Nakamura H, Makiguchi T, Atomura D, Yamatsu Y, Shirabe K, Yokoo S. Changes in Skin Perfusion Pressure After Hyperbaric Oxygen Therapy Following Revascularization in Patients With Critical Limb Ischemia: A Preliminary Study. <i>International Journal of Lower Extremity Wounds</i> . 2020;19(1):57-62. | 2 |
| 108 | Nakanishi T, Tsujii M, Lino T, Sudo A. Analysis of skeletal muscles during regeneration due to reduced circulation in ob/ob mice and effect of edaravone for oxidative stress on C2C12 myoblast. <i>Journal of Orthopaedic Research Conference</i> . 2016;34(Supplement 1). | 5 |
| 109 | Nazer B, Ghahghaie F, Kashima R, Khokhlova T, Perez C, Crum L, et al. Therapeutic ultrasound promotes reperfusion and angiogenesis in a rat model of peripheral arterial disease. <i>Circulation Journal</i> . 2015;79(9):2043-9. | 6 |
| 110 | Nct. Acute Effects of Leg Heating on Skeletal Muscle Blood Flow. https://clinicaltrials.gov/show/NCT03763357 . 2018. | 2 |
| 111 | Nct. Adjunctive Hyperbaric Oxygen Therapy (HBOT) for Lower Extremity Diabetic Ulcer. https://clinicaltrials.gov/show/NCT03675269 . 2018. | 2 |
| 112 | Nct. AVAZZIA Electrical Stimulation. https://clinicaltrials.gov/show/NCT03821675 . 2019. | 2 |
| 113 | Nct. Comparative Study Evaluating Performance of Celliant Fibers on Tissue Oxygenation. https://clinicaltrials.gov/show/NCT02798640 . 2016. | 2 |
| 114 | Nct. Comparison of Prevena Negative Pressure Incision Management System vs. Standard Dressing After Vascular Surgery. https://clinicaltrials.gov/show/NCT02389023 . 2015. | 5 |
| 115 | Nct. EO2 Oxygen Delivery To Study Success Rate of Surgically Closed Wounds. https://clinicaltrials.gov/show/NCT03960463 . 2019. | 5 |
| 116 | Nct. Iontophoresis of Treprostinil to Enhance Wound Healing in Diabetic Foot Skin Ulcers. https://clinicaltrials.gov/show/NCT03654989 . 2018. | 2 |
| 117 | Nct. RCT to Compare Restrata and NPWT to Heal Complex DFUs. https://clinicaltrials.gov/show/NCT04405050 . 2020. | 2 |
| 118 | Nct. Safety and Efficacy of Magnesium Sulphate as an Adjuvant to Levobupivacaine in Ultra-sound Guided Transversus Abdominis Plan Block in Pediatric Abdominal Cancer Surgery. https://clinicaltrials.gov/show/NCT03979599 . 2019. | 1 |
| 119 | Nct. The Prognostic Impact of Using High-dose Hydralazine in Severe Systolic Heart Failure With Hemodynamically Significant Mitral Regurgitation. https://clinicaltrials.gov/show/NCT04217135 . 2020. | 1 |
| 120 | Nishimoto S, Aini K, Fukuda D, Higashikuni Y, Tanaka K, Hirata Y, et al. Activation of Toll-Like Receptor 9 Impairs Blood Flow Recovery After Hind-Limb Ischemia. <i>Frontiers in Cardiovascular Medicine</i> . 2018;5 (144). | 6 |
| 121 | Nishimoto S, Kawai K, Sotsuka Y, Fujita K, Fujiwara T, Tonooka M, et al. Should aggressive minor amputation be performed to treat ischemic gangrene? <i>Wound Repair and Regeneration</i> . 2014;22 (1):A20. | 5 |

| 연번 | 서지정보 | 배제 사유 |
|-----|--|-------|
| 122 | Nogami K, Niihara H, Nakatani T, Kishimoto K, Nakamura M, Matsuki S, et al. A case of Buerger's disease effectively treated by sympathetic block using clipping. [Japanese]. Nishinohon Journal of Dermatology. 2017;79(1):28-33. | 7 |
| 123 | Ohmine T, Iwasa K, Yamaoka T. Strategy of Revascularization for Critical Limb Ischemia Due to Infragenicular Lesions-Which Should Be Selected Firstly, Bypass Surgery or Endovascular Therapy? Avd. 2015;8(4):275-81. | 2 |
| 124 | Ohtake T, Mochida Y, Ishioka K, Oka M, Maesato K, Moriya H, et al. Autologous Granulocyte Colony-Stimulating Factor-Mobilized Peripheral Blood CD34 Positive Cell Transplantation for Hemodialysis Patients with Critical Limb Ischemia: A Prospective Phase II Clinical Trial. Stem Cells Translational Medicine. 2018;7(11):774-82. | 2 |
| 125 | Ohtake T, Mochida Y, Ishioka K, Oka M, Maesato K, Moriya H, et al. Effect of autologous G-CSF-mobilized cd34+ cell transplantation in hemodialysis patients with critical limb ischemia. Nephrology Dialysis Transplantation. 2017;32 (Supplement 3):iii309. | 5 |
| 126 | Ohtake T, Sato M, Nakazawa R, Kondoh M, Miyaji T, Moriya H, et al. Randomized pilot trial between prostaglandin I2 analog and anti-platelet drugs on peripheral arterial disease in hemodialysis patients. Therapeutic Apheresis & Dialysis: Official Peer-Reviewed Journal of the International Society for Apheresis, the Japanese Society for Apheresis, the Japanese Society for Dialysis Therapy. 2014;18(1):1-8. | 2 |
| 127 | Omer SM, Krishna SM, Li J, Moxon JV, Nsengiyumva V, Golledge J. The efficacy of extraembryonic stem cells in improving blood flow within animal models of lower limb ischaemia. Heart. 2016;102(1):69-74. | 4 |
| 128 | Orbay H, Hong H, Koch JM, Valdovinos HF, Hacker TA, Theuer CP, et al. Pravastatin stimulates angiogenesis in a murine hindlimb ischemia model: A positron emission tomography imaging study with 64Cu-NOTA-TRC105. American Journal of Translational Research. 2014;6(1):54-63. | 6 |
| 129 | Osawa Y, Ogawa A, Shimotori T. Ankle-brachial index and skin perfusion pressure : Which is the stronger predictor of life prognosis among maintenance-phase hemodialysis patients? Nephrology Dialysis Transplantation. 2017;32 (Supplement 3):iii320. | 5 |
| 130 | Pek SL, Tavintharan S, Wang X, Lim SC, Woon K, Yeoh LY, et al. Elevation of a novel angiogenic factor, leucine-rich-alpha2-glycoprotein (LRG1), is associated with arterial stiffness, endothelial dysfunction, and peripheral arterial disease in patients with type 2 diabetes. Journal of Clinical Endocrinology & Metabolism. 2015;100(4):1586-93. | 2 |
| 131 | Pek SL, Tavintharan S, Wang X, Lim SC, Woon K, Yeoh LY, et al. Elevation of a novel angiogenic factor, leucine-rich-alpha2-glycoprotein (LRG1), is associated with arterial stiffness, endothelial dysfunction, and peripheral arterial disease in patients with type 2 diabetes. Journal of Clinical Endocrinology and Metabolism. 2015;100(4):1586-93. | 2 |
| 132 | Pek SL, Lim SC, Ang K, Kwan PY, Tang WE, Sum CF, et al. Endothelin-1 predicts incident diabetic peripheral neuropathy in Type 2 Diabetes: a cohort study. European Journal of Endocrinology. 2020;182(4):429-38. | 2 |
| 133 | Pelletier CC, Jolivot A, Kalbacher E, Panaye M, Du Colombier PB, Juillard L. Effects of perodialytic cycling on the peripheral microcirculation in chronic hemodialysis patients: Preliminary results of activdial study. Nephrology Dialysis Transplantation. 2014;3(3):iii489. | 5 |
| 134 | Peravali R, Gunnels L, Alleboina S, Gerling IC, Dokun AO. Type 1 diabetes alters ischemia-induced gene expression. Journal of Clinical and Translational Endocrinology. 2019;15:19-24. | 6 |
| 135 | Pettersen EM, Avdal J, Hisdal J, Torp H, Seternes A. Validation of a novel ultrasound Doppler monitoring device (earlybird) for detection of microvascular circulatory changes. Clinical Hemorheology & Microcirculation. 2020;74(4):429-40. | 1 |
| 136 | Pitts J. Skin perfusion pressure: A case study demonstrating microcirculatory blood flow. Journal of Diagnostic Medical Sonography. 2014;30(4):213-6. | 4 |
| 137 | Poon S, Lu X, Smith RAA, Ho P, Bhakoo K, Nurcombe V, et al. Improved recovery from limb ischaemia by delivery of an affinity-isolated heparan sulphate. Angiogenesis. 2018;21(4):777-91. | 6 |

| 연번 | 서지정보 | 배제 사유 |
|-----|--|-------|
| 138 | Prochazka V, Gumulec J, Jaluvka F, Kusunova P, Jonszta T, Czerny D, et al. No-option critical limb ischemia revascularization with stem cell therapy. <i>Italian Journal of Vascular and Endovascular Surgery</i> . 2014;21(4):191-200. | 4 |
| 139 | Rabin I, Shpolanski U, Leibovitz A, Bass A. Preprogrammed oscillations improve lower limb blood flow and walking distance in patients with peripheral arterial disease. <i>Israel Medical Association Journal: Imaj</i> . 2014;16(7):423-6. | 2 |
| 140 | Rahemi H, Armstrong DG, Enriquez A, Owl J, Talal TK, Najafi B. Lace Up for Healthy Feet: The Impact of Shoe Closure on Plantar Stress Response. <i>Journal of Diabetes Science & Technology</i> . 2017;11(4):678-84. | 1 |
| 141 | Rajendran S, Kevil CG. Cystathionine gamma-lyase regulates autophagy in ischemic endothelial cells. <i>Circulation Conference: Resuscitation Science Symposium, ReSS</i> . 2017;136(Supplement 1). | 5 |
| 142 | Rathsman B, Jensen-Urstad K, Nystrom T. Intensified insulin treatment is associated with improvement in skin microcirculation and ischaemic foot ulcer in patients with type 1 diabetes mellitus: a long-term follow-up study. <i>Diabetologia</i> . 2014;57(8):1703-10. | 2 |
| 143 | Rishi MT, Selvaraju V, Thirunavukkarasu M, Shaikh IA, Takeda K, Fong GH, et al. Deletion of prolyl hydroxylase domain proteins (PHD1, PHD3) stabilizes hypoxia inducible factor-1 alpha, promotes neovascularization, and improves perfusion in a murine model of hind-limb ischemia. <i>Microvascular Research</i> . 2015;97:181-8. | 6 |
| 144 | Rogers RK, Montero-Baker M, Biswas M, Morrison J, Braun J. Assessment of foot perfusion: Overview of modalities, review of evidence, and identification of evidence gaps. <i>Vascular Medicine</i> . 2020;25(3):235-45. | 5 |
| 145 | Rosenson RS, Chen Q, Najera SD, Krishnan P, Lee ML, Cho DJ. Ticagrelor improves blood viscosity-dependent microcirculatory flow in patients with lower extremity arterial disease: the Hema-kinesis clinical trial. <i>Cardiovascular Diabetology</i> . 2019;18(1):77. | 2 |
| 146 | Rosenson RS, Chen Q, Najera SD, Lee ML, Cho DJ. Ticagrelor and the Prevention of Microvascular Complications in Diabetes Patients with Lower Extremity Arterial Disease: Rationale and Design of the Hema-Kinesis Trial. <i>Cardiovascular Drugs & Therapy</i> . 2018;32(5):443-51. | 5 |
| 147 | Rother U, Grussler A, Griesbach C, Almasi-Sperling V, Lang W, Meyer A. Safety of medical compression stockings in patients with diabetes mellitus or peripheral arterial disease. <i>BMJ Open Diabetes Research & Care</i> . 2020;8(1):06. | 2 |
| 148 | Rother U, Krenz K, Lang W, Horch RE, Schmid A, Heinz M, et al. Immediate changes of angiosome perfusion during tibial angioplasty. <i>Journal of Vascular Surgery</i> . 2017;65(2):422-30. | 1 |
| 149 | Rundback JH, Armstrong EJ, Contos B, Iida O, Jacobs D, Jaff MR, et al. Key Concepts in Critical Limb Ischemia: Selected Proceedings from the 2015 Vascular Interventional Advances Meeting. <i>Annals of Vascular Surgery</i> . 2017;38:191-205. | 5 |
| 150 | Ruzsa Z, Rona S, Toth GG, Sotonyi P, Bertrand OF, Nemes B, et al. Fractional flow reserve in below the knee arteries with critical limb ischemia and validation against gold-standard morphologic, functional measures and long term clinical outcomes. <i>Cardiovascular Revascularization Medicine</i> . 2018;19(2):175-81. | 3 |
| 151 | Schmidt CA, Amorese AJ, Ryan TE, Goldberg EJ, Tarpey MD, Green TD, et al. Strain-Dependent Variation in Acute Ischemic Muscle Injury. <i>American Journal of Pathology</i> . 2018;188(5):1246-62. | 6 |
| 152 | Schwarz EI, Schlatzer C, Stehli J, Kaufmann PA, Bloch KE, Stradling JR, et al. Effect of CPAP Withdrawal on myocardial perfusion in OSA: a randomized controlled trial. <i>Respirology (Carlton, Vic)</i> . 2016;21(6):1126-33. | 1 |
| 153 | Seinturier C, Blaise S, Pichot O, Guigard S, Genty C, Jean Luc M, et al. Is contrast enhanced ultrasonography a useful tool for the evaluation of muscular microcirculation in patients with critical limb ischaemia? <i>Vasa</i> . 2017;46(5):389-94. | 2 |
| 154 | Sekiya N, Ichioka S. Efficacy of Ultrasonography at the Ankle Level for Estimation of Pedal Microcirculation. <i>Avd</i> . 2015;8(3):198-202. | 2 |
| 155 | Shemilt R, Khan F. The effects and mechanisms of FAR infrared on the cardiovascular system. <i>Microcirculation (new york, NY : 1994)</i> . 2017;24(7). | 5 |

| 연번 | 서지정보 | 배제 사유 |
|-----|--|-------|
| 156 | Shishehbor MH, Rundback J, Bunte M, Hammad TA, Miller L, Patel PD, et al. SDF-1 plasmid treatment for patients with peripheral artery disease (STOP-PAD): Randomized, double-blind, placebo-controlled clinical trial. <i>Vascular Medicine</i> . 2019;24(3):200-7. | 2 |
| 157 | Shishehbor MH, White CJ, Gray BH, Menard MT, Lookstein R, Rosenfield K, et al. Critical Limb Ischemia: An Expert Statement. <i>Journal of the American College of Cardiology</i> . 2016;68(18):2002-15. | 5 |
| 158 | Shoji K, Yanishi K, Shiraishi H, Yamabata S, Yukawa A, Teramukai S, et al. Establishment of optimal exercise therapy using near-infrared spectroscopy monitoring of tissue muscle oxygenation after therapeutic angiogenesis for patients with critical limb ischemia: A multicenter, randomized, controlled trial. <i>Contemporary Clinical Trials Communications</i> . 2020;17:100542. | 2 |
| 159 | Smith Jorgensen T, Gottlieb H, Brorson S, Hellsten Y, Hoier B. Passive movement training as a treatment for non-healing diabetic foot ulcers: a microcirculatory perspective. <i>Microcirculation (New York, NY : 1994)</i> . 2019;26(4). | 5 |
| 160 | Smith Jorgensen T, Gottlieb H, Brorson S, Hellsten Y, Hoier B. Passive movement training as a treatment for non-healing diabetic foot ulcers: A microcirculatory perspective. <i>Microcirculation Conference: 11th World Congress for Microcirculation, WCM</i> . 2018;26(4). | 5 |
| 161 | Stefanie Reich-Schupke S, Stucker M. Safety of VenoTrain® angioflow in patients with chronic venous insufficiency (CVI) and peripheral arterial occlusive disease (PAD). <i>Vasa - european journal of vascular medicine</i> . 2018;47:9-. | 5 |
| 162 | Stromberg T, Sjoberg F, Bergstrand S. Temporal and spatiotemporal variability in comprehensive forearm skin microcirculation assessment during occlusion protocols. <i>Microvascular Research</i> . 2017;113:50-5. | 1 |
| 163 | Sundby OH, Hoiseth LO, Mathiesen I, Jorgensen JJ, Weedon-Fekjaer H, Hisdal J. Application of intermittent negative pressure on the lower extremity and its effect on macro- and microcirculation in the foot of healthy volunteers. <i>Physiological Reports</i> . 2016;4(17):09. | 1 |
| 164 | Sundby OH, Hoiseth LO, Mathiesen I, Weedon-Fekjaer H, Sundhagen JO, Hisdal J. The acute effects of lower limb intermittent negative pressure on foot macro-and microcirculation in patients with peripheral arterial disease. <i>PLoS ONE</i> . 2017;12 (6) (e0179001). | 2 |
| 165 | Suzuki K, Birnbaum Z, Lockhart R. Skin Perfusion Pressure and Wound Closure Time in Lower Extremity Wounds. <i>The journal of the American College of Clinical Wound Specialists</i> . 2017;9(1-3):14-8. | 3 |
| 166 | Takayama K, Myouchin K, Kichikawa K. Usefulness of laser Doppler tissue flowmeter monitoring during endovascular therapy for peripheral artery disease. <i>CardioVascular and Interventional Radiology</i> . 2016;39 (3 Supplement 1):S304. | 5 |
| 167 | Tara S, Miyamoto M, Takagi G, Kirinoki-Ichikawa S, Tezuka A, Hada T, et al. Low-energy extracorporeal shock wave therapy improves microcirculation blood flow of ischemic limbs in patients with peripheral arterial disease: pilot study. <i>Journal of Nippon Medical School = Nihon Ika Daigaku Zasshi</i> . 2014;81(1):19-27. | 2 |
| 168 | Tsunekawa K, Nagai F, Kato T, Takashimizu I, Yanagisawa D, Yuzuriha S. Hallucal thenar index: A new index to detect peripheral arterial disease using laser speckle flowgraphy. <i>Vascular</i> . 2020;1708538120938935. | 2 |
| 169 | Uchkin IG, Zudin AM, Bagdasarian AG, Fedorovich AA. Effect of drug therapy for chronic obliterating diseases of lower-limb arteries on the state of the microcirculatory bed. <i>Angiologija i sosudistaia khirurgiia = angiology and vascular surgery</i> . 2014;20(2):27-36. | 2 |
| 170 | van den Hoven P, Ooms S, van Manen L, van der Bogt KEA, van Schaik J, Hamming JF, et al. A systematic review of the use of near-infrared fluorescence imaging in patients with peripheral artery disease. <i>Journal of Vascular Surgery</i> . 2019;70(1):286-97.e1. | 4 |
| 171 | Varetto G, Verzini F, Trucco A, Frola E, Spalla F, Gibello L, et al. Oxygen Delivery Therapy with EPIFLO Reduces Wound Hyperperfusion in Patients with Chronic Leg Ulcers: A Laser Speckle Contrast Analysis. <i>Annals of Vascular Surgery</i> . 2020;64:246-52. | 2 |

| 연번 | 서지정보 | 배제 사유 |
|-----|--|-------|
| 172 | Wang CJ, Wu CT, Yang YJ, Liu RT, Kuo YR. Long-term outcomes of extracorporeal shockwave therapy for chronic foot ulcers. <i>Journal of surgical research</i> . 2014;189(2):366-72. | 2 |
| 173 | Wang Z, Hasan R, Firwana B, Elraiyah T, Tsapas A, Prokop L, et al. A systematic review and meta-analysis of tests to predict wound healing in diabetic foot. <i>Journal of Vascular Surgery</i> . 2016;63(2 Suppl):29S-36S.e1-2. | 4 |
| 174 | Weir GR, Hiske S, Van Marle J, Cronje FJ. Arterial disease ulcers, Part 1: Clinical diagnosis and investigation. <i>Journal of the Dermatology Nurses' Association</i> . 2015;7(1):30-7. | 5 |
| 175 | Weir GR, Smart H, van Marle J, Cronje FJ. Arterial disease ulcers, part 1: clinical diagnosis and investigation. <i>Advances in Skin & Wound Care</i> . 2014;27(9):421-8; quiz 9-30. | 5 |
| 176 | Welten SMJ, Bastiaansen AJNM, De Jong RCM, De Vries MR, Peters EAB, Boonstra MC, et al. Inhibition of 14q32 MicroRNAs miR-329, miR-487b, miR-494, and miR-495 increases neovascularization and blood flow recovery after ischemia. <i>Circulation Research</i> . 2014;115(8):696-708. | 6 |
| 177 | Wu FL, Wang WT, Liao F, Elliott J, Jain S, Jan YK. Effects of walking speeds and durations on plantar skin blood flow responses. <i>Microvascular Research</i> . 2020;128:103936. | 2 |
| 178 | Wu FL, Wang WT, Liao F, Liu Y, Li J, Jan YK. Microvascular Control Mechanism of the Plantar Foot in Response to Different Walking Speeds and Durations: Implication for the Prevention of Foot Ulcers. <i>International Journal of Lower Extremity Wounds</i> . 2020:1534734620915360. | 1 |
| 179 | Wu SC, Crews RT, Skratsky M, Overstreet J, Yalla SV, Winder M, et al. Control of lower extremity edema in patients with diabetes: Double blind randomized controlled trial assessing the efficacy of mild compression diabetic socks. <i>Diabetes Research & Clinical Practice</i> . 2017;127:35-43. | 2 |
| 180 | Xu P, Liu X, Song Q, Chen G, Wang D, Zhang H, et al. Patient-specific structural effects on hemodynamics in the ischemic lower limb artery. <i>Scientific Reports</i> . 2016;6:39225. | 2 |
| 181 | Yamamoto K, Hoshina K, Akai A, Isaji T, Akagi D, Miyahara T, et al. Results of surgical interventions for critical limb ischemia due to vasculitis or collagen-tissue related disease. <i>International Angiology</i> . 2017;36(4):332-9. | 2 |
| 182 | Yamamoto Y, Inoue Y, Igari K, Toyofuku T, Kudo T, Uetake H. Assessment of the Severity of Ischaemia and the Outcomes of Revascularisation in Peripheral Arterial Disease Patients Based on the Skin Microcirculatory Response to a Thermal Load Test. <i>Ejves Short Reports</i> . 2019;42:21-5. | 2 |
| 183 | Yata T, Sano M, Kayama T, Naruse E, Yamamoto N, Inuzuka K, et al. Utility of a Finger-Mounted Tissue Oximeter with Near-Infrared Spectroscopy to Evaluate Limb Ischemia in Patients with Peripheral Arterial Disease. <i>Avd</i> . 2019;12(1):36-43. | 2 |
| 184 | Yotsu RR, Pham NM, Oe M, Nagase T, Sanada H, Hara H, et al. Comparison of characteristics and healing course of diabetic foot ulcers by etiological classification: neuropathic, ischemic, and neuro-ischemic type. <i>Journal of Diabetes & its Complications</i> . 2014;28(4):528-35. | 2 |
| 185 | You J, Sun J, Ma T, Yang Z, Wang X, Zhang Z, et al. Curcumin induces therapeutic angiogenesis in a diabetic mouse hindlimb ischemia model via modulating the function of endothelial progenitor cells. <i>Stem Cell Research and Therapy</i> . 2017;8 (1) (182). | 6 |
| 186 | Yu COL, Leung KS, Fung KP, Lam FFY, Ng ESK, Lau KM, et al. The characterization of a full-thickness excision open foot wound model in n5-streptozotocin (STZ)-induced type 2 diabetic rats that mimics diabetic foot ulcer in terms of reduced blood circulation, higher C-reactive protein, elevated inflammation, and reduced cell proliferation. <i>Experimental Animals</i> . 2017;66(3):259-69. | 6 |
| 187 | Yu J, Dardik A. A murine model of hind limb ischemia to study angiogenesis and arteriogenesis. <i>Methods in Molecular Biology</i> . 2018;1717:135-43. | 6 |

| 연번 | 서지정보 | 배제 사유 |
|-----|---|----------|
| 188 | Yu S, Hu SCS, Yu HS, Chin YY, Cheng YC, Lee CH. Early sign of microangiopathy in systemic sclerosis: The significance of cold stress test in dynamic laser Doppler flowmetry. <i>Clinical Hemorheology and Microcirculation</i> . 2019;71(3):373-8. | 4 |
| 189 | Zaleska MT, Olszewski WL, Ross J. The long-term arterial assist intermittent pneumatic compression generating venous flow obstruction is responsible for improvement of arterial flow in ischemic legs. <i>PLoS ONE [Electronic Resource]</i> . 2019;14(12):e0225950. | 2 |
| 190 | Zhang B, Zou Y, Fan Z, Song L, Yang M, Niu G. Effect of catheter-based peripheral sympathetic denervation on peripheral artery sympathetic tone of New Zealand rabbits. [Chinese]. <i>Chinese Journal of Interventional Imaging and Therapy</i> . 2018;15(3):171-4. | 7 |
| 191 | Zhang J, Wang Q, Rao G, Qiu J, He R. Curcumin improves perfusion recovery in experimental peripheral arterial disease by upregulating microRNA-93 expression. <i>Experimental and Therapeutic Medicine</i> . 2019;17(1):798-802. | 6 |
| 192 | Zhang MJ, Sansbury BE, Hellmann J, Baker JF, Guo L, Parmer CM, et al. Resolvin D2 enhances postischemic revascularization while resolving inflammation. <i>Circulation</i> . 2016;134(9):666-80. | 6 |
| 193 | 김석현, 황지현, 모진아, 설아람. 압력 커프를 이용한 레이저 도플러 미세혈류 평가. <i>신의료기술평가 보고서</i> . 2015;1(45):1-77. | 4 |
| 194 | 정찬희, 조윤영, 최덕현, 김보연, 김철희, 목지오. Relationship of Sarcopenia with Microcirculation Measured by Skin Perfusion Pressure in Patients with Type 2 Diabetes. <i>Endocrinology and Metabolism</i> . 2020;35(3):578-86. | 2 |