

**<문헌배제사유>**

1. 광범위한 중화상 환자를 대상으로 하지 않은 연구
2. 자기유래배양피부이식술이 관심중재가 아닌 연구
3. 적절한 의료결과가 하나 이상 보고되지 않은 연구
4. 동물 실험(non-human) 및 전임상시험 연구
5. 원저가 아닌 연구(narrative review, expert opinion, editorial, letter articles, book chapters, comments, proceedings 등)
6. 사전에 정의한 연구유형이 아닌 연구(SR, 증례보고)
7. 한국어나 영어로 출판되지 않은 연구
8. 중복문헌

**1. 국외 DB**

연번	서지정보	배제 사유
1	Bairagi A, Griffin B, Banani T, McPhail SM, Kimble R, Tyack Z. A systematic review and meta-analysis of randomized trials evaluating the efficacy of autologous skin cell suspensions for re-epithelialization of acute partial thickness burn injuries and split-thickness skin graft donor sites. <i>Burns</i> . 2021;16:16.	6
2	Meuli JN, Pantet O, Berger MM, Waselle L, Raffoul W. Massive burns: retrospective analysis of changes in outcomes indicators across 18 years. <i>J Burn Care Res</i> . 2021;22:22.	6
3	Frese L, Darwiche SE, von Rechenberg B, Hoerstrup SP, Giovanoli P, Calcagni M. Thermal conditioning improves quality and speed of keratinocyte sheet production for burn wound treatment. <i>Cytotherapy</i> . 2021;05:05.	4
4	Dearman BL, Greenwood JE. Scale-up of a Composite Cultured Skin using a novel Bioreactor Device in a Porcine Wound Model. <i>J Burn Care Res</i> . 2021;26:26.	4
5	Goh BKL, Chua AWC, Chew KY, Kang GC, Chiang LW, Tan BK, et al. The use of negative-pressure wound therapy over a cultured epithelial autograft for full-thickness wounds secondary to purpura fulminans in an infant. <i>Arch</i> . 2020;20:20.	1
6	Elseth A, Nunez Lopez O. Wound Grafts. StatPearls Publishing. 2020;01:01.	6
7	Sonthalia S, Kachhawa D. Jodhpur Technique. StatPearls Publishing. 2020;01:01.	6
8	Rajput FA, Zeltser R. Aortic Valve Replacement. StatPearls Publishing. 2020;01:01.	6
9	Cooper-Jones B, Visintini S. A Noncultured Autologous Skin Cell Spray Graft for the Treatment of Burns. Canadian Agency for Drugs and Technologies in Health. 2016;168:1-11.	6
10	Blackstone BN, Powell HM. Morphogenesis and Biomechanics of Engineered Skin Cultured Under Uniaxial Strain. <i>Adv Wound Care (New Rochelle)</i> . 2012;1(2):69-74.	4
11	Supp DM. Genomic Reprogramming and Skin-Like Maturation of Engineered Human Skin Substitutes. <i>Adv Wound Care (New Rochelle)</i> . 2012;1(2):63-8.	4
12	Rasmussen CA, Allen-Hoffmann BL. Chimeric Human Skin Substitute Tissue: A Novel Treatment Option for the Delivery of Autologous Keratinocytes. <i>Adv Wound Care (New Rochelle)</i> . 2012;1(2):57-62.	4
13	Koller R, Bierochs B, Meissl G, Rab M, Frey M. The use of allogeneic cultivated keratinocytes for the early coverage of deep dermal burns - indications, results and problems. <i>Cell Tissue Bank</i> . 2002;3(1):11-4.	2
14	Enzo E, Secone Seconetti A, Forcato M, Tenedini E, Polito MP, Sala I, et al. Single-keratinocyte transcriptomic analyses identify different clonal types and proliferative potential mediated by FOXM1 in human epidermal stem cells. <i>Nat Commun</i> . 2021;12(1):2505.	4

연번	서지정보	배제 사유
15	Figueiredo FC, Glanville JM, Arber M, Carr E, Rydevik G, Hogg J, et al. A systematic review of cellular therapies for the treatment of limbal stem cell deficiency affecting one or both eyes. <i>Ocular surf.</i> 2021;20:48-61.	6
16	Anitua E, Pino A, Aspe L, Martinez M, Garcia A, Goni F, et al. Anti-inflammatory effect of different PRGF formulations on cutaneous surface. <i>J Tissue Viability.</i> 2021;30(2):183-9.	6
17	Ozhathil DK, Tay MW, Wolf SE, Branski LK. A Narrative Review of the History of Skin Grafting in Burn Care. <i>Medicina (Kaunas).</i> 2021;57(4):15.	6
18	Selver OB, Gurdal M, Yagci A, Egrilmez S, Palamar M, Cavusoglu T, et al. Multi-parametric evaluation of autologous cultivated Limbal epithelial cell transplantation outcomes of Limbal stem cell deficiency due to chemical burn. <i>BMC ophthalmol.</i> 2020;20(1):325.	1
19	Shen C, Deng H, Sun T, Cai J, Li D, Li L, et al. Use of Fresh Scalp Allografts From Living Relatives for Extensive Deep Burns in Children: A Clinical Study Over 7 Years. <i>J Burn Care Res.</i> 2021;42(2):323-30.	2
20	Wu CJ, Li JJ, Liao WC, Wang TH, Perng CK, Ma H, et al. Using Various Skin Graft Techniques in Major Burn Reconstruction: A Lesson Learned From a Taiwanese Cornstarch Explosion. <i>Ann Plast Surg.</i> 2021;86(2S Suppl 1):S30-S4.	2
21	Ahn HN, Kang HS, Park SJ, Park MH, Chun W, Cho E. Safety and efficacy of basic fibroblast growth factors for deep second-degree burn patients. <i>Burns.</i> 2020;46(8):1857-66.	2
22	Baus A, Chatard M, Bich CS, Trouillas M, Cirodde A, Bey E, et al. [Current place of cultured epithelial autografts in the management of massive burns and future prospects: Literature review]. <i>Ann Chir Plast Esthet.</i> 2021;66(1):10-8.	7
23	Molnar JA, Walker N, Steele TN, Craig C, Williams J, Carter JE, et al. Initial Experience With Autologous Skin Cell Suspension for Treatment of Deep Partial-Thickness Facial Burns. <i>J Burn Care Res.</i> 2020;41(5):1045-51.	2
24	Desanlis A, Albouy M, Rousselle P, Thepot A, Santos MD, Auxenfans C, et al. Validation of an implantable bioink using mechanical extraction of human skin cells: First steps to a 3D bioprinting treatment of deep second degree burn. <i>J Tissue Eng Regen Med.</i> 2021;15(1):37-48.	4
25	Labus W, Kitala D, Klama-Baryla A, Szapski M, Smetek W, Kraut M, et al. A new approach to the production of a biovital skin graft based on human acellular dermal matrix produced in-house, in vitro revitalized internally by human fibroblasts and keratinocytes on the surface. <i>J Biomed Mater Res B Appl Biomater.</i> 2020;108(4):1281-94.	4
26	Monnier S, Abdel-Sayed P, Roessingh AB, Hirt-Burri N, Chemali M, Applegate LA, et al. Surgical Management Evolution Between 2 Massive Burn Cases at 17-Year Interval: Contribution of Cell Therapies in Improving the Surgical Care. <i>Cell Transplant.</i> 2020;29:963689720973642.	6
27	Malara MM, Blackstone BN, Baumann ME, Bailey JK, Supp DM, Powell HM. Cultured Epithelial Autograft Combined with Micropatterned Dermal Template Forms Rete Ridges In Vivo. <i>Tissue Eng Part A.</i> 2020;26(21-22):1138-46.	4
28	Xue DD, Zhou L, Yang Y, Ma SY. Autologous scalp skin grafting to treat toxic epidermal necrolysis in a patient with a large skin injury: A case report. <i>World j.</i> 2021;9(7):1646-53.	6
29	Mogha P, Srivastava A, Kumar S, Das S, Kureel S, Dwivedi A, et al. Hydrogel scaffold with substrate elasticity mimicking physiological-niche promotes proliferation of functional keratinocytes. <i>RSC Adv.</i> 2019;9(18):10174-83.	4

연번	서지정보	배제 사유
30	Fujiwara O, Prasai A, Perez-Bello D, El Ayadi A, Petrov IY, Esenaliev RO, et al. Adipose-derived stem cells improve grafted burn wound healing by promoting wound bed blood flow. <i>Burns Trauma</i> . 2020;8:tkaa009.	4
31	Andreone A, de Hollander D. A case report on the effect of micrografting in the healing of chronic and complex burn wounds. <i>Int</i> . 2020;10(1):15-20.	6
32	Samoila O, Gocan D. Clinical Outcomes From Cultivated Allogenic Stem Cells vs. Oral Mucosa Epithelial Transplants in Total Bilateral Stem Cells Deficiency. <i>Front Med (Lausanne)</i> . 2020;7:43.	6
33	Wang C, Zhang F, Lineaweaver WC. Clinical Applications of Allograft Skin in Burn Care. <i>Ann Plast Surg</i> . 2020;84(3S Suppl 2):S158-S60.	6
34	Tseng SCG, Chen SY, Mead OG, Tighe S. Niche regulation of limbal epithelial stem cells: HC-HA/PTX3 as surrogate matrix niche. <i>Exp Eye Res</i> . 2020;199:108181.	4
35	Ottoman C, Buntrock G, Gatz K, Hartmann B, Aarabi G, Kaschwich M, et al. SkinDot: A modified full-skin transplantation technique. <i>Ann Anat</i> . 2020;229:151454.	6
36	Jalili RB, Pourghadiri A, Li Y, Cleversey C, Kilani RT, Ghahary A. Split Thickness Grafts Grow From Bottom Up in Large Skin Injuries. <i>J Burn Care Res</i> . 2019;40(6):727-33.	4
37	Tjin MS, Chua AWC, Tryggvason K. Chemically defined and xenogeneic-free culture method for human epidermal keratinocytes on laminin-based matrices. <i>Nat</i> . 2020;15(2):694-711.	4
38	Sarkozyova N, Dragunova J, Bukovcan P, Ferancikova N, Breza J, Zilinska Z, et al. Preparation and processing of human allogenic dermal matrix for utilization in reconstructive surgical procedures. <i>Bratisl Lek Listy</i> . 2020;121(6):386-94.	4
39	Magne B, Dedier M, Nivet M, Coulomb B, Banzet S, Lataillade JJ, et al. IL-1beta-Primed Mesenchymal Stromal Cells Improve Epidermal Substitute Engraftment and Wound Healing via Matrix Metalloproteinases and Transforming Growth Factor-beta1. <i>J Invest Dermatol</i> . 2020;140(3):688-98.e21.	4
40	Meuli M, Hartmann-Fritsch F, Hugging M, Marino D, Saglini M, Hynes S, et al. A Cultured Autologous Dermo-epidermal Skin Substitute for Full-Thickness Skin Defects: A Phase I, Open, Prospective Clinical Trial in Children. <i>Plast Reconstr Surg</i> . 2019;144(1):188-98.	2
41	Nanassy AD, Glat PM, Burkey BA, Misra AC, Meyer LK, Gates L, et al. The Use of an Autologous Cell Harvesting and Processing Device to Decrease Surgical Procedures and Expedite Healing in Two Pediatric Burn Patients. <i>Wounds</i> . 2019;31(12):316-21.	6
42	Nanba D. Human keratinocyte stem cells: From cell biology to cell therapy. <i>J Dermatol Sci</i> . 2019;96(2):66-72.	6
43	Li ZQ, Wen HD, Li SL, Niu LB, Wu Q. [Clinical therapeutic observation on sequential treatment of chronic wounds with repeated recurrence of hidradenitis suppurativa in buttocks]. <i>Zhonghua Shao Shang Za Zhi</i> . 2019;35(10):757-60.	7
44	Zhou X, Ning K, Ling B, Chen X, Cheng H, Lu B, et al. Multiple Injections of Autologous Adipose-Derived Stem Cells Accelerate the Burn Wound Healing Process and Promote Blood Vessel Regeneration in a Rat Model. <i>Stem Cells Dev</i> . 2019;28(21):1463-72.	4
45	Borderie VM, Ghoubay D, Georgeon C, Borderie M, de Sousa C, Legendre A, et al. Long-Term Results of Cultured Limbal Stem Cell Versus Limbal Tissue Transplantation in Stage III Limbal Deficiency. <i>Stem cells transl</i> . 2019;8(12):1230-41.	1
46	Lonnqvist S, Junker JPE, Sedell M, Nyman E, Kratz G. Tracking keratinocytes and melanocytes using carboxyfluorescein hydroxysuccinimidyl ester staining. <i>PLoS ONE</i> . 2019;14(8):e0221878.	4
47	Lo CH, Chong E, Akbarzadeh S, Brown WA, Cleland H. A systematic review: Current trends and take rates of cultured epithelial autografts in the treatment of patients with burn injuries. <i>Wound Repair Regen</i> . 2019;27(6):693-701.	6

연번	서지정보	배제 사유
48	Cortez Ghio S, Le-Bel G, Lavoie A, Larouche D, Germain L. Isolation and Culture of Human Keratinocytes. <i>Methods Mol Biol.</i> 2019;1993:3-13.	5
49	Alexaline MM, Magne B, Zuleta Rodriguez A, Nivet M, Bacqueville D, Lataillade JJ, et al. Influence of fibrin matrices and their released factors on epidermal substitute phenotype and engraftment. <i>J Tissue Eng Regen Med.</i> 2019;13(8):1362-74.	4
50	Schlottmann F, Strauss S, Hake K, Vogt PM, Bucan V. Down-Regulation of MHC Class I Expression in Human Keratinocytes Using Viral Vectors Containing US11 Gene of Human Cytomegalovirus and Cultivation on Bovine Collagen-Elastin Matrix (Matriderm <sup>&lt;sup&gt; R&lt;/sup&gt;</sup> ): Potential Approach for an Immune-Privileged Skin Substitute. <i>Int.</i> 2019;20(9):26.	4
51	Skog M, Sivler P, Steinvall I, Aili D, Sjoberg F, Elmasry M. The Effect of Enzymatic Digestion on Cultured Epithelial Autografts. <i>Cell Transplant.</i> 2019;28(5):638-44.	4
52	Ter Horst B, Moakes RJA, Chouhan G, Williams RL, Moiemmen NS, Grover LM. A gellan-based fluid gel carrier to enhance topical spray delivery. <i>Acta Biomater.</i> 2019;89:166-79.	4
53	Fan YF, Chen C, Li JL, Huang N, Cui SY. [Five patients with severe burns complicated by fungal infection]. <i>Zhonghua Shao Shang Za Zhi.</i> 2019;35(3):221-3.	7
54	Kitala D, Klama-Baryla A, Labus W, Ples M, Misiuga M, Kraut M, et al. Amniotic cells share clusters of differentiation of fibroblasts and keratinocytes, influencing their ability to proliferate and aid in wound healing while impairing their angiogenesis capability. <i>Eur J Pharmacol.</i> 2019;854:167-78.	4
55	Cheshire P, Zhafira AS, Banakh I, Rahman MM, Carmichael I, Herson M, et al. Xeno-free expansion of adult keratinocytes for clinical application: the use of human-derived feeder cells and serum. <i>Cell Tissue Res.</i> 2019;376(3):389-400.	4
56	Holmes JHt, Molnar JA, Shupp JW, Hickerson WL, King BT, Foster KN, et al. Demonstration of the safety and effectiveness of the RECELL <sup>&lt;sup&gt; R&lt;/sup&gt;</sup> System combined with split-thickness meshed autografts for the reduction of donor skin to treat mixed-depth burn injuries. <i>Burns.</i> 2019;45(4):772-82.	2
57	Lo CH, Akbarzadeh S, McLean C, Ives A, Paul E, Brown WA, et al. Wound healing after cultured epithelial autografting in patients with massive burn injury: A cohort study. <i>J Plast Reconstr Aesthet Surg.</i> 2019;72(3):427-37.	6
58	Kitala D, Klama-Baryla A, Kraut M, Lowicki D, Labus W, Glik J, et al. Does the transplantation of keratinocytes really reduce the risk of death? Survival analysis of patients hospitalized at the Dr Stanislaw Sakiel Centre for Burns Treatment in 2008-2015. <i>Adv.</i> 2019;28(1):5-10.	2
59	Hayashi M, Yoshitake K, Tokunaka R, Yoshida Y, Oshima M, Tatsuta S, et al. Combination of meshed dermis graft and cultured epithelial autograft for massive burns: Three case reports. <i>Medicine (Baltimore).</i> 2018;97(48):e13313.	6
60	Amini-Nik S, Dolp R, Eylert G, Datu AK, Parousis A, Blakeley C, et al. Stem cells derived from burned skin - The future of burn care. <i>EBioMedicine.</i> 2018;37:509-20.	4
61	Qi YJ, Wang X, Jiao Y, Gong HM, Pan Y, Jiang DY. [Preparation of bioactive denatured acellular dermal matrix from burn mice riched in mice bone marrow mesenchymal stem cells]. <i>Zhonghua Shao Shang Za Zhi.</i> 2018;34(12):895-900.	7
62	Blackstone BN, Hahn JM, McFarland KL, DeBruler DM, Supp DM, Powell HM. Inflammatory response and biomechanical properties of coaxial scaffolds for engineered skin in vitro and post-grafting. <i>Acta Biomater.</i> 2018;80:247-57.	4
63	Klama-Baryla A, Kitala D, Labus W, Kraut M, Glik J, Nowak M, et al. Autologous and Allogeneic Skin Cell Grafts in the Treatment of Severely Burned Patients: Retrospective Clinical Study. <i>Transplant Proc.</i> 2018;50(7):2179-87.	2

연번	서지정보	배제 사유
64	Hassanzadeh H, Matin MM, Naderi-Meshkin H, Bidkhori HR, Mirahmadi M, Raeesolmohaddeseen M, et al. Using paracrine effects of Ad-MSCs on keratinocyte cultivation and fabrication of epidermal sheets for improving clinical applications. <i>Cell Tissue Bank</i> . 2018;19(4):531-47.	4
65	Bremond-Gignac D, Copin H, Benkhalifa M. Corneal epithelial stem cells for corneal injury. <i>Expert Opin Biol Ther</i> . 2018;18(9):997-1003.	6
66	Kruse CR, Sakthivel D, Sinha I, Helm D, Sorensen JA, Eriksson E, et al. Evaluation of the efficacy of cell and micrograft transplantation for full-thickness wound healing. <i>J Surg Res</i> . 2018;227:35-43.	4
67	Efanov JI, Tchiloemba B, Duong A, Belisle A, Izadpanah A, Coeugnet E, et al. Use of bilaminar grafts as life-saving interventions for severe burns: A single-center experience. <i>Burns</i> . 2018;44(5):1336-45.	2
68	Xu X, Lai L, Zhang X, Chen J, Chen J, Wang F, et al. Autologous chyle fat grafting for the treatment of hypertrophic scars and scar-related conditions. <i>Stem Cell Res Ther</i> . 2018;9(1):64.	2
69	Chua AWC, Khoo YC, Truong TTH, Woo E, Tan BK, Chong SJ. From skin allograft coverage to allograft-micrograft sandwich method: A retrospective review of severe burn patients who received conjunctive application of cultured epithelial autografts. <i>Burns</i> . 2018;44(5):1302-7.	2
70	Basu S, Mohan S, Bhalekar S, Singh V, Sangwan V. Simple limbal epithelial transplantation (SLET) in failed cultivated limbal epithelial transplantation (CLET) for unilateral chronic ocular burns. <i>Br J Ophthalmol</i> . 2018;102(12):1640-5.	1
71	Fan SM, Chen PH, Tseng YT, Hong JB, Chen W, Tsai TF, et al. Preclinical evaluation of melanocyte transplantation by chitosan-based melanocyte spheroid patch to skin prepared by controlled sunburn blistering. <i>J Biomed Mater Res B Appl Biomater</i> . 2018;106(7):2535-43.	4
72	Chiba T, Ishida N, Kohda F, Furue M. Air exposure may be associated with the histological differentiation of a cultured epidermal autograft (JACE). <i>Australas J Dermatol</i> . 2018;59(3):e244-e6.	5
73	Esteban-Vives R, Corcos A, Choi MS, Young MT, Over P, Ziembicki J, et al. Cell-spray auto-grafting technology for deep partial-thickness burns: Problems and solutions during clinical implementation. <i>Burns</i> . 2018;44(3):549-59.	2
74	Gholipourmalekabadi M, Samadikuchaksaraei A, Seifalian AM, Urbanska AM, Ghanbarian H, Hardy JG, et al. Silk fibroin/amniotic membrane 3D bi-layered artificial skin. <i>Biomed</i> . 2018;13(3):035003.	4
75	Watt SM, Pleat JM. Stem cells, niches and scaffolds: Applications to burns and wound care. <i>Adv Drug Deliv Rev</i> . 2018;123:82-106.	2
76	Ter Horst B, Chouhan G, Moiemens NS, Grover LM. Advances in keratinocyte delivery in burn wound care. <i>Adv Drug Deliv Rev</i> . 2018;123:18-32.	5
77	Labus W, Glik J, Klama-Baryla A, Kitala D, Kraut M, Maj M, et al. Atomic force microscopy in the production of a biovital skin graft based on human acellular dermal matrix produced in-house and in vitro cultured human fibroblasts. <i>J Biomed Mater Res B Appl Biomater</i> . 2018;106(2):726-33.	4
78	Jeschke MG, Shahrokhi S, Finnerty CC, Branski LK, Dibildox M, Organization ABA, et al. Wound Coverage Technologies in Burn Care: Established Techniques. <i>J Burn Care Res</i> . 2018;39(3):313-8.	5
79	Nasoori A, Hoomand R. Maggot debridement therapy for an electrical burn injury with instructions for the use of <i>Lucilia sericata</i> larvae. <i>J Wound Care</i> . 2017;26(12):734-41.	6

연번	서지정보	배제 사유
80	Lykov AP, Bondarenko NA, Poveshchenko OV, Miller TV, Poveshchenko AF, Surovtseva MA, et al. Prospect of Using Cell Product for the Therapy of Skin Defects in Diabetes Mellitus. <i>Bull Exp Biol Med.</i> 2017;164(2):266-8.	4
81	Rettinger CL, Fletcher JL, Carlsson AH, Chan RK. Accelerated epithelialization and improved wound healing metrics in porcine full-thickness wounds transplanted with full-thickness skin micrografts. <i>Wound Repair Regen.</i> 2017;25(5):816-27.	4
82	Lataillade JJ, Magne B, Bey E, Leclerc T, Trouillas M. [Skin engineering for severe burns]. <i>Transfus Clin Biol.</i> 2017;24(3):245-50.	7
83	Velasquillo C, Silva-Bermudez P, Vazquez N, Martinez A, Espadin A, Garcia-Lopez J, et al. In vitro and in vivo assessment of lactic acid-modified chitosan scaffolds for potential treatment of full-thickness burns. <i>J Biomed Mater Res A.</i> 2017;105(10):2875-91.	4
84	Greenwood JE. The evolution of acute burn care - retiring the split skin graft. <i>Ann R Coll Surg Engl.</i> 2017;99(6):432-8.	5
85	Suzuki D, Pinto F, Senoo M. Inhibition of TGF-beta signaling promotes expansion of human epidermal keratinocytes in feeder cell co-culture. <i>Wound Repair Regen.</i> 2017;25(3):526-31.	4
86	Zieger MA, Ochoa M, Rahimi R, Campana G, Tholpady S, Ziaie B, et al. Skin Regeneration Using Dermal Substrates that Contain Autologous Cells and Silver Nanoparticles to Promote Antibacterial Activity: In Vitro Studies. <i>Mil Med.</i> 2017;182(S1):376-82.	4
87	Yoon C, Lee J, Jeong H, Lee S, Sohn T, Chung S. Rapid preparation of a noncultured skin cell suspension that promotes wound healing. <i>Cell Tissue Bank.</i> 2017;18(2):131-41.	2
88	Sakamoto M, Morimoto N, Inoie M, Takahagi M, Ogino S, Jinno C, et al. Cultured Human Epidermis Combined With Meshed Skin Autografts Accelerates Epithelialization and Granulation Tissue Formation in a Rat Model. <i>Ann Plast Surg.</i> 2017;78(6):651-8.	4
89	Martin YH, Jubin K, Smalley S, Wong JPF, Brown RA, Metcalfe AD. A novel system for expansion and delivery of human keratinocytes for the treatment of severe cutaneous injuries using microcarriers and compressed collagen. <i>J Tissue Eng Regen Med.</i> 2017;11(11):3124-33.	4
90	Beaudoin Cloutier C, Goyer B, Perron C, Guignard R, Larouche D, Moulin VJ, et al. In Vivo Evaluation and Imaging of a Bilayered Self-Assembled Skin Substitute Using a Decellularized Dermal Matrix Grafted on Mice. <i>Tissue Eng Part A.</i> 2017;23(7-8):313-22.	4
91	Fasolo A, Pedrotti E, Passilongo M, Marchini G, Monterosso C, Zampini R, et al. Safety outcomes and long-term effectiveness of ex vivo autologous cultured limbal epithelial transplantation for limbal stem cell deficiency. <i>Br J Ophthalmol.</i> 2017;101(5):640-9.	1
92	Johnstone P, Kwei JS, Filobbos G, Lewis D, Jeffery S. Successful application of keratinocyte suspension using autologous fibrin spray. <i>Burns.</i> 2017;43(3):e27-e30.	6
93	Chen B, Xu M, Li B. The clinical experience for treating post-burn depigmentation with tiny epidermal particles graft. <i>Int Wound J.</i> 2017;14(1):165-71.	2
94	Nilforoushzadeh MA, Kazemikhoo N, Mokmeli S, Zare S, Dahmardehei M, Vaghar Doost R, et al. An Open-Label Study of Low-Level Laser Therapy Followed by Autologous Fibroblast Transplantation for Healing Grade 3 Burn Wounds in Diabetic Patients. <i>J.</i> 2019;10(Suppl 1):S7-S12.	2
95	Nuri T, Ueda K, Fujimori Y. Ten-year Follow-up After Treating Extended Burn Scar Contracture with an Autologous Cultured Dermal Substitute. <i>Plast.</i> 2018;6(6):e1782.	6
96	Rapp SJ, Schwentker AR, Visscher MO, Van Aalst J, Pan BS. Effects of Autologous Fat and ASCs on Swine Hypertrophic Burn Scars: A Multimodal Quantitative Analysis. <i>Plast.</i> 2017;5(11):e1547.	4

연번	서지정보	배제 사유
97	Bolton L. Can Skin Cells Improve Outcomes for Patients With Deep Burns? <i>Wounds</i> . 2020;32(6):174-7.	5
98	Kljenak A, Tominac Trcin M, Bujic M, Dolenc T, Jevak M, Mrcic G, et al. Fibrin gel as a scaffold for skin substitute - production and clinical experience. <i>Acta Clin</i> . 2016;55(2):279-89.	4
99	Busch KH, Bender R, Walezko N, Aziz H, Altintas MA, Aust MC. [Autologous Skin Cell Transplantation and Medical Needling for Repigmentation of Depigmented Burn Scars on UV-protected and UV-exposed Skin]. <i>Handchir Mikrochir Plast Chir</i> . 2016;48(6):346-53.	7
100	Novosadova EV, Manuilova ES, Arsenyeva EL, Grivennikov IA, Myasoedov NF. Fibroblast-like cells as an effective feeder for the cultivation and derivation of new lines of human induced pluripotent stem cells. <i>Dokl Biochem Biophys</i> . 2016;470(1):353-6.	2
101	Kushnerev E, Shawcross SG, Sothirachagan S, Carley F, Brahma A, Yates JM, et al. Regeneration of Corneal Epithelium With Dental Pulp Stem Cells Using a Contact Lens Delivery System. <i>Invest Ophthalmol Vis Sci</i> . 2016;57(13):5192-9.	1
102	Busin M, Breda C, Bertolin M, Bovone C, Ponzin D, Ferrari S, et al. Corneal Epithelial Stem Cells Repopulate the Donor Area within 1 Year from Limbus Removal for Limbal Autograft. <i>Ophthalmology</i> . 2016;123(12):2481-8.	1
103	Esteban-Vives R, Young MT, Zhu T, Beiriger J, Pekor C, Ziembicki J, et al. Calculations for reproducible autologous skin cell-spray grafting. <i>Burns</i> . 2016;42(8):1756-65.	2
104	Esteban-Vives R, Choi MS, Young MT, Over P, Ziembicki J, Corcos A, et al. Second-degree burns with six etiologies treated with autologous noncultured cell-spray grafting. <i>Burns</i> . 2016;42(7):e99-e106.	6
105	Surrao DC, Boon K, Borys B, Sinha S, Kumar R, Biernaskie J, et al. Large-scale expansion of human skin-derived precursor cells (hSKPs) in stirred suspension bioreactors. <i>Biotechnol Bioeng</i> . 2016;113(12):2725-38.	2
106	Busch KH, Bender R, Walezko N, Aziz H, Altintas MA, Aust MC. Combination of medical needling and non-cultured autologous skin cell transplantation (ReNovaCell) for repigmentation of hypopigmented burn scars. <i>Burns</i> . 2016;42(7):1556-66.	2
107	Matsumura H, Harunari N, Ikeda H. First experience using cultured epidermal autografts in Taiwan for burn victims of the Formosa Fun Coast Water Park explosion, as part of Japanese medical assistance. <i>Burns</i> . 2016;42(3):697-703.	6
108	Vazirani J, Ali MH, Sharma N, Gupta N, Mittal V, Atallah M, et al. Autologous simple limbal epithelial transplantation for unilateral limbal stem cell deficiency: multicentre results. <i>Br J Ophthalmol</i> . 2016;100(10):1416-20.	1
109	Mittal V, Jain R, Mittal R, Vashist U, Narang P. Successful management of severe unilateral chemical burns in children using simple limbal epithelial transplantation (SLET). <i>Br J Ophthalmol</i> . 2016;100(8):1102-8.	1
110	Esteban-Vives R, Young MT, Ziembicki J, Corcos A, Gerlach JC. Effects of wound dressings on cultured primary keratinocytes. <i>Burns</i> . 2016;42(1):81-90.	2
111	Akershoek JJ, Vlig M, Talhout W, Boekema BK, Richters CD, Beelen RH, et al. Cell therapy for full-thickness wounds: are fetal dermal cells a potential source? <i>Cell Tissue Res</i> . 2016;364(1):83-94.	4
112	Caliari-Oliveira C, Yaochite JN, Ramalho LN, Palma PV, Carlos D, Cunha Fde Q, et al. Xenogeneic Mesenchymal Stromal Cells Improve Wound Healing and Modulate the Immune Response in an Extensive Burn Model. <i>Cell Transplant</i> . 2016;25(2):201-15.	4
113	Kadoya K, Amano S, Nishiyama T, Inomata S, Tsunenaga M, Kumagai N, et al. Changes in fibrillin-1 expression, elastin expression and skin surface texture at sites of cultured epithelial autograft transplantation onto wounds from burn scar excision. <i>Int Wound J</i> . 2016;13(5):780-6.	3

연번	서지정보	배제 사유
114	Kadoya K, Amano S, Nishiyama T, Inomata S, Tsunenaga M, Kumagai N, et al. Changes in the expression of epidermal differentiation markers at sites where cultured epithelial autografts were transplanted onto wounds from burn scar excision. <i>Int Wound J.</i> 2016;13(3):412-7.	3
115	McHeik JN, Barrault C, Pedretti N, Garnier J, Juchaux F, Levard G, et al. Foreskin-isolated keratinocytes provide successful extemporaneous autologous paediatric skin grafts. <i>J Tissue Eng Regen Med.</i> 2016;10(3):252-60.	2
116	Moravvej H, Hormozi AK, Hosseini SN, Sorouri R, Mozafari N, Ghazisaidi MR, et al. Comparison of the Application of Allogeneic Fibroblast and Autologous Mesh Grafting With the Conventional Method in the Treatment of Third-Degree Burns. <i>J Burn Care Res.</i> 2016;37(1):e90-5.	2
117	Lei J, Hou C, Duan P, Hao Z, Zhai Y, Meng Y. [Clinical application of modified skin soft tissue expansion in early repair of devastating wound on the head due to electrical burn]. <i>Zhonghua Shao Shang Za Zhi.</i> 2015;31(6):406-9.	7
118	Labus W, Kawecki M, Glik J, Maj M, Kitala D, Misiuga M, et al. Own Experience From The Use Of A Substitute Of An Allogeneic Acellular Dermal Matrix Revitalized With In Vitro Cultured Skin Cells In Clinical Practice. <i>Pol Przeql Chir.</i> 2015;87(10):513-21.	6
119	Chugh RM, Chaturvedi M, Yerneni LK. An evaluation of the choice of feeder cell growth arrest for the production of cultured epidermis. <i>Burns.</i> 2015;41(8):1788-95.	6
120	Dai LG, Fu KY, Hsieh PS, Hung YM, Wang YW, Hsia LC, et al. Evaluation of Wound Healing Efficacy of an Antimicrobial Spray Dressing at Skin Donor Sites. <i>Wounds.</i> 2015;27(8):224-8.	2
121	Dhamodaran K, Subramani M, Jeyabalan N, Ponnalagu M, Chevour P, Shetty R, et al. Characterization of ex vivo cultured limbal, conjunctival, and oral mucosal cells: A comparative study with implications in transplantation medicine. <i>Mol Vis.</i> 2015;21:828-45.	2
122	Mahjour SB, Fu X, Yang X, Fong J, Sefat F, Wang H. Rapid creation of skin substitutes from human skin cells and biomimetic nanofibers for acute full-thickness wound repair. <i>Burns.</i> 2015;41(8):1764-74.	4
123	Holan V, Trosan P, Cejka C, Javorkova E, Zajicova A, Hermankova B, et al. A Comparative Study of the Therapeutic Potential of Mesenchymal Stem Cells and Limbal Epithelial Stem Cells for Ocular Surface Reconstruction. <i>Stem cells transl.</i> 2015;4(9):1052-63.	4
124	Sabeh G, Sabe M, Ishak S, Sweid R, Ayoubi M, Chahal AM. [Sequential grafts of cutaneous cells: First results of a new procedure and review of the literature]. <i>J Med Liban.</i> 2015;63(2):47-58.	7
125	Esteban-Vives R, Young M, Over P, Schmelzer E, Corcos A, Ziembicki J, et al. In vitro keratinocyte expansion for cell transplantation therapy is associated with differentiation and loss of basal layer derived progenitor population. <i>Differentiation.</i> 2015;89(5):137-45.	4
126	Wu SH, Huang SH, Lo YC, Chai CY, Lee SS, Chang KP, et al. Autologous adipose-derived stem cells attenuate muscular atrophy and protect spinal cord ventral horn motor neurons in an animal model of burn injury. <i>Cytherapy.</i> 2015;17(8):1066-75.	4
127	Boekema BK, Breederveld RS. Response to Letter to the Editor: 'Human skin allografts: Storage medium at 4degreeC and viability'. <i>Burns.</i> 2015;41(6):1370.	5
128	Gaucher S, Khaznadar Z, Jarraya M. Human skin allograft: storage medium at 4degreeC and viability. <i>Burns.</i> 2015;41(6):1369-70.	5
129	Hanifi N, Halim AS, Aleas CF, Singh J, Marzuki M, Win TT, et al. Epidermal Regeneration of Cultured Autograft, Allograft, and Xenograft Keratinocytes Transplanted on Full-Thickness Wounds in Rabbits. <i>Exp Clin Transplant.</i> 2015;13(3):273-8.	4



연번	서지정보	배제 사유
130	Pedrotti E, Passilongo M, Fasolo A, Nubile M, Parisi G, Mastropasqua R, et al. In Vivo Confocal Microscopy 1 Year after Autologous Cultured Limbal Stem Cell Grafts. <i>Ophthalmology</i> . 2015;122(8):1660-8.	1
131	Paul M, Kaur P, Herson M, Cheshire P, Cleland H, Akbarzadeh S. Use of Clotted Human Plasma and Aprotinin in Skin Tissue Engineering: A Novel Approach to Engineering Composite Skin on a Porous Scaffold. <i>Tissue Eng Part C Methods</i> . 2015;21(10):1098-104.	4
132	Bobba S, Chow S, Watson S, Di Girolamo N. Clinical outcomes of xeno-free expansion and transplantation of autologous ocular surface epithelial stem cells via contact lens delivery: a prospective case series. <i>Stem Cell Res Ther</i> . 2015;6:23.	1
133	Bardag-Gorce F, Oliva J, Wood A, Hoft R, Pan D, Thropay J, et al. Carrier-free cultured autologous oral mucosa epithelial cell sheet (CAOMECS) for corneal epithelium reconstruction: a histological study. <i>Ocular surf</i> . 2015;13(2):150-63.	4
134	Alexaline MM, Trouillas M, Nivet M, Bourreau E, Leclerc T, Duhamel P, et al. Bioengineering a human plasma-based epidermal substitute with efficient grafting capacity and high content in clonogenic cells. <i>Stem cells transl</i> . 2015;4(6):643-54.	4
135	Sood R, Roggy DE, Zieger MJ, Nazim M, Hartman BC, Gibbs JT. A comparative study of spray keratinocytes and autologous meshed split-thickness skin graft in the treatment of acute burn injuries. <i>Wounds</i> . 2015;27(2):31-40.	2
136	Boekema BK, Boekestijn B, Breederveld RS. Evaluation of saline, RPMI and DMEM/F12 for storage of split-thickness skin grafts. <i>Burns</i> . 2015;41(4):848-52.	2
137	Sati A, Basu S, Sangwan VS, Vemuganti GK. Correlation between the histological features of corneal surface pannus following ocular surface burns and the final outcome of cultivated limbal epithelial transplantation. <i>Br J Ophthalmol</i> . 2015;99(4):477-81.	1
138	Clover AJ, Kumar AH, Isakson M, Whelan D, Stocca A, Gleeson BM, et al. Allogeneic mesenchymal stem cells, but not culture modified monocytes, improve burn wound healing. <i>Burns</i> . 2015;41(3):548-57.	4
139	McHeik JN, Barrault C, Pedretti N, Garnier J, Juchaux F, Levard G, et al. Study of proliferation and 3D epidermal reconstruction from foreskin, auricular and trunk keratinocytes in children. <i>Burns</i> . 2015;41(2):352-8.	4
140	Boyce ST, Zimmerman RL, Supp DM. Tumorigenicity Testing in Athymic Mice of Cultured Human Melanocytes for Transplantation in Engineered Skin Substitutes. <i>Cell Transplant</i> . 2015;24(8):1423-9.	4
141	Keskar V, Jamale T, Karegar M, Parikh H, Jawale S, Mahajan D, et al. Late-Onset Cytomegalovirus Gastritis in Low-Risk Renal Allograft Recipients. <i>Exp Clin Transplant</i> . 2015;13(3):279-82.	4
142	Auxenfans C, Menet V, Catherine Z, Shipkov H, Lacroix P, Bertin-Maghit M, et al. Cultured autologous keratinocytes in the treatment of large and deep burns: a retrospective study over 15 years. <i>Burns</i> . 2015;41(1):71-9.	2
143	Nuutila K, Peura M, Suomela S, Hukkanen M, Siltanen A, Harjula A, et al. Recombinant human collagen III gel for transplantation of autologous skin cells in porcine full-thickness wounds. <i>J Tissue Eng Regen Med</i> . 2015;9(12):1386-93.	4
144	Golinski P, Menke H, Hofmann M, Valesky E, Butting M, Kippenberger S, et al. Development and Characterization of an Engraftable Tissue-Cultured Skin Autograft: Alternative Treatment for Severe Electrical Injuries. <i>Cells Tissues Organs</i> . 2014;200(3-4):227-39.	6
145	Takami Y, Yamaguchi R, Ono S, Hyakusoku H. Clinical application and histological properties of autologous tissue-engineered skin equivalents using an acellular dermal matrix. <i>J Nippon Med Sch</i> . 2014;81(6):356-63.	2

연번	서지정보	배제 사유
146	Ye Z, Xie W, Long Z, Wang H, Liu S, Xie Q, et al. [Analysis of diagnosis and management of 21 patients with Marjolin's ulcers]. <i>Zhonghua Shao Shang Za Zhi</i> . 2014;30(6):491-4.	7
147	Lenihan C, Rogers C, Metcalfe AD, Martin YH. The effect of isolation and culture methods on epithelial stem cell populations and their progeny-toward an improved cell expansion protocol for clinical application. <i>Cytotherapy</i> . 2014;16(12):1750-9.	4
148	Amescua G, Atallah M, Nikpoor N, Galor A, Perez VL. Modified simple limbal epithelial transplantation using cryopreserved amniotic membrane for unilateral limbal stem cell deficiency. <i>Am J Ophthalmol</i> . 2014;158(3):469-75.e2.	1
149	Marino D, Reichmann E, Meuli M. Skingeneering. <i>Eur J Pediatr Surg</i> . 2014;24(3):205-13.	6
150	Dearman BL, Li A, Greenwood JE. Optimization of a polyurethane dermal matrix and experience with a polymer-based cultured composite skin. <i>J Burn Care Res</i> . 2014;35(5):437-48.	4
151	Idrus RB, Rameli MA, Low KC, Law JX, Chua KH, Latiff MB, et al. Full-thickness skin wound healing using autologous keratinocytes and dermal fibroblasts with fibrin: bilayered versus single-layered substitute. <i>Adv Skin Wound Care</i> . 2014;27(4):171-80.	4
152	Shahrokhi S, Arno A, Jeschke MG. The use of dermal substitutes in burn surgery: acute phase. <i>Wound Repair Regen</i> . 2014;22(1):14-22.	5
153	Sander EA, Lynch KA, Boyce ST. Development of the mechanical properties of engineered skin substitutes after grafting to full-thickness wounds. <i>J Biomech Eng</i> . 2014;136(5):051008.	4
154	Fang T, Lineaweaver WC, Sailes FC, Kisner C, Zhang F. Clinical application of cultured epithelial autografts on acellular dermal matrices in the treatment of extended burn injuries. <i>Ann Plast Surg</i> . 2014;73(5):509-15.	6
155	Vazirani J, Basu S, Kenia H, Ali MH, Kacham S, Mariappan I, et al. Unilateral partial limbal stem cell deficiency: contralateral versus ipsilateral autologous cultivated limbal epithelial transplantation. <i>Am J Ophthalmol</i> . 2014;157(3):584-90.e1-2.	1
156	Boekema BK, Vlig M, Olde Damink L, Middelkoop E, Eummelen L, Buhren AV, et al. Effect of pore size and cross-linking of a novel collagen-elastin dermal substitute on wound healing. <i>J</i> . 2014;25(2):423-33.	4
157	Xiao H, Li C, Zhou X, Wang X, Wu Z, Zhang L, et al. A new method of microskin autografting with a Vaseline-based moisture dressing on granulation tissue. <i>Burns</i> . 2014;40(2):337-46.	2
158	Pellegrini G, Rama P, Di Rocco A, Panaras A, De Luca M. Concise review: hurdles in a successful example of limbal stem cell-based regenerative medicine. <i>Stem Cells</i> . 2014;32(1):26-34.	6
159	Nam JJ, Chung KK, King BT, Jones JA, Cancio LC, Baer DG, et al. Citation classics in the burn literature during the past 55 years. <i>J Burn Care Res</i> . 2014;35(2):176-85.	6
160	Gaddipati S, Muralidhar R, Sangwan VS, Mariappan I, Vemuganti GK, Balasubramanian D. Oral epithelial cells transplanted on to corneal surface tend to adapt to the ocular phenotype. <i>Indian J Ophthalmol</i> . 2014;62(5):644-8.	1
161	Sharma S, Tandon R, Mohanty S, Kashyap S, Vanathi M. Phenotypic evaluation of severely damaged ocular surface after reconstruction by cultured limbal epithelial cell transplantation. <i>Ophthalmic Res</i> . 2013;50(1):59-64.	1
162	Bateman C. 'Boston' skin cloning an exception - burns surgeons. <i>Samj, S</i> . 2013;103(6):358-9.	5
163	Pellegrini G, Rama P, Matuska S, Lambiase A, Bonini S, Pocobelli A, et al. Biological parameters determining the clinical outcome of autologous cultures of limbal stem cells. <i>Regen</i> . 2013;8(5):553-67.	1

연번	서지정보	배제 사유
164	Zamora DO, Natesan S, Becerra S, Wrice N, Chung E, Suggs LJ, et al. Enhanced wound vascularization using a dsASCs seeded FPEG scaffold. <i>Angiogenesis</i> . 2013;16(4):745-57.	4
165	Allouni A, Papini R, Lewis D. Spray-on-skin cells in burns: a common practice with no agreed protocol. <i>Burns</i> . 2013;39(7):1391-4.	6
166	Nair RP, Krishnan LK. Identification of p63+ keratinocyte progenitor cells in circulation and their matrix-directed differentiation to epithelial cells. <i>Stem Cell Res Ther</i> . 2013;4(2):38.	2
167	Mineo A, Suzuki R, Kuroyanagi Y. Development of an artificial dermis composed of hyaluronic acid and collagen. <i>J Biomater Sci Polym Ed</i> . 2013;24(6):726-40.	4
168	Sejpal K, Ali MH, Maddileti S, Basu S, Ramappa M, Kekunnaya R, et al. Cultivated limbal epithelial transplantation in children with ocular surface burns. <i>JAMA Ophthalmol</i> . 2013;131(6):731-6.	1
169	Frew Q, Philp B, Shelley O, Myers S, Navsaria H, Dziewulski P. The use of Biobrane R as a delivery method for cultured epithelial autograft in burn patients. <i>Burns</i> . 2013;39(5):876-80.	6
170	Matsumura H, Gondo M, Imai R, Shibata D, Watanabe K. Chronological histological findings of cultured epidermal autograft over bilayer artificial dermis. <i>Burns</i> . 2013;39(4):705-13.	6
171	Rignault-Clerc S, Biemann C, Delodder F, Raffoul W, Waeber B, Liudet L, et al. Functional late outgrowth endothelial progenitors isolated from peripheral blood of burned patients. <i>Burns</i> . 2013;39(4):694-704.	4
172	Menon S, Li Z, Harvey JG, Holland AJ. The use of the Meek technique in conjunction with cultured epithelial autograft in the management of major paediatric burns. <i>Burns</i> . 2013;39(4):674-9.	2
173	Lee SM, Kim MK, Shin MS, Wee WR. Matrix metalloproteinase-9 expression in the Seoul-type keratoprosthesis implanted corneas with concurrent cultivated autologous oral mucosal epithelial cell transplantation. <i>Graefes Arch Clin Exp Ophthalmol</i> . 2013;251(2):619-22.	5
174	Busch KH, Bender R, Walezko N, Aziz H, Altintas MA, Aust MC. Combination of medical needling and non-cultured autologous skin cell transplantation (renovacell) for repigmentation of hypopigmented burn scars in children and young people. <i>Ann</i> . 2016;29(2):116-22.	2
175	Larouche D, Cantin-Warren L, Desgagne M, Guignard R, Martel I, Ayoub A, et al. Improved Methods to Produce Tissue-Engineered Skin Substitutes Suitable for the Permanent Closure of Full-Thickness Skin Injuries. <i>BioRes</i> . 2016;5(1):320-9.	4
176	Hayashi M, Muramatsu H, Nakano M, Yamamoto N, Tokunaka R, Umezawa K, et al. Changes in the Dermal Structure during Cultured Epidermal Autograft Engraftment Process. <i>Plast</i> . 2016;4(9):e870.	3
177	Chua AW, Khoo YC, Tan BK, Tan KC, Foo CL, Chong SJ. Skin tissue engineering advances in severe burns: review and therapeutic applications. <i>Burns Trauma</i> . 2016;4:3.	6
178	Singh M, Nuutlia K, Chauhan AS, Eriksson E. Invasive Squamous Cell Carcinoma in Full-thickness Burn Wounds After Treatment with Cultured Epithelial Autografts. <i>Plast</i> . 2015;3(7):e460.	5
179	Taghiabadi E, Mohammadi P, Aghdami N, Falah N, Orouji Z, Nazari A, et al. Treatment of Hypertrophic Scar in Human with Autologous Transplantation of Cultured Keratinocytes and Fibroblasts along with Fibrin Glue. <i>Cell j</i> . 2015;17(1):49-58.	2
180	Urushihara H, Kawakami K. Academic Clinical Trials and Drug Regulations in Japan: Impacts of Introducing the Investigational New Drug System. <i>Ther</i> . 2014;48(4):463-72.	2

연번	서지정보	배제 사유
181	Moghaddam AS, Raji A, Movaffagh J, Yazdi AT, Mahmoudi M. Effects of autologous keratinocyte cell spray with and without chitosan on third degree burn healing: an animal experiment. <i>Wounds</i> . 2014;26(4):109-17.	4
182	McHeik JN, Barrault C, Levard G, Morel F, Bernard FX, Lecron JC. Epidermal healing in burns: autologous keratinocyte transplantation as a standard procedure: update and perspective. <i>Plast</i> . 2014;2(9):e218.	6
183	Rogers AD, Allorto NL, Adams S, Adams KG, Hudson DA, Rode H. Isn't it time for a cadaver skin bank in South Africa? <i>Ann</i> . 2013;26(3):142-6.	6
184	Bisson F, Rochefort E, Lavoie A, Larouche D, Zaniolo K, Simard-Bisson C, et al. Irradiated human dermal fibroblasts are as efficient as mouse fibroblasts as a feeder layer to improve human epidermal cell culture lifespan. <i>Int</i> . 2013;14(3):4684-704.	4
185	De Corte P, Verween G, Verbeken G, Rose T, Jennes S, De Coninck A, et al. Feeder layer- and animal product-free culture of neonatal foreskin keratinocytes: improved performance, usability, quality and safety. <i>Cell Tissue Bank</i> . 2012;13(1):175-89.	4
186	Verbeken G, Verween G, De Vos D, Pascual B, De Corte P, Richters C, et al. Glycerol treatment as recovery procedure for cryopreserved human skin allografts positive for bacteria and fungi. <i>Cell Tissue Bank</i> . 2012;13(1):1-7.	2
187	Orsag M, Bukovcan P, Koller J. Cultured keratinocytes and their possible applications. <i>Acta Chir Plast</i> . 2012;54(2):67-70.	4
188	Barrandon Y, Grasset N, Zaffalon A, Gorostidi F, Claudinot S, Droz-Georget SL, et al. Capturing epidermal stemness for regenerative medicine. <i>Semin Cell Dev Biol</i> . 2012;23(8):937-44.	2
189	Leonardi D, Oberdoerfer D, Fernandes MC, Meurer RT, Pereira-Filho GA, Cruz P, et al. Mesenchymal stem cells combined with an artificial dermal substitute improve repair in full-thickness skin wounds. <i>Burns</i> . 2012;38(8):1143-50.	4
190	Corradini F, Venturi B, Pellegrini G, De Luca M. Methods for characterization/manipulation of human corneal stem cells and their applications in regenerative medicine. <i>Methods Mol Biol</i> . 2012;916:357-72.	1
191	Dragunova J, Koller J, Cucorova V. [Employment of current results of tissue engineering in the development of skin substitutes]. <i>Cas Lek Cesk</i> . 2012;151(6):286-93.	7
192	Fritz JR, Phillips BT, Conkling N, Fourman M, Melendez MM, Bhatnagar D, et al. Comparison of native porcine skin and a dermal substitute using tensiometry and digital image speckle correlation. <i>Ann Plast Surg</i> . 2012;69(4):462-7.	4
193	Chen HC, Yeh LK, Tsai YJ, Lai CH, Chen CC, Lai JY, et al. Expression of angiogenesis-related factors in human corneas after cultivated oral mucosal epithelial transplantation. <i>Invest Ophthalmol Vis Sci</i> . 2012;53(9):5615-23.	1
194	Lee H. Outcomes of sprayed cultured epithelial autografts for full-thickness wounds: a single-centre experience. <i>Burns</i> . 2012;38(6):931-6.	6
195	Li Z, Overend C, Maitz P, Kennedy P. Quality evaluation of meshed split-thickness skin grafts stored at 4degreeC in isotonic solutions and nutrient media by cell cultures. <i>Burns</i> . 2012;38(6):899-907.	2
196	Sangwan VS, Basu S, MacNeil S, Balasubramanian D. Simple limbal epithelial transplantation (SLET): a novel surgical technique for the treatment of unilateral limbal stem cell deficiency. <i>Br J Ophthalmol</i> . 2012;96(7):931-4.	1
197	Basu S, Ali H, Sangwan VS. Clinical outcomes of repeat autologous cultivated limbal epithelial transplantation for ocular surface burns. <i>Am J Ophthalmol</i> . 2012;153(4):643-50, 50.e1-2.	1
198	Greenwood JE, Dearman BL. Comparison of a sealed, polymer foam biodegradable temporizing matrix against Integra R dermal regeneration template in a porcine wound model. <i>J Burn Care Res</i> . 2012;33(1):163-73.	4

연번	서지정보	배제 사유
199	Eldardiri M, Martin Y, Roxburgh J, Lawrence-Watt DJ, Sharpe JR. Wound contraction is significantly reduced by the use of microcarriers to deliver keratinocytes and fibroblasts in an in vivo pig model of wound repair and regeneration. <i>Tissue Eng Part A</i> . 2012;18(5-6):587-97.	4
200	Sakrak T, Kose AA, Kivanc O, Ozer MC, Cosan DT, Soyocak A, et al. The effects of combined application of autogenous fibroblast cell culture and full-tissue skin graft (FTSG) on wound healing and contraction in full-thickness tissue defects. <i>Burns</i> . 2012;38(2):225-31.	4
201	Dragunova J, Kabat P, Koller J, Jarabinska V. Experience gained during the long term cultivation of keratinocytes for treatment of burns patients. <i>Cell Tissue Bank</i> . 2012;13(3):471-8.	4
202	Marchini G, Pedrotti E, Pedrotti M, Barbaro V, Di Iorio E, Ferrari S, et al. Long-term effectiveness of autologous cultured limbal stem cell grafts in patients with limbal stem cell deficiency due to chemical burns. <i>Clin Experiment Ophthalmol</i> . 2012;40(3):255-67.	1
203	Coolen NA, Verkerk M, Reijnen L, Vlig M, Van Den Bogardt AJ, Breetveld M, et al. Culture of Keratinocytes for Transplantation without the Need of Feeder Layer Cells. <i>Cell Transplant</i> . 2007;16(6):649-61.	4
204	Lu G, Cai L, Zhao P, Gu Z, Zhu Y. Mixed Suspension of Cultured Autologous and Allogenic Keratinocytes in Fibrin Glue for the Treatment of Full-thickness Burns. <i>Wounds</i> . 2011;23(2):32-7.	2
205	Kazmi B, Inglefield CJ, Lewis MP. Autologous cell therapy: current treatments and future prospects. <i>Wounds</i> . 2009;21(9):234-42.	5
206	Saaq M, Zaib S, Ahmad S. Early excision and grafting versus delayed excision and grafting of deep thermal burns up to 40% total body surface area: a comparison of outcome. <i>Ann</i> . 2012;25(3):143-7.	2
207	Chan RK, Zamora DO, Wrice NL, Baer DG, Renz EM, Christy RJ, et al. Development of a vascularized skin construct using adipose-derived stem cells from debrided burned skin. <i>Stem Cells Int</i> . 2012;2012:841203.	2
208	Jubin K, Martin Y, Lawrence-Watt DJ, Sharpe JR. A fully autologous co-culture system utilising non-irradiated autologous fibroblasts to support the expansion of human keratinocytes for clinical use. <i>Cytotechnology</i> . 2011;63(6):655-62.	4
209	Khadjibayev AM, Fayazov AD, Djabriyev DA, Kamilov UR. Surgical treatment of deep burns. <i>Ann</i> . 2008;21(3):150-2.	2
210	Thepot A, Desanlis A, Venet E, Thivillier L, Justin V, Morel AP, et al. Assessment of transformed properties in vitro and of tumorigenicity in vivo in primary keratinocytes cultured for epidermal sheet transplantation. <i>J Skin Cancer</i> . 2011;2011:936546.	4
211	McKegney M, Taggart I, Grant MH. The influence of crosslinking agents and diamines on the pore size, morphology and the biological stability of collagen sponges and their effect on cell penetration through the sponge matrix. <i>J</i> . 2001;12(9):833-44.	4
212	Dobrowolski D, Wylegala E, Orzechowska-Wylegala B, Wowra B, Wroblewska-Czajka E. Application of autologous cultivated corneal epithelium for corneal limbal stem cell insufficiency--short-term results. <i>Klin Oczna</i> . 2011;113(10-12):346-51.	1
213	Dobrowolski D, Wylegala E, Orzechowska-Wylegala B, Wowra B, Bielecka A, Obuchowicz E, et al. Culture morphology of the autologous cultivated corneal epithelium. <i>Klin Oczna</i> . 2011;113(7-9):249-53.	1
214	Yamato M, Okano T. [The history and current status of clinical applications of stem cells]. <i>Nippon Rinsho</i> . 2011;69(12):2103-8.	7
215	Keck M, Haluza D, Selig HF, Jahl M, Lumenta DB, Kamolz LP, et al. Adipose tissue engineering: three different approaches to seed preadipocytes on a collagen-elastin matrix. <i>Ann Plast Surg</i> . 2011;67(5):484-8.	4

연번	서지정보	배제 사유
216	Sangwan VS, Basu S, Vemuganti GK, Sejpal K, Subramaniam SV, Bandyopadhyay S, et al. Clinical outcomes of xeno-free autologous cultivated limbal epithelial transplantation: a 10-year study. <i>Br J Ophthalmol.</i> 2011;95(11):1525-9.	1
217	Basu S, Mohamed A, Chaurasia S, Sejpal K, Vemuganti GK, Sangwan VS. Clinical outcomes of penetrating keratoplasty after autologous cultivated limbal epithelial transplantation for ocular surface burns. <i>Am J Ophthalmol.</i> 2011;152(6):917-24.e1.	1
218	Varkey M, Ding J, Tredget EE. Differential collagen-glycosaminoglycan matrix remodeling by superficial and deep dermal fibroblasts: potential therapeutic targets for hypertrophic scar. <i>Biomaterials.</i> 2011;32(30):7581-91.	4
219	Natesan S, Wrice NL, Baer DG, Christy RJ. Debrided skin as a source of autologous stem cells for wound repair. <i>Stem Cells.</i> 2011;29(8):1219-30.	4
220	Takeda K, Nakamura T, Inatomi T, Sotozono C, Watanabe A, Kinoshita S. Ocular surface reconstruction using the combination of autologous cultivated oral mucosal epithelial transplantation and eyelid surgery for severe ocular surface disease. <i>Am J Ophthalmol.</i> 2011;152(2):195-201.e1.	1
221	Satake Y, Higa K, Tsubota K, Shimazaki J. Long-term outcome of cultivated oral mucosal epithelial sheet transplantation in treatment of total limbal stem cell deficiency. <i>Ophthalmology.</i> 2011;118(8):1524-30.	1
222	Anderson JR, Fear MW, Phillips JK, Dawson LF, Wallace H, Wood FM, et al. A preliminary investigation of the reinnervation and return of sensory function in burn patients treated with INTEGRA R. <i>Burns.</i> 2011;37(7):1101-8.	2
223	Yim H, Yang HT, Cho YS, Seo CH, Lee BC, Ko JH, et al. Clinical study of cultured epithelial autografts in liquid suspension in severe burn patients. <i>Burns.</i> 2011;37(6):1067-71.	6
224	Leclerc T, Thepenier C, Jault P, Bey E, Peltzer J, Trouillas M, et al. Cell therapy of burns. <i>Cell Prolif.</i> 2011;44 Suppl 1:48-54.	6
225	Gerlach JC, Johnen C, Ottomann C, Brautigam K, Plettig J, Belfekroun C, et al. Method for autologous single skin cell isolation for regenerative cell spray transplantation with non-cultured cells. <i>Int J Artif Organs.</i> 2011;34(3):271-9.	2
226	Posluszny JA, Jr., Conrad P, Halerz M, Shankar R, Gamelli RL. Surgical burn wound infections and their clinical implications. <i>J Burn Care Res.</i> 2011;32(2):324-33.	3
227	Lugo LM, Lei P, Andreadis ST. Vascularization of the dermal support enhances wound re-epithelialization by in situ delivery of epidermal keratinocytes. <i>Tissue Eng Part A.</i> 2011;17(5-6):665-75.	4
228	Bargues L, Prat M, Leclerc T, Bey E, Lataillade JJ. [Present and future of cell therapy in burns]. <i>Pathol Biol (Paris).</i> 2011;59(3):e49-56.	7
229	Chaouat M, Zakine G, Mimoun M. [Principles of the local treatment: Surgical processing]. <i>Pathol Biol (Paris).</i> 2011;59(3):e57-61.	7
230	Lataillade JJ, Bey E, Thepenier C, Prat M, Leclerc T, Bargues L. [Skin engineering for burns treatment]. <i>Bull Acad Natl Med.</i> 2010;194(7):1339-51.	7
231	Ueda M. Sprayed cultured mucosal epithelial cell for deep dermal burns. <i>J Craniofac Surg.</i> 2010;21(6):1729-32.	2
232	Niu Z, Yu J, Jin Z, Zhao J, Shi K, Liu J. Nonantibiotic therapy for multidrug-resistant <i>Pseudomonas aeruginosa</i> in a patient with chronic burn wounds. <i>Am J Med Sci.</i> 2010;340(6):521-3.	6
233	Forouzandeh F, Jalili RB, Hartwell RV, Allan SE, Boyce S, Supp D, et al. Local expression of indoleamine 2,3-dioxygenase suppresses T-cell-mediated rejection of an engineered bilayer skin substitute. <i>Wound Repair Regen.</i> 2010;18(6):614-23.	4
234	Castagnoli C, Fumagalli M, Alotto D, Cambieri I, Casarin S, Ostorero A, et al. Preparation and characterization of a novel skin substitute. <i>J Biomed Biotechnol.</i> 2010.	4

연번	서지정보	배제 사유
235	Hayama Y, Ueda K, Kuroyanagi Y. Optimum preservation for autologous cultured dermal substitutes. <i>Ann Plast Surg.</i> 2010;65(3):338-43.	4
236	Sood R, Roggy D, Zieger M, Balledux J, Chaudhari S, Koumanis DJ, et al. Cultured epithelial autografts for coverage of large burn wounds in eighty-eight patients: the Indiana University experience. <i>J Burn Care Res.</i> 2010;31(4):559-68.	6
237	Agay D, Scherthan H, Forcheron F, Grenier N, Herodin F, Meineke V, et al. Multipotent mesenchymal stem cell grafting to treat cutaneous radiation syndrome: development of a new minipig model. <i>Exp Hematol.</i> 2010;38(10):945-56.	4
238	Rama P, Matuska S, Paganoni G, Spinelli A, De Luca M, Pellegrini G. Limbal stem-cell therapy and long-term corneal regeneration. <i>N Engl J Med.</i> 2010;363(2):147-55.	1
239	Gibot L, Galbraith T, Huot J, Auger FA. A preexisting microvascular network benefits in vivo revascularization of a microvascularized tissue-engineered skin substitute. <i>Tissue Eng Part A.</i> 2010;16(10):3199-206.	4
240	Pauklin M, Fuchsluger TA, Westekemper H, Steuhl KP, Meller D. Midterm results of cultivated autologous and allogeneic limbal epithelial transplantation in limbal stem cell deficiency. <i>Dev Ophthalmol.</i> 2010;45:57-70.	1
241	Baradaran-Rafii A, Ebrahimi M, Kanavi MR, Taghi-Abadi E, Aghdami N, Eslani M, et al. Midterm outcomes of autologous cultivated limbal stem cell transplantation with or without penetrating keratoplasty. <i>Cornea.</i> 2010;29(5):502-9.	1
242	Liu J, Bian Z, Kuijpers-Jagtman AM, Von den Hoff JW. Skin and oral mucosa equivalents: construction and performance. <i>Orthod Craniofac Res.</i> 2010;13(1):11-20.	6
243	Leon-Villapalos J, Eldardiri M, Dziewulski P. The use of human deceased donor skin allograft in burn care. <i>Cell Tissue Bank.</i> 2010;11(1):99-104.	2
244	Klingenberg JM, McFarland KL, Friedman AJ, Boyce ST, Aronow BJ, Supp DM. Engineered human skin substitutes undergo large-scale genomic reprogramming and normal skin-like maturation after transplantation to athymic mice. <i>J Invest Dermatol.</i> 2010;130(2):587-601.	4
245	Oliveira-Prado R, Caldas IR, Pereira GM, Andrade MV, Cunha-Melo JR. Swine pericardium as dermal substrate for human keratinocyte culture. <i>Br J Dermatol.</i> 2010;162(2):251-7.	4
246	Gallego L, Junquera L, Villarreal P, Pena I, Meana A. Use of cultured human epithelium for coverage: a defect of radial forearm free flap donor site. <i>Med Oral Patol Oral Cir Bucal.</i> 2010;15(1):e58-60.	6
247	James SE, Booth S, Dheansa B, Mann DJ, Reid MJ, Shevchenko RV, et al. Sprayed cultured autologous keratinocytes used alone or in combination with meshed autografts to accelerate wound closure in difficult-to-heal burns patients. <i>Burns.</i> 2010;36(3):e10-20.	2
248	Scuderi N, Anniboletti T, Carlesimo B, Onesti MG. Clinical application of autologous three-cellular cultured skin substitutes based on esterified hyaluronic acid scaffold: our experience. <i>In Vivo.</i> 2009;23(6):991-1003.	1
249	Guenou H, Nissan X, Larcher F, Feteira J, Lemaitre G, Saidani M, et al. Human embryonic stem-cell derivatives for full reconstruction of the pluristratified epidermis: a preclinical study. <i>Lancet.</i> 2009;374(9703):1745-53.	4
250	Sheridan R. Closure of the excised burn wound: autografts, semipermanent skin substitutes, and permanent skin substitutes. <i>Clin Plast Surg.</i> 2009;36(4):643-51.	6
251	Ma DH, Kuo MT, Tsai YJ, Chen HC, Chen XL, Wang SF, et al. Transplantation of cultivated oral mucosal epithelial cells for severe corneal burn. <i>Eye.</i> 2009;23(6):1442-50.	1
252	Inoue S, Kijima H, Kidokoro M, Tanaka M, Suzuki Y, Motojuku M, et al. The effectiveness of basic fibroblast growth factor in fibrin-based cultured skin substitute in vivo. <i>J Burn Care Res.</i> 2009;30(3):514-9.	4

연번	서지정보	배제 사유
253	Bargues L, Boyer S, Leclerc T, Duhamel P, Bey E. [Incidence and microbiology of infectious complications with the use of artificial skin Integra in burns]. <i>Ann Chir Plast Esthet.</i> 2009;54(6):533-9.	7
254	Lumenta DB, Kamolz LP, Frey M. Adult burn patients with more than 60% TBSA involved-Meek and other techniques to overcome restricted skin harvest availability--the Viennese Concept. <i>J Burn Care Res.</i> 2009;30(2):231-42.	2
255	Auxenfans C, Builles N, Andre V, Lequeux C, Fievet A, Rose S, et al. [Porous matrix and primary-cell culture: a shared concept for skin and cornea tissue engineering]. <i>Pathol Biol (Paris).</i> 2009;57(4):290-8.	7
256	Dorai AA, Lim CK, Fareha AC, Halim AS. Cultured epidermal autografts in combination with MEEK Micrografting technique in the treatment of major burn injuries. <i>Med J Malaysia.</i> 2008;63 Suppl A:44.	5
257	Green H. The birth of therapy with cultured cells. <i>Bioessays.</i> 2008;30(9):897-903.	5
258	Lineen E, Namias N. Biologic dressing in burns. <i>J Craniofac Surg.</i> 2008;19(4):923-8.	6
259	Bass MJ, Phillips LG. Economics of pediatric burns. <i>J Craniofac Surg.</i> 2008;19(4):888-90.	6
260	Haertsch PA. Re: "The use of a non-cultured autologous cell suspension and Integra) dermal regeneration template to repair full thickness skin wounds in a porcine skin model: a one-step process" [ <i>Burns</i> 2007;33(6):693-700]. <i>Burns.</i> 2008;34(6):899; author reply 900-1.	5
261	Halmy C, Nadai Z, Juhasz Z, Pesthy P, Marczell Z, Szetei K, et al. [Skin replacement with biological and biosynthetic materials following burn injury]. <i>Orv Hetil.</i> 2008;149(20):915-9.	7
262	Warner PM, Yakuboff KP, Kagan RJ, Boyce S, Warden GD. An 18-year experience in the management of congenital nevomelanocytic nevi. <i>Ann Plast Surg.</i> 2008;60(3):283-7.	1
263	Wilshaw SP, Kearney J, Fisher J, Ingham E. Biocompatibility and potential of acellular human amniotic membrane to support the attachment and proliferation of allogeneic cells. <i>Tissue Eng Part A.</i> 2008;14(4):463-72.	4
264	Eves PC, Bullett NA, Haddow D, Beck AJ, Layton C, Way L, et al. Simplifying the delivery of melanocytes and keratinocytes for the treatment of vitiligo using a chemically defined carrier dressing. <i>J Invest Dermatol.</i> 2008;128(6):1554-64.	4
265	Bannasch H, Stark GB, Knam F, Horch RE, Fohn M. Decellularized dermis in combination with cultivated keratinocytes in a short- and long-term animal experimental investigation. <i>J Eur Acad Dermatol Venereol.</i> 2008;22(1):41-9.	4
266	Chua AW, Ma DR, Song IC, Phan TT, Lee ST, Song C. In vitro evaluation of fibrin mat and Tegaderm wound dressing for the delivery of keratinocytes--implications of their use to treat burns. <i>Burns.</i> 2008;34(2):175-80.	4
267	Mary O'Brien C, Breuning E, Webb J, Balderson D, Nancarrow J. A prospective randomised controlled trial to investigate the effect of local anaesthetic in vivo on cell culture. <i>J Plast Reconstr Aesthet Surg.</i> 2008;61(10):1232-4.	2
268	Zweifel CJ, Contaldo C, Kohler C, Jandali A, Kunzi W, Giovanoli P. Initial experiences using non-cultured autologous keratinocyte suspension for burn wound closure. <i>J Plast Reconstr Aesthet Surg.</i> 2008;61(11):e1-4.	4
269	Fredriksson C, Kratz G, Huss F. Transplantation of cultured human keratinocytes in single cell suspension: a comparative in vitro study of different application techniques. <i>Burns.</i> 2008;34(2):212-9.	4
270	Fohn M, Bannasch H. Artificial skin. <i>Methods Mol Med.</i> 2007;140:167-82.	5
271	Hernon CA, Harrison CA, Thornton DJ, MacNeil S. Enhancement of keratinocyte performance in the production of tissue-engineered skin using a low-calcium medium. <i>Wound Repair Regen.</i> 2007;15(5):718-26.	4



연번	서지정보	배제 사유
272	Spiekstra SW, Breetveld M, Rustemeyer T, Scheper RJ, Gibbs S. Wound-healing factors secreted by epidermal keratinocytes and dermal fibroblasts in skin substitutes. <i>Wound Repair Regen.</i> 2007;15(5):708-17.	4
273	Luengo Gimeno F, Lavigne V, Gatto S, Croxatto JO, Correa L, Gallo JE. Advances in corneal stem-cell transplantation in rabbits with severe ocular alkali burns. <i>J Cataract Refract Surg.</i> 2007;33(11):1958-65.	4
274	Pellegrini G, De Luca M, Arsenijevic Y. Towards therapeutic application of ocular stem cells. <i>Semin Cell Dev Biol.</i> 2007;18(6):805-18.	1
275	Lataillade JJ, Doucet C, Bey E, Carsin H, Huet C, Clairand I, et al. New approach to radiation burn treatment by dosimetry-guided surgery combined with autologous mesenchymal stem cell therapy. <i>Regen.</i> 2007;2(5):785-94.	6
276	Myers SR, Partha VN, Soranzo C, Price RD, Navsaria HA. Hyalomatrix: a temporary epidermal barrier, hyaluronan delivery, and neodermis induction system for keratinocyte stem cell therapy. <i>Tissue Eng.</i> 2007;13(11):2733-41.	4
277	Hata K. Current issues regarding skin substitutes using living cells as industrial materials. <i>J Artif Organs.</i> 2007;10(3):129-32.	6
278	Pham C, Greenwood J, Cleland H, Woodruff P, Maddern G. Bioengineered skin substitutes for the management of burns: a systematic review. <i>Burns.</i> 2007;33(8):946-57.	6
279	Chai JK, Sheng ZY, Ma ZF, Yang HM, Liu Q, Liang LM. Growth activity of epidermal cells from different parts of human body. <i>Chin Med J.</i> 2007;120(16):1444-7.	4
280	Fu X, Fang L, Li H, Li X, Cheng B, Sheng Z. Adipose tissue extract enhances skin wound healing. <i>Wound Repair Regen.</i> 2007;15(4):540-8.	4
281	Gardner RL. Stem cells and regenerative medicine: principles, prospects and problems. <i>C R Biol.</i> 2007;330(6-7):465-73.	4
282	Wood FM, Stoner ML, Fowler BV, Fear MW. The use of a non-cultured autologous cell suspension and Integra dermal regeneration template to repair full-thickness skin wounds in a porcine model: a one-step process. <i>Burns.</i> 2007;33(6):693-700.	4
283	Gustafson CJ, Birgisson A, Junker J, Huss F, Salemark L, Johnson H, et al. Employing human keratinocytes cultured on macroporous gelatin spheres to treat full thickness-wounds: an in vivo study on athymic rats. <i>Burns.</i> 2007;33(6):726-35.	4
284	Marazzi M, De Angelis A, Ravizza A, Ordanini MN, Falcone L, Chiaratti A, et al. Successful management of deep facial burns in a patient with extensive third-degree burns: the role of a nanocrystalline silver dressing in facilitating resurfacing. <i>Int Wound J.</i> 2007;4(1):8-14.	6
285	Sobral CS, Gragnani A, Morgan J, Ferreira LM. Inhibition of proliferation of <i>Pseudomonas aeruginosa</i> by KGF in an experimental burn model using human cultured keratinocytes. <i>Burns.</i> 2007;33(5):613-20.	4
286	Atiyeh BS, Costagliola M. Cultured epithelial autograft (CEA) in burn treatment: three decades later. <i>Burns.</i> 2007;33(4):405-13.	6
287	Burd A. Evaluating the use of hydrogel sheet dressings in comprehensive burn wound care. <i>Ostomy Wound Manage.</i> 2007;53(3):52-62.	2
288	Kalyanaraman B, Boyce S. Assessment of an automated bioreactor to propagate and harvest keratinocytes for fabrication of engineered skin substitutes. <i>Tissue Eng.</i> 2007;13(5):983-93.	4
289	Fatima A, Vemuganti GK, Iftekhar G, Rao GN, Sangwan VS. In vivo survival and stratification of cultured limbal epithelium. <i>Clin Experiment Ophthalmol.</i> 2007;35(1):96-8.	1
290	Magnusson M, Papini RP, Rea SM, Reed CC, Wood FM. Cultured autologous keratinocytes in suspension accelerate epithelial maturation in an in vivo wound model as measured by surface electrical capacitance. <i>Plast Reconstr Surg.</i> 2007;119(2):495-9.	4

연번	서지정보	배제 사유
291	Chiarini A, Dal Pra I, Armato U. In vitro and in vivo characteristics of frozen/thawed neonatal pig split-skin strips: a novel biologically active dressing for areas of severe, acute or chronic skin loss. <i>Int J Mol Med.</i> 2007;19(2):245-55.	4
292	Price RD, Das-Gupta V, Leigh IM, Navsaria HA. A comparison of tissue-engineered hyaluronic acid dermal matrices in a human wound model. <i>Tissue Eng.</i> 2006;12(10):2985-95.	1
293	De Luca M, Pellegrini G, Green H. Regeneration of squamous epithelia from stem cells of cultured grafts. <i>Regen.</i> 2006;1(1):45-57.	1
294	Hernon CA, Dawson RA, Freedlander E, Short R, Haddow DB, Brotherston M, et al. Clinical experience using cultured epithelial autografts leads to an alternative methodology for transferring skin cells from the laboratory to the patient. <i>Regen.</i> 2006;1(6):809-21.	2
295	Hirsch T, von Peter S, Dubin G, Mittler D, Jacobsen F, Lehnhardt M, et al. Adenoviral gene delivery to primary human cutaneous cells and burn wounds. <i>Mol Med.</i> 2006;12(9-10):199-207.	4
296	Fatima A, Sangwan VS, Iftekhhar G, Reddy P, Matalia H, Balasubramanian D, et al. Technique of cultivating limbal derived corneal epithelium on human amniotic membrane for clinical transplantation. <i>J Postgrad Med.</i> 2006;52(4):257-61.	1
297	Muangman P, Deubner H, Honari S, Heimbach DM, Engrav LH, Klein MB, et al. Correlation of clinical outcome of integra application with microbiologic and pathological biopsies. <i>J Trauma.</i> 2006;61(5):1212-7.	2
298	Fujimori Y, Ueda K, Fumimoto H, Kubo K, Kuroyanagi Y. Skin regeneration for children with burn scar contracture using autologous cultured dermal substitutes and superthin auto-skin grafts: preliminary clinical study. <i>Ann Plast Surg.</i> 2006;57(4):408-14.	2
299	Ehrenreich M, Ruszczak Z. Update on tissue-engineered biological dressings. <i>Tissue Eng.</i> 2006;12(9):2407-24.	6
300	Wang TW, Sun JS, Wu HC, Tsuang YH, Wang WH, Lin FH. The effect of gelatin-chondroitin sulfate-hyaluronic acid skin substitute on wound healing in SCID mice. <i>Biomaterials.</i> 2006;27(33):5689-97.	4
301	Norris MM, Aksan A, Sugimachi K, Toner M. 3-O-methyl-D-glucose improves desiccation tolerance of keratinocytes. <i>Tissue Eng.</i> 2006;12(7):1873-9.	4
302	Shealy FG, Jr., DeLoach ED. Experience with the use of apligraf to heal complicated surgical and nonsurgical wounds in a private practice setting. <i>Adv Skin Wound Care.</i> 2006;19(6):310-22.	2
303	Kuroyanagi Y. [Regenerative medicine for skin]. <i>Nippon Ronen Igakkai Zasshi.</i> 2006;43(3):326-9.	7
304	Gutierrez P, Taggart S. Cultured epithelial autografts. <i>Aust Nurs J.</i> 2006;13(11):37.	5
305	Wood FM, Kolybaba ML, Allen P. The use of cultured epithelial autograft in the treatment of major burn wounds: eleven years of clinical experience. <i>Burns.</i> 2006;32(5):538-44.	6
306	Wood FM, Kolybaba ML, Allen P. The use of cultured epithelial autograft in the treatment of major burn injuries: a critical review of the literature. <i>Burns.</i> 2006;32(4):395-401.	6
307	Boyce ST, Kagan RJ, Greenhalgh DG, Warner P, Yakuboff KP, Palmieri T, et al. Cultured skin substitutes reduce requirements for harvesting of skin autograft for closure of excised, full-thickness burns. <i>J Trauma.</i> 2006;60(4):821-9.	2
308	Ozturk E, Ergun MA, Ozturk Z, Nurozler AB, Kecici K, Ozdemir N, et al. Chitosan-coated alginate membranes for cultivation of limbal epithelial cells to use in the restoration of damaged corneal surfaces. <i>Int J Artif Organs.</i> 2006;29(2):228-38.	4

연번	서지정보	배제 사유
309	Sangwan VS, Matalia HP, Vemuganti GK, Fatima A, Ifthekar G, Singh S, et al. Clinical outcome of autologous cultivated limbal epithelium transplantation. <i>Indian J Ophthalmol.</i> 2006;54(1):29-34.	1
310	Llames S, Garcia E, Garcia V, del Rio M, Larcher F, Jorcano JL, et al. Clinical results of an autologous engineered skin. <i>Cell Tissue Bank.</i> 2006;7(1):47-53.	2
311	Theopold C, Eriksson E. The need for aggressive follow-up after cultured epidermal autograft-grafted full-thickness burn. <i>Plast Reconstr Surg.</i> 2006;117(2):708.	5
312	Johnen C, Hartmann B, Steffen I, Brautigam K, Witascheck T, Toman N, et al. Skin cell isolation and expansion for cell transplantation is limited in patients using tobacco, alcohol, or are exhibiting diabetes mellitus. <i>Burns.</i> 2006;32(2):194-200.	4
313	Caissie R, Gingras M, Champigny MF, Berthod F. In vivo enhancement of sensory perception recovery in a tissue-engineered skin enriched with laminin. <i>Biomaterials.</i> 2006;27(15):2988-93.	4
314	Harrison CA, Gossiel F, Bullock AJ, Sun T, Blumsohn A, Mac Neil S. Investigation of keratinocyte regulation of collagen I synthesis by dermal fibroblasts in a simple in vitro model. <i>Br J Dermatol.</i> 2006;154(3):401-10.	4
315	Pallua N, von Bulow S. [Methods of burn treatment. Part II: Technical aspects]. <i>Chirurg.</i> 2006;77(2):179-86; quiz 87-8.	7
316	Drewa T, Nadolski B, Czaplewski A, Drewa L. [Application of autologous and allogenic keratynocyte cultures in burn management]. <i>Przeegl Lek.</i> 2005;62(9):934-7.	7
317	Kuroyanagi Y. [Advances in regenerative medicine for skin]. <i>Nippon Ronen Igakkai Zasshi.</i> 2005;42(6):609-15.	7
318	Braye F, Hautier A, Bouez C, Damour O. [Skin substitutes reconstructed in the laboratory: application in burn treatment]. <i>Pathol Biol (Paris).</i> 2005;53(10):613-7.	7
319	Smiley AK, Klingenberg JM, Aronow BJ, Boyce ST, Kitzmiller WJ, Supp DM. Microarray analysis of gene expression in cultured skin substitutes compared with native human skin. <i>J Invest Dermatol.</i> 2005;125(6):1286-301.	4
320	Harrison CA, Dalley AJ, Mac Neil S. A simple in vitro model for investigating epithelial/mesenchymal interactions: keratinocyte inhibition of fibroblast proliferation and fibronectin synthesis. <i>Wound Repair Regen.</i> 2005;13(6):543-50.	4
321	Coruh A, Tosun Z, Ozbebit U. Close relative intermingled skin allograft and autograft use in the treatment of major burns in adults and children. <i>J Burn Care Rehabil.</i> 2005;26(6):471-7.	2
322	Wang HJ, Chou TD, Tsou TL, Chen TM, Chen SL, Chen SG, et al. The application of new biosynthetic artificial skin for long-term temporary wound coverage. <i>Burns.</i> 2005;31(8):991-7.	4
323	Horch RE, Kopp J, Kneser U, Beier J, Bach AD. Tissue engineering of cultured skin substitutes. <i>J Cell Mol Med.</i> 2005;9(3):592-608.	6
324	Hohlfeld J, de Buys Roessingh A, Hirt-Burri N, Chaubert P, Gerber S, Scaletta C, et al. Tissue engineered fetal skin constructs for paediatric burns. <i>Lancet.</i> 2005;366(9488):840-2.	2
325	Schwerdtner O, Damaskos T, Kage A, Weitzel-Kage D, Klein M. Autologous epidermal cells can induce wound closure of neurotrophic ulceration caused by trigeminal trophic syndrome. <i>Int J Oral Maxillofac Surg.</i> 2005;34(4):443-5.	1
326	Supp DM, Boyce ST. Engineered skin substitutes: practices and potentials. <i>Clin Dermatol.</i> 2005;23(4):403-12.	6
327	Rab M, Koller R, Ruzicka M, Burda G, Kamolz LP, Bierochs B, et al. Should dermal scald burns in children be covered with autologous skin grafts or with allogeneic cultivated keratinocytes?--"The Viennese concept". <i>Burns.</i> 2005;31(5):578-86.	2
328	Barthel R, Aberdam D. Epidermal stem cells. <i>J Eur Acad Dermatol Venereol.</i> 2005;19(4):405-13.	6

연번	서지정보	배제 사유
329	Sangwan VS, Murthy SI, Vemuganti GK, Bansal AK, Gangopadhyay N, Rao GN. Cultivated corneal epithelial transplantation for severe ocular surface disease in vernal keratoconjunctivitis. <i>Cornea</i> . 2005;24(4):426-30.	1
330	Tremblay PL, Hudon V, Berthod F, Germain L, Auger FA. Inosculation of tissue-engineered capillaries with the host's vasculature in a reconstructed skin transplanted on mice. <i>Am J Transplant</i> . 2005;5(5):1002-10.	4
331	Duncan CO, Shelton RM, Navsaria H, Balderson DS, Papini RP, Barralet JE. In vitro transfer of keratinocytes: comparison of transfer from fibrin membrane and delivery by aerosol spray. <i>J Biomed Mater Res B Appl Biomater</i> . 2005;73(2):221-8.	4
332	Sangwan VS, Matalia HP, Vemuganti GK, Iftheekar G, Fatima A, Singh S, et al. Early results of penetrating keratoplasty after cultivated limbal epithelium transplantation. <i>Arch Ophthalmol</i> . 2005;123(3):334-40.	1
333	Qing C, Cao YP, Shi JX, Zhou H, Tao J, Zheng ZX, et al. [Experimental study on the role of cytokines and keratinocytes in the survival mechanism of auto and allogeneic mixed skin grafting]. <i>Zhonghua Shao Shang Za Zhi</i> . 2004;20(6):336-9.	7
334	Hunziker T. [Autologous cultured skin substitutes]. <i>Hautarzt</i> . 2004;55(11):1077-84; quiz 85.	7
335	Mis B, Rolland E, Ronfard V. Combined use of a collagen-based dermal substitute and a fibrin-based cultured epithelium: a step toward a total skin replacement for acute wounds. <i>Burns</i> . 2004;30(7):713-9.	4
336	Supp DM, Karpinski AC, Boyce ST. Expression of human beta-defensins HBD-1, HBD-2, and HBD-3 in cultured keratinocytes and skin substitutes. <i>Burns</i> . 2004;30(7):643-8.	4
337	Dvorankova B, Broz L, Pafcuga I, Kapounkova Z, Konigova R. The role of skin bank in the treatment of severely burnt patients. <i>Acta Chir Plast</i> . 2004;46(2):51-5.	5
338	Theopold C, Hoeller D, Velander P, Demling R, Eriksson E. Graft site malignancy following treatment of full-thickness burn with cultured epidermal autograft. <i>Plast Reconstr Surg</i> . 2004;114(5):1215-9.	6
339	Nakamura T, Inatomi T, Sotozono C, Amemiya T, Kanamura N, Kinoshita S. Transplantation of cultivated autologous oral mucosal epithelial cells in patients with severe ocular surface disorders. <i>Br J Ophthalmol</i> . 2004;88(10):1280-4.	1
340	Fu XB, Fang LJ, Wang YX, Sun TZ, Cheng B. [Enhancing the repair quality of skin injury on porcine after autografting with the bone marrow mesenchymal stem cells]. <i>Chung Hua I Hsueh Tsa Chih</i> . 2004;84(11):920-4.	7
341	Harkin DG, Barnard Z, Gillies P, Ainscough SL, Apel AJ. Analysis of p63 and cytokeratin expression in a cultivated limbal autograft used in the treatment of limbal stem cell deficiency. <i>Br J Ophthalmol</i> . 2004;88(9):1154-8.	1
342	Nakamura T, Inatomi T, Sotozono C, Koizumi N, Kinoshita S. Successful primary culture and autologous transplantation of corneal limbal epithelial cells from minimal biopsy for unilateral severe ocular surface disease. <i>Acta Ophthalmol Scand</i> . 2004;82(4):468-71.	1
343	Chester DL, Balderson DS, Papini RP. A review of keratinocyte delivery to the wound bed. <i>J Burn Care Rehabil</i> . 2004;25(3):266-75.	6
344	Supp DM, Karpinski AC, Boyce ST. Vascular endothelial growth factor overexpression increases vascularization by murine but not human endothelial cells in cultured skin substitutes grafted to athymic mice. <i>J Burn Care Rehabil</i> . 2004;25(4):337-45.	4
345	Kopp J, Jeschke MG, Bach AD, Kneser U, Horch RE. Applied tissue engineering in the closure of severe burns and chronic wounds using cultured human autologous keratinocytes in a natural fibrin matrix. <i>Cell Tissue Bank</i> . 2004;5(2):89-96.	6
346	Ehrlich HP. Understanding experimental biology of skin equivalent: from laboratory to clinical use in patients with burns and chronic wounds. <i>Am J Surg</i> . 2004;187(5A):29S-33S.	4

연번	서지정보	배제 사유
347	Haertsch P. Re: Initial experience with cultured epithelial autografts in massively burnt patients. ANZ J Surg. 2004;74(3):175.	5
348	Llames SG, Del Rio M, Larcher F, Garcia E, Garcia M, Escamez MJ, et al. Human plasma as a dermal scaffold for the generation of a completely autologous bioengineered skin. Transplantation. 2004;77(3):350-5.	4
349	Vernez M, Raffoul W, Gailloud-Matthieu MC, Egloff D, Senechaud I, Panizzon RG, et al. Quantitative assessment of cell viability and apoptosis in cultured epidermal autografts: application to burn therapy. Int J Artif Organs. 2003;26(9):793-803.	6
350	Bannasch H, Fohn M, Unterberg T, Bach AD, Weyand B, Stark GB. Skin tissue engineering. Clin Plast Surg. 2003;30(4):573-9.	6
351	Currie LJ, Martin R, Sharpe JR, James SE. A comparison of keratinocyte cell sprays with and without fibrin glue. Burns. 2003;29(7):677-85.	4
352	Kadoya K, Amano S, Inomata S, Tsunenaga M, Matsuzaki K, Oshima H, et al. Evaluation of autologous cultured epithelium as replacement skin after tattoo excision: correlation between skin texture and histological features. Br J Dermatol. 2003;149(2):377-80.	1
353	Davey RB. Initial experience with cultured epithelial autografts in massively burnt patients. ANZ J Surg. 2003;73(8):662-3.	5
354	Sangwan VS, Vemuganti GK, Iftekhar G, Bansal AK, Rao GN. Use of autologous cultured limbal and conjunctival epithelium in a patient with severe bilateral ocular surface disease induced by acid injury: a case report of unique application. Cornea. 2003;22(5):478-81.	1
355	Kogan L, Govrin-Yehudain J. Vertical (two-layer) skin grafting: new reserves for autologic skin. Ann Plast Surg. 2003;50(5):514-6.	2
356	Neovius EB, Kratz G. Tissue engineering by cocultivating human elastic chondrocytes and keratinocytes. Tissue Eng. 2003;9(2):365-9.	4
357	Huang YC, Wang TW, Sun JS, Lin FH. Cultured keratinocytes and dermal fibroblasts on a double-layer scaffold with bi-medium culture system. Biomed Sci Instrum. 2003;39:500-5.	4
358	Cao Y, Zhou H, Tao J, Zheng Z, Li N, Shen B, et al. Keratinocytes induce local tolerance to skin graft by activating interleukin-10-secreting T cells in the context of costimulation molecule B7-H1. Transplantation. 2003;75(8):1390-6.	4
359	Vacher D. [Autologous epidermal sheets production for skin cellular therapy]. Ann Pharm Fr. 2003;61(3):203-6.	7
360	Counter CM, Press W, Compton CC. Telomere shortening in cultured autografts of patients with burns. Lancet. 2003;361(9366):1345-6.	5
361	Oshima H, Inoue H, Matsuzaki K, Tanabe M, Kumagai N. Permanent restoration of human skin treated with cultured epithelium grafting--wound healing by stem cell based tissue engineering. Hum Cell. 2002;15(3):118-28.	4
362	Solomon DE. An in vitro examination of an extracellular matrix scaffold for use in wound healing. Int J Exp Pathol. 2002;83(5):209-16.	4
363	Dhennin C. [Skin substitutes and skin culture]. Rev Prat. 2002;52(20):2249-52.	7
364	Xie W, Wang L, Tan H, Wang D, Liu J, Hu B, et al. [Microskin grafting by spraying in burn management]. Zhonghua Shao Shang Za Zhi. 2002;18(1):26-8.	7
365	Elliott M, Vandervord J. Initial experience with cultured epithelial autografts in massively burnt patients. ANZ J Surg. 2002;72(12):893-5.	6
366	Soler C. [Genevriev Biotechnology Center: production of autologous epidermal sheets (Epibase)]. Ann Dermatol Venereol. 2002;129(10 Pt 2):1239-41.	7
367	Dubertret L. [Cells, tissues and organs for wound cicatrization]. Ann Dermatol Venereol. 2002;129(10 Pt 2):1235-8.	7

연번	서지정보	배제 사유
368	Rose CM, Reilly KJ, Haith LR, Patton ML, Guilday RJ, Cawley MJ, et al. Emergence of resistance of vancomycin-resistant <i>Enterococcus faecium</i> in a thermal injury patient treated with quinupristin-dalfopristin and cultured epithelial autografts for wound closure. <i>Burns</i> . 2002;28(7):696-8.	6
369	Grueterich M, Espana EM, Touhami A, Ti SE, Tseng SC. Phenotypic study of a case with successful transplantation of ex vivo expanded human limbal epithelium for unilateral total limbal stem cell deficiency. <i>Ophthalmology</i> . 2002;109(8):1547-52.	1
370	Ti SE, Anderson D, Touhami A, Kim C, Tseng SC. Factors affecting outcome following transplantation of ex vivo expanded limbal epithelium on amniotic membrane for total limbal deficiency in rabbits. <i>Invest Ophthalmol Vis Sci</i> . 2002;43(8):2584-92.	4
371	Pouliot R, Larouche D, Auger FA, Juhasz J, Xu W, Li H, et al. Reconstructed human skin produced in vitro and grafted on athymic mice. <i>Transplantation</i> . 2002;73(11):1751-7.	4
372	Supp DM, Wilson-Landy K, Boyce ST. Human dermal microvascular endothelial cells form vascular analogs in cultured skin substitutes after grafting to athymic mice. <i>Faseb J</i> . 2002;16(8):797-804.	4
373	Nanchahal J, Dover R, Otto WR. Allogeneic skin substitutes applied to burns patients. <i>Burns</i> . 2002;28(3):254-7.	2
374	Ueda M, Sumi Y, Hibino Y, Hata K. Transplantation of cultured epithelial cells. <i>Methods Mol Biol</i> . 2002;188:197-211.	2
375	Kumagai N. Clinical application of autologous cultured epithelia for the treatment of burns and disfigurement of skin surfaces. <i>Methods Mol Biol</i> . 2002;188:185-96.	6
376	Boyce ST, Warden GD. Principles and practices for treatment of cutaneous wounds with cultured skin substitutes. <i>Am J Surg</i> . 2002;183(4):445-56.	4
377	Dhennin C. [Local treatment of burns]. <i>Pathol Biol (Paris)</i> . 2002;50(2):109-17.	7
378	Boyce ST, Kagan RJ, Yakuboff KP, Meyer NA, Rieman MT, Greenhalgh DG, et al. Cultured skin substitutes reduce donor skin harvesting for closure of excised, full-thickness burns. <i>Ann Surg</i> . 2002;235(2):269-79.	2
379	Shelton J. Cultured autologous keratinocyte sheets: use on large area burns. <i>J Wound Care</i> . 2001;10(8):307-9.	5
380	Donati L. Infection and chemotherapy: recent experiences, proposed strategies and perspectives in plastic and reconstructive surgery. <i>J Chemother</i> . 2001;13 Spec No 1(1):129-33.	6
381	Braye F, Dumortier R, Bertin-Maghit M, Girbon JP, Tissot E, Damour O. [Cultured epidermis for the treatment of severe burns. A 2-year study (18 patients)]. <i>Ann Chir Plast Esthet</i> . 2001;46(6):599-606.	7
382	Rama P, Bonini S, Lambiase A, Golisano O, Paterna P, De Luca M, et al. Autologous fibrin-cultured limbal stem cells permanently restore the corneal surface of patients with total limbal stem cell deficiency. <i>Transplantation</i> . 2001;72(9):1478-85.	1
383	Kremer M, Lang E, Berger A. Organotypical engineering of differentiated composite-skin equivalents of human keratinocytes in a collagen-GAG matrix (INTEGRA Artificial Skin) in a perfusion culture system. <i>Langenbecks Arch Surg</i> . 2001;386(5):357-63.	4
384	Guillot FL. [Production of autologous keratinocytes for therapeutic purposes within a pharmaceutical company]. <i>J Soc Biol</i> . 2001;195(1):75-7.	7
385	Qaryoute S, Mirdad I, Hamail AA. Usage of autograft and allograft skin in treatment of burns in children. <i>Burns</i> . 2001;27(6):599-602.	2
386	Matouskova E, Broz L, Pokorna E, Konigova R. Necrobiotic process causing burn wound conversion may be prevented by allogeneic keratinocytes delivered by the recombined human/pig skin. <i>Folia Biol (Praha)</i> . 2001;47(4):135-42.	4
387	Ojeh NO, Frame JD, Navsaria HA. In vitro characterization of an artificial dermal scaffold. <i>Tissue Eng</i> . 2001;7(4):457-72.	4

연번	서지정보	배제 사유
388	Svensjo T, Yao F, Pomahac B, Eriksson E. Autologous keratinocyte suspensions accelerate epidermal wound healing in pigs. <i>J Surg Res.</i> 2001;99(2):211-21.	4
389	Sheridan RL, Morgan JR, Cusick JL, Petras LM, Lydon MM, Tompkins RG. Initial experience with a composite autologous skin substitute. <i>Burns.</i> 2001;27(5):421-4.	2
390	Kangesu T. Hunterian lecture. Study of dermal grafts and cultured autologous keratinocytes in an experimental model. <i>Ann R Coll Surg Engl.</i> 2001;83(3):197-202.	5
391	Blot SI, Monstrey SJ, Hoste EA, Carsin H, et al. Cultured epithelial autografts in extensive burn coverage of severely traumatized patients: a five year single-center experience with 30 patients. <i>Burns</i> 2000;26:379-387. <i>Burns.</i> 2001;27(4):418-9.	5
392	Lam PK, Chan ES. Integra and cultured epithelium. <i>J Burn Care Rehabil.</i> 2001;22(2):197-8.	5
393	Kuroyanagi Y, Yamada N, Yamashita R, Uchinuma E. Tissue-engineered product: allogeneic cultured dermal substitute composed of spongy collagen with fibroblasts. <i>Artif Organs.</i> 2001;25(3):180-6.	2
394	Navarro FA, Stoner ML, Park CS, Huertas JC, Lee HB, Wood FM, et al. Sprayed keratinocyte suspensions accelerate epidermal coverage in a porcine microwound model. <i>J Burn Care Rehabil.</i> 2000;21(6):513-8.	4
395	Ronfard V, Rives JM, Neveux Y, Carsin H, Barrandon Y. Long-term regeneration of human epidermis on third degree burns transplanted with autologous cultured epithelium grown on a fibrin matrix. <i>Transplantation.</i> 2000;70(11):1588-98.	4
396	Basti S, Rao SK. Current status of limbal conjunctival autograft. <i>Curr Opin Ophthalmol.</i> 2000;11(4):224-32.	6
397	Konigova R, Matouskova E, Broz L. Burn wound coverage and burn wound closure. <i>Acta Chir Plast.</i> 2000;42(2):64-8.	4
398	Bannasch H, Horch RE, Tanczos E, Stark GB. [Treatment of chronic wounds with cultured autologous keratinocytes as suspension in fibrin glue]. <i>Zentralbl Chir.</i> 2000;125 Suppl 1:79-81.	7
399	Stark GB, Bannasch H, Schaefer DJ, Bittner K, Bach A, Voigt M. [Tissue engineering: possibilities and perspectives]. <i>Zentralbl Chir.</i> 2000;125 Suppl 1:69-73.	7
400	Kremer M, Lang E, Berger AC. Evaluation of dermal-epidermal skin equivalents ('composite-skin') of human keratinocytes in a collagen-glycosaminoglycan matrix(Integra artificial skin). <i>Br J Plast Surg.</i> 2000;53(6):459-65.	4
401	Loss M, Wedler V, Kunzi W, Meuli-Simmen C, Meyer VE. Artificial skin, split-thickness autograft and cultured autologous keratinocytes combined to treat a severe burn injury of 93% of TBSA. <i>Burns.</i> 2000;26(7):644-52.	6
402	Tsai RJ, Li LM, Chen JK. Reconstruction of damaged corneas by transplantation of autologous limbal epithelial cells. <i>N Engl J Med.</i> 2000;343(2):86-93.	1
403	Braye F, Pascal P, Bertin-Maghit M, Colpart JJ, Tissot E, Damour O. Advantages of using a bank of allogenic keratinocytes for the rapid coverage of extensive and deep second-degree burns. <i>Med Biol Eng Comput.</i> 2000;38(2):248-52.	2
404	Logsetty S, Heimbach DM. Modern techniques for wound coverage of the thermally injured upper extremity. <i>Hand Clin.</i> 2000;16(2):205-14.	6
405	Braye F, Oddou L, Bertin-Maghit M, Belgacem S, Damour O, Spitalier P, et al. Widely meshed autograft associated with cultured autologous epithelium for the treatment of major burns in children: report of 12 cases. <i>Eur J Pediatr Surg.</i> 2000;10(1):35-40.	2
406	Amshel CE, Caruso DM. Vietnamese "coining": a burn case report and literature review. <i>J Burn Care Rehabil.</i> 2000;21(2):112-4.	6
407	Lamme EN, Van Leeuwen RT, Brandsma K, Van Marle J, Middelkoop E. Higher numbers of autologous fibroblasts in an artificial dermal substitute improve tissue regeneration and modulate scar tissue formation. <i>J Pathol.</i> 2000;190(5):595-603.	4

연번	서지정보	배제 사유
408	Boyce ST, Supp AP, Wickett RR, Hoath SB, Warden GD. Assessment with the dermal torque meter of skin pliability after treatment of burns with cultured skin substitutes. <i>J Burn Care Rehabil.</i> 2000;21(1 Pt 1):55-63.	3
409	Stoner ML, Wood FM. The treatment of hypopigmented lesions with cultured epithelial autograft. <i>J Burn Care Rehabil.</i> 2000;21(1 Pt 1):50-4.	6
410	Supp DM, Supp AP, Bell SM, Boyce ST. Enhanced vascularization of cultured skin substitutes genetically modified to overexpress vascular endothelial growth factor. <i>J Invest Dermatol.</i> 2000;114(1):5-13.	4
411	Supp AP, Wickett RR, Swope VB, Harriger MD, Hoath SB, Boyce ST. Incubation of cultured skin substitutes in reduced humidity promotes cornification in vitro and stable engraftment in athymic mice. <i>Wound Repair Regen.</i> 1999;7(4):226-37.	4
412	Pieptu D, Luchian S. [The skin substitute center--a need in the treatment of patients with severe burns]. <i>Rev Med Chir Soc Med Nat Iasi.</i> 1999;103(1-2):251-5.	7
413	Chistolini P, De Angelis G, De Luca M, Pellegrini G, Ruspantini I. Analysis of the mechanical properties of in vitro reconstructed epidermis: preliminary results. <i>Med Biol Eng Comput.</i> 1999;37(5):670-2.	4
414	Caruso DM, Schuh WH, Al-Kasspooles MF, Chen MC, Schiller WR. Cultured composite autografts as coverage for an extensive body surface area burn: case report and review of the technology. <i>Burns.</i> 1999;25(8):771-9.	6
415	Boyce ST, Kagan RJ, Meyer NA, Yakuboff KP, Warden GD. The 1999 clinical research award. Cultured skin substitutes combined with Integra Artificial Skin to replace native skin autograft and allograft for the closure of excised full-thickness burns. <i>J Burn Care Rehabil.</i> 1999;20(6):453-61.	6
416	Voigt M, Schauer M, Schaefer DJ, Andree C, Horch R, Stark GB. Cultured epidermal keratinocytes on a microspherical transport system are feasible to reconstitute the epidermis in full-thickness wounds. <i>Tissue Eng.</i> 1999;5(6):563-72.	4
417	Lam PK, Chan ES, To EW, Lau CH, Yen SC, King WW. Development and evaluation of a new composite Laserskin graft. <i>J Trauma.</i> 1999;47(5):918-22.	4
418	Hunziker T, Limat A. Cultured keratinocyte grafts. <i>Curr.</i> 1999;27:57-64.	6
419	Pellegrini G, Ranno R, Stracuzzi G, Bondanza S, Guerra L, Zambruno G, et al. The control of epidermal stem cells (holoclones) in the treatment of massive full-thickness burns with autologous keratinocytes cultured on fibrin. <i>Transplantation.</i> 1999;68(6):868-79.	4
420	Aikawa N, Aoki K, Yamazaki M. [Recent advances in the management of severely burned patients]. <i>Nippon Geka Gakkai Zasshi.</i> 1999;100(7):424-9.	7
421	Oreffo RO, Triffitt JT. Future potentials for using osteogenic stem cells and biomaterials in orthopedics. <i>Bone.</i> 1999;25(2 Suppl):5S-9S.	6
422	Boranic M, Jakic-Razumovic J, Stanovic S, Kljenak A, Fattorini I. [Skin cell culture: utilization in plastic surgery and laboratory studies]. <i>Lijec Vjesn.</i> 1999;121(4-5):137-43.	7
423	Takeda T, Tanaka H, Matsuda H, Shimazaki S. Enzyme-free method for cultured skin grafting. <i>J Invest Surg.</i> 1999;12(2):125-7.	4
424	Maggi SP, Soler PM, Smith PD, Hill DP, Ko F, Robson MC. The efficacy of 5% Sulfamylon solution for the treatment of contaminated explanted human meshed skin grafts. <i>Burns.</i> 1999;25(3):237-41.	4
425	Chalumeau M, Saulnier JP, Ainaud P, Lebever H, Stephanazzi J, Lecoadou A, et al. Initial general management and surgery of six extensively burned children treated with cultured epidermal autografts. <i>J Pediatr Surg.</i> 1999;34(4):602-5.	6
426	Chu AJ, Prasad JK. Up-regulation by human recombinant transforming growth factor beta-1 of collagen production in cultured dermal fibroblasts is mediated by the inhibition of nitric oxide signaling. <i>J Am Coll Surg.</i> 1999;188(3):271-80.	4



연번	서지정보	배제 사유
427	Osborne CS, Reid WH, Grant MH. Investigation into the biological stability of collagen/chondroitin-6-sulphate gels and their contraction by fibroblasts and keratinocytes: the effect of crosslinking agents and diamines. <i>Biomaterials</i> . 1999;20(3):283-90.	4
428	Harris PA, di Francesco F, Barisoni D, Leigh IM, Navsaria HA. Use of hyaluronic acid and cultured autologous keratinocytes and fibroblasts in extensive burns. <i>Lancet</i> . 1999;353(9146):35-6.	5
429	Boyce ST, Supp AP, Swope VB, Warden GD. Topical sulfamylon reduces engraftment of cultured skin substitutes on athymic mice. <i>J Burn Care Rehabil</i> . 1999;20(1 Pt 1):33-6.	4
430	Duquesnoy RJ, Liu K, Fu XF, Murase N, Ye Q, Demetris AJ. Evidence for heat shock protein immunity in a rat cardiac allograft model of chronic rejection. <i>Transplantation</i> . 1999;67(1):156-64.	4
431	Pellegrini G, Bondanza S, Guerra L, De Luca M. Cultivation of human keratinocyte stem cells: current and future clinical applications. <i>Med Biol Eng Comput</i> . 1998;36(6):778-90.	6
432	Boyce ST. Skin substitutes from cultured cells and collagen-GAG polymers. <i>Med Biol Eng Comput</i> . 1998;36(6):791-800.	4
433	Bozinko GM, Lowe KE, Reigart CL. A new option for burn victims. <i>Rn</i> . 1998;61(11):37-9.	5
434	Kirsner RS. The use of Apligraf in acute wounds. <i>J Dermatol</i> . 1998;25(12):805-11.	6
435	Compton CC, Nadire KB, Regauer S, Simon M, Warland G, O'Connor NE, et al. Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation. <i>Differentiation</i> . 1998;64(1):45-53.	3
436	van Zuijlen PP, van Leeuwen RT, Middelkoop E. Practical sources for autologous fibroblasts to prepare a bioengineered dermal equivalent. <i>Burns</i> . 1998;24(7):687.	2
437	Sheridan R, Choucair R, Donelan M, Lydon M, Petras L, Tompkins R. Acellular allodermis in burns surgery: 1-year results of a pilot trial. <i>J Burn Care Rehabil</i> . 1998;19(6):528-30.	2
438	Black AF, Berthod F, L'Heureux N, Germain L, Auger FA. In vitro reconstruction of a human capillary-like network in a tissue-engineered skin equivalent. <i>Faseb J</i> . 1998;12(13):1331-40.	4
439	Soejima K, Nozaki M, Kobayashi M, Negishi N. Studies of surface microarchitecture using a hand-held video microscope in cases of cultured epithelial autografts. <i>Ann Plast Surg</i> . 1998;41(3):270-4.	3
440	Pandya AN, Woodward B, Parkhouse N. The use of cultured autologous keratinocytes with integra in the resurfacing of acute burns. <i>Plast Reconstr Surg</i> . 1998;102(3):825-8; discussion 9-30.	6
441	Matsumura H, Meyer NA, Mann R, Heimbach DM. Melting graft-wound syndrome. <i>J Burn Care Rehabil</i> . 1998;19(4):292-5.	2
442	Kumagai N. [Clinical application of cultured epithelium]. <i>Hum Cell</i> . 1998;11(1):61-8.	7
443	Orgill DP, Butler C, Regan JF, Barlow MS, Yannas IV, Compton CC. Vascularized collagen-glycosaminoglycan matrix provides a dermal substrate and improves take of cultured epithelial autografts. <i>Plast Reconstr Surg</i> . 1998;102(2):423-9.	4
444	Hunt JP, Hunter CT, Brownstein M, Hultman CS, deSerres S, Bracey L, et al. Host priming, not target antigen type, decides rejection rate in mice primed with MHC II "knockout" cultured keratinocytes. <i>J Surg Res</i> . 1998;76(1):32-6.	4
445	Hultman CS, Hunt JP, Yamamoto H, Giannopoulos A, deSerres S, Frelinger JA, et al. Immunogenicity of cultured keratinocyte allografts deficient in major histocompatibility complex antigens. <i>J Trauma</i> . 1998;45(1):25-33; discussion -4.	4
446	Garner WL. Epidermal regulation of dermal fibroblast activity. <i>Plast Reconstr Surg</i> . 1998;102(1):135-9.	4
447	Kirsner RS, Falanga V, Eaglstein WH. The development of bioengineered skin. <i>Trends Biotechnol</i> . 1998;16(6):246-9.	6

연번	서지정보	배제 사유
448	Coulomb B, Friteau L, Baruch J, Guilbaud J, Chretien-Marquet B, Glicenstein J, et al. Advantage of the presence of living dermal fibroblasts within in vitro reconstructed skin for grafting in humans. <i>Plast Reconstr Surg.</i> 1998;101(7):1891-903.	4
449	Osborne CS, Barbenel JC, Smith D, Savakis M, Grant MH. Investigation into the tensile properties of collagen/chondroitin-6-sulphate gels: the effect of crosslinking agents and diamines. <i>Med Biol Eng Comput.</i> 1998;36(1):129-34.	4
450	Wright KA, Nadire KB, Busto P, Tubo R, McPherson JM, Wentworth BM. Alternative delivery of keratinocytes using a polyurethane membrane and the implications for its use in the treatment of full-thickness burn injury. <i>Burns.</i> 1998;24(1):7-17.	4
451	Benathan M, Labidi-Ubaldi F. [Living epidermal and dermal substitutes for treatment of severely burned patients]. <i>Rev Med Suisse Romande.</i> 1998;118(2):149-53.	7
452	Pages JC, Benathan M, Imhof M, Deglise B, Favarger N. [Treatment of the severely burned patient with cultured autologous epidermis: the experience at Lausanne]. <i>Rev Med Suisse Romande.</i> 1998;118(2):145-8.	7
453	Tanaka H, Takami Y, Tokunaga T, Takeda T, Shimazaki S. [Current treatment burn wounds: alternative wound coverage methods]. <i>Nippon Geka Gakkai Zasshi.</i> 1998;99(1):14-20.	7
454	Eming SA, Medalie DA, Tompkins RG, Yarmush ML, Morgan JR. Genetically modified human keratinocytes overexpressing PDGF-A enhance the performance of a composite skin graft. <i>Hum Gene Ther.</i> 1998;9(4):529-39.	4
455	Haith LR, Jr., Moss BE, Kraut JD, Patton ML, Germain TJ. Trilaminar skin coverage combined with cultured epithelial autografts--a new technique for treating the elderly patient with burn injuries. <i>J Burn Care Rehabil.</i> 1998;19(1 Pt 1):59-61.	6
456	Klein L, Mericka P, Strakova H, Jebavy L, Nozickova M, Blaha M, et al. Biological skin covers in treatment of two cases of the Lyell's syndrome. <i>Ann Transplant.</i> 1997;2(1):45-8.	1
457	Wang X, Wang J, Wu J. [Manufacture and application of a new composite allograft]. <i>Chung Kuo Hsiu Fu Chung Chien Wai Ko Tsa Chih.</i> 1997;11(2):100-2.	7
458	Carsin H, Ainaud P, Le Bever H, Rives JM, Le Coadou A, Stephanazzi J. [Objectives, results and future prospects of burn treatment in 1997]. <i>Bull Acad Natl Med.</i> 1997;181(7):1307-19; discussion 19-20.	7
459	Malakhov SF, Paramonov BA, Emel'ianov AV, Vasil'ev AV, Terskikh VV. [New approaches to the treatment of severe burns: the transplantation of keratinocytes grown in culture]. <i>Voен Med Zh.</i> 1997;318(9):16-9, 80.	7
460	Sheridan RL, Choucair RJ. Acellular allogenic dermis does not hinder initial engraftment in burn wound resurfacing and reconstruction. <i>J Burn Care Rehabil.</i> 1997;18(6):496-9.	2
461	Raghunath M, Meuli M. Cultured epithelial autografts: diving from surgery into matrix biology. <i>Pediatr Surg Int.</i> 1997;12(7):478-83.	5
462	Meuli M, Raghunath M. Burns (Part 2). Tops and flops using cultured epithelial autografts in children. <i>Pediatr Surg Int.</i> 1997;12(7):471-7.	6
463	Paddle-Ledinek JE, Cruickshank DG, Masterton JP. Skin replacement by cultured keratinocyte grafts: an Australian experience. <i>Burns.</i> 1997;23(3):204-11.	6
464	Matouskova E, Bucek S, Vogtova D, Vesely P, Chaloupkova A, Broz L, et al. Treatment of burns and donor sites with human allogeneic keratinocytes grown on acellular pig dermis. <i>Br J Dermatol.</i> 1997;136(6):901-7.	2
465	Schlotterer M. [Value of cutaneous allografts in the treatment of severe burns]. <i>Chirurgie.</i> 1997;122(1):6-8; discussion 9.	7
466	Myers SR, Grady J, Soranzo C, Sanders R, Green C, Leigh IM, et al. A hyaluronic acid membrane delivery system for cultured keratinocytes: clinical "take" rates in the porcine kerato-dermal model. <i>J Burn Care Rehabil.</i> 1997;18(3):214-22.	4

연번	서지정보	배제 사유
467	Kumagai N, Oshima H, Tanabe M, Ishida H, Uchikoshi T. Favorable donor site for epidermal cultivation for the treatment of burn scars with autologous cultured epithelium. <i>Ann Plast Surg.</i> 1997;38(5):506-13.	1
468	Janezic TF. Burns following petrol sniffing. <i>Burns.</i> 1997;23(1):78-80.	6
469	Munster AM. Whither [corrected] skin replacement? <i>Burns.</i> 1997;23(1):v.	5
470	Pellegrini G, Traverso CE, Franzi AT, Zingirian M, Cancedda R, De Luca M. Long-term restoration of damaged corneal surfaces with autologous cultivated corneal epithelium. <i>Lancet.</i> 1997;349(9057):990-3.	1
471	Tsai CY, Ueda M, Hata K, Horie K, Hibino Y, Sugimura Y, et al. Clinical results of cultured epithelial cell grafting in the oral and maxillofacial region. <i>J Craniomaxillofac Surg.</i> 1997;25(1):4-8.	1
472	Purdue GF, Hunt JL, Still JM, Jr., Law EJ, Herndon DN, Goldfarb IW, et al. A multicenter clinical trial of a biosynthetic skin replacement, Dermagraft-TC, compared with cryopreserved human cadaver skin for temporary coverage of excised burn wounds. <i>J Burn Care Rehabil.</i> 1997;18(1 Pt 1):52-7.	2
473	Hansbrough JF, Mozingo DW, Kealey GP, Davis M, Gidner A, Gentzkow GD. Clinical trials of a biosynthetic temporary skin replacement, Dermagraft-Transitional Covering, compared with cryopreserved human cadaver skin for temporary coverage of excised burn wounds. <i>J Burn Care Rehabil.</i> 1997;18(1 Pt 1):43-51.	4
474	Kakibuchi M, Hosokawa K, Fujikawa M, Yoshikawa K. The use of cultivated epidermal cell sheets in skin grafting. <i>J Wound Care.</i> 1996;5(10):487-90.	6
475	Spilker G, Reifenrath MW, Kaiser HW. [Skin cultivation in treatment of severely burned patients]. <i>Langenbecks Arch Chir Suppl Kongressbd.</i> 1996;113:1149-55.	7
476	Raff T, Hartmann B, Wagner H, Germann G. Experience with the modified Meek technique. <i>Acta Chir Plast.</i> 1996;38(4):142-6.	2
477	Nunez-Gutierrez H, Castro-Munozledo F, Kuri-Harcuch W. Combined use of allograft and autograft epidermal cultures in therapy of burns. <i>Plast Reconstr Surg.</i> 1996;98(6):929-39; discussion 40-1.	2
478	Caruso DM, Gregory MW, Schiller WR. The use of skin from a monozygotic twin combined with cultured epithelial autografts as coverage for a large surface area burn: a case report and review of the literature. <i>J Burn Care Rehabil.</i> 1996;17(5):432-4.	6
479	Still J, Scheirer R, Law E. Acute cholecystectomy performed through cultured epithelial autografts in a patient with burn injuries: a case report. <i>J Burn Care Rehabil.</i> 1996;17(5):429-31.	6
480	Holder IA, Boyce ST. Formulation of 'idealized' topical antimicrobial mixtures for use with cultured skin grafts. <i>J Antimicrob Chemother.</i> 1996;38(3):457-63.	4
481	Wood FM, Stoner M. Implication of basement membrane development on the underlying scar in partial-thickness burn injury. <i>Burns.</i> 1996;22(6):459-62.	6
482	Wood F, Liddiard K, Skinner A, Ballentyne J. Scar management of cultured epithelial autograft. <i>Burns.</i> 1996;22(6):451-4.	6
483	Somers T, Verbeken G, Vanhalle S, Delaey B, Duinslaeger L, Govaerts P, et al. Lysates from cultured allogeneic keratinocytes stimulate wound healing after tympanoplasty. <i>Acta Otolaryngol (Stockh).</i> 1996;116(4):589-93.	2
484	Xu W, Germain L, Goulet F, Auger FA. Permanent grafting of living skin substitutes: surgical parameters to control for successful results. <i>J Burn Care Rehabil.</i> 1996;17(1):7-13.	4
485	Raghunath M, Hopfner B, Aeschlimann D, Luthi U, Meuli M, Altermatt S, et al. Cross-linking of the dermo-epidermal junction of skin regenerating from keratinocyte autografts. Anchoring fibrils are a target for tissue transglutaminase. <i>J Clin Invest.</i> 1996;98(5):1174-84.	4

연번	서지정보	배제 사유
486	Vicanova J, Mommaas AM, Mulder AA, Koerten HK, Ponec M. Impaired desquamation in the in vitro reconstructed human epidermis. <i>Cell Tissue Res.</i> 1996;286(1):115-22.	5
487	Sheridan RL, Tompkins RG. Recent clinical experience with cultured autologous epithelium. <i>Br J Plast Surg.</i> 1996;49(1):72-4.	5
488	Rennekampff HO, Hansbrough JF, Kiessig V, Abiezzi S, Woods V, Jr. Wound closure with human keratinocytes cultured on a polyurethane dressing overlaid on a cultured human dermal replacement. <i>Surgery.</i> 1996;120(1):16-22.	4
489	Hultman CS, Brinson GM, Siltharm S, deSerres S, Cairns BA, Peterson HD, et al. Allogeneic fibroblasts used to grow cultured epidermal autografts persist in vivo and sensitize the graft recipient for accelerated second-set rejection. <i>J Trauma.</i> 1996;41(1):51-8; discussion 8-60.	4
490	Duinslaeger L, Verbeken G, Reper P, Delaey B, Vanhalle S, Vanderkelen A. Lyophilized keratinocyte cell lysates contain multiple mitogenic activities and stimulate closure of meshed skin autograft-covered burn wounds with efficiency similar to that of fresh allogeneic keratinocyte cultures. <i>Plast Reconstr Surg.</i> 1996;98(1):110-7.	3
491	Dawson RA, Goberdhan NJ, Freedlander E, MacNeil S. Influence of extracellular matrix proteins on human keratinocyte attachment, proliferation and transfer to a dermal wound model. <i>Burns.</i> 1996;22(2):93-100.	4
492	Erdmann D, Hussmann J, Kucan JO. Treatment of a severe alkali burn. <i>Burns.</i> 1996;22(2):141-6.	6
493	Goulet F, Poitras A, Rouabhia M, Cusson D, Germain L, Auger FA. Stimulation of human keratinocyte proliferation through growth factor exchanges with dermal fibroblasts in vitro. <i>Burns.</i> 1996;22(2):107-12.	4
494	Raghunath M, Bachi T, Meuli M, Altermatt S, Gobet R, Bruckner-Tuderman L, et al. Fibrillin and elastin expression in skin regenerating from cultured keratinocyte autografts: morphogenesis of microfibrils begins at the dermo-epidermal junction and precedes elastic fiber formation. <i>J Invest Dermatol.</i> 1996;106(5):1090-5.	4
495	Eming SA, Yarmush ML, Morgan JR. Enhanced function of cultured epithelium by genetic modification: Cell-based synthesis and delivery of growth factors. <i>Biotechnol Bioeng.</i> 1996;52(1):15-23.	4
496	Gallico GG, O'Connor NE. Engineering a skin replacement. <i>Tissue Eng.</i> 1995;1(3):231-40.	5
497	Goretsky MJ, Supp AP, Greenhalgh DG, Warden GD, Boyce ST. Surface electrical capacitance as an index of epidermal barrier properties of composite skin substitutes and skin autografts. <i>Wound Repair Regen.</i> 1995;3(4):419-25.	3
498	Putland M, Snelling CF, Macdonald I, Tron VA. Histologic comparison of cultured epithelial autograft and meshed expanded split-thickness skin graft. <i>J Burn Care Rehabil.</i> 1995;16(6):627-40.	3
499	Neveux Y, Rives JM, Le Breton C, Gentilhomme E, Saint-Blancar P, Carsin H. Clinical interest of cutaneous models reproduced in vitro for severe burn treatment: histopathological and ultrastructural study. <i>Cell Biol Toxicol.</i> 1995;11(3-4):173-8.	3
500	Wang HJ, Chen TM, Cheng LF, Cheng TY, Tung YM. Human keratinocyte culture using porcine pituitary extract in serum-free medium. <i>Burns.</i> 1995;21(7):503-6.	4
501	Matsuzaki K, Kumagai N, Fukushi S, Ohshima H, Tanabe M, Ishida H. Cultured epithelial autografting on meshed skin graft scars: evaluation of skin elasticity. <i>J Burn Care Rehabil.</i> 1995;16(5):496-502.	6
502	Boyce ST, Goretsky MJ, Greenhalgh DG, Kagan RJ, Rieman MT, Warden GD. Comparative assessment of cultured skin substitutes and native skin autograft for treatment of full-thickness burns. <i>Ann Surg.</i> 1995;222(6):743-52.	2

연번	서지정보	배제 사유
503	Harriger MD, Warden GD, Greenhalgh DG, Kagan RJ, Boyce ST. Pigmentation and microanatomy of skin regenerated from composite grafts of cultured cells and biopolymers applied to full-thickness burn wounds. <i>Transplantation</i> . 1995;59(5):702-7.	4
504	Boyce ST, Supp AP, Harriger MD, Greenhalgh DG, Warden GD. Topical nutrients promote engraftment and inhibit wound contraction of cultured skin substitutes in athymic mice. <i>J Invest Dermatol</i> . 1995;104(3):345-9.	4
505	Rouabhia M, Germain L, Bergeron J, Auger FA. Allogeneic-syngeneic cultured epithelia. A successful therapeutic option for skin regeneration. <i>Transplantation</i> . 1995;59(9):1229-35.	4
506	Henckel von Donnersmarck G, Muhlbauer W, Hofter E, Hartinger A. [Use of keratinocyte cultures in treatment of severe burns--experiences up to now, outlook for further subsequent developments]. <i>Unfallchirurg</i> . 1995;98(4):229-32.	7
507	Pitzler D, Bisgwa F, Partecke BD. [Basic comments on differential surgical therapy of severe burns]. <i>Unfallchirurg</i> . 1995;98(4):174-9.	7
508	Cairns BA, deSerres S, Brady LA, Hultman CS, Meyer AA. Xenogeneic mouse fibroblasts persist in human cultured epidermal grafts: a possible mechanism of graft loss. <i>J Trauma</i> . 1995;39(1):75-9; discussion 9-80.	4
509	Muhlbauer W, Henckel von Donnersmarck G, Hofter E, Hartinger A. [Keratinocyte culture and transplantation in burns]. <i>Chirurg</i> . 1995;66(4):271-6.	7
510	Germann G, Raff T. [Homograft transplantation in severely burned patients. Principles, indications and possibilities]. <i>Chirurg</i> . 1995;66(4):260-70.	7
511	Wong L. The many uses of allograft skin. <i>Ostomy Wound Manage</i> . 1995;41(4):36-8, 40-2.	5
512	Rives JM, Cantaloube D, Ainaud P, Barandon Y, Carsin H. [Role of autografts of cultured epidermis in the treatment of deep burns of the face. Preliminary results]. <i>Ann Chir Plast Esthet</i> . 1995;40(3):286-92.	7
513	Hultman CS, Cairns BA, deSerres S, Frelinger JA, Meyer AA. Burn injury impairs second-set rejection and CTL reactivity in mice primed by cultured keratinocyte allografts. <i>Transplantation</i> . 1995;60(6):584-9.	4
514	Lopez Gutierrez JC, Ros Z, Vallejo D, Perdiguero M, Soto C, Tovar J. Cultured epidermal autograft in the management of critical pediatric burn patients. <i>Eur J Pediatr Surg</i> . 1995;5(3):174-6.	6
515	Damour O, Gueugniaud PY, Berthin-Maghit M, Rousselle P, Berthod F, Sahuc F, et al. A dermal substrate made of collagen--GAG--chitosan for deep burn coverage: first clinical uses. <i>Clin Mater</i> . 1994;15(4):273-6.	6
516	Teepe RG, Burger A, Ponec M. Immunohistochemical studies on regeneration in cultured epidermal autografts used to treat full-thickness burn wounds. <i>Clin Exp Dermatol</i> . 1994;19(1):16-22.	4
517	Hickerson WL, Compton C, Fletchall S, Smith LR. Cultured epidermal autografts and allodermis combination for permanent burn wound coverage. <i>Burns</i> . 1994;20 Suppl 1:S52-5; discussion S5-6.	6
518	Stark GB, Kaiser HW. Cologne Burn Centre experience with glycerol-preserved allogeneic skin: Part II: Combination with autologous cultured keratinocytes. <i>Burns</i> . 1994;20 Suppl 1:S34-8.	6
519	Moghtader JC, Edlich RF, Mintz PD, Zachmann GC, Himel HN. The use of recombinant human erythropoietin and cultured epithelial autografts in a Jehovah's Witness with a major thermal injury. <i>Burns</i> . 1994;20(2):176-7.	6
520	Hansbrough JF, Morgan J, Greenleaf G, Underwood J. Development of a temporary living skin replacement composed of human neonatal fibroblasts cultured in Biobrane, a synthetic dressing material. <i>Surgery</i> . 1994;115(5):633-44.	4

연번	서지정보	배제 사유
521	Kaiser HW, Stark GB, Kopp J, Balcerkiewicz A, Spilker G, Kreysel HW. Cultured autologous keratinocytes in fibrin glue suspension, exclusively and combined with STS-allograft (preliminary clinical and histological report of a new technique). Burns. 1994;20(1):23-9.	6
522	Krupp S, Wiesner L, Krstic R, Pescia G, Winistorfer B. Mid-term results with cultured epidermal autografts, allogenic skin transplants and cyclosporin A medication. Burns. 1994;20(1):15-20.	6
523	Malakhov SF, Paramonov BA, Vasiliev AV, Terskikh VV. Preliminary report of the clinical use of cultured allogeneic keratinocytes. Burns. 1994;20(5):463-6.	6
524	Angres G, Maier H, Maier K, Weidauer H. [The use of autologous keratinocytes in ear surgery]. Hno. 1994;42(9):565-9.	7
525	Petzoldt JL, Leigh IM, Duffy PG, Masters JR. Culture and characterisation of human urothelium in vivo and in vitro. Urol Res. 1994;22(2):67-74.	4
526	Fratianne R, Papay F, Housini I, Lang C, Schafer IA. Keratinocyte allografts accelerate healing of split-thickness donor sites: applications for improved treatment of burns. J Burn Care Rehabil. 1993;14(2 Pt 1):148-54.	2
527	Rue LW, 3rd, Cioffi WG, McManus WF, Pruitt BA, Jr. Wound closure and outcome in extensively burned patients treated with cultured autologous keratinocytes. J Trauma. 1993;34(5):662-7; discussion 7-8.	6
528	Walter PH. Burn wound management. AACN Clin Issues Crit Care Nurs. 1993;4(2):378-87.	6
529	Boyce ST, Greenhalgh DG, Kagan RJ, Housinger T, Sorrell JM, Childress CP, et al. Skin anatomy and antigen expression after burn wound closure with composite grafts of cultured skin cells and biopolymers. Plast Reconstr Surg. 1993;91(4):632-41.	3
530	Bolgiani A, Patino O, Benaim F. The use of the modified Thomas splint for a patient with cultured epidermal autografts. J Burn Care Rehabil. 1993;14(4):466-70.	6
531	Malakhov SF, Terskikh VV, Bautin EA, Vasil'ev AV, Paramonov BA. [The autotransplantation of keratinocytes grown outside the body to treat extensive burns]. Vestn Khir Im I I Grek. 1993;150(3-4):59-62.	7
532	Yadav JK, Singhvi AM, Kumar N, Garg S. Topical phenytoin in the treatment of split-thickness skin autograft donor sites: a comparative study with polyurethane membrane drape and conventional dressing. Burns. 1993;19(4):306-10.	2
533	Brychta P, Michalek V, Adler J, Rihova H, Suchanek I. [Use of cultured epidermal autotransplantation in a severely burned patient]. Rozhl Chir. 1993;72(5):216-8.	7
534	Cairns BA, deSerres S, Kilpatrick K, Frelinger JA, Meyer AA. Cultured keratinocyte allografts fail to induce sensitization in vivo. Surgery. 1993;114(2):416-22.	4
535	Compton CC, Hickerson W, Nadire K, Press W. Acceleration of skin regeneration from cultured epithelial autografts by transplantation to homograft dermis. J Burn Care Rehabil. 1993;14(6):653-62.	3
536	Kuroyanagi Y, Kenmochi M, Ishihara S, Takeda A, Shiraishi A, Ootake N, et al. A cultured skin substitute composed of fibroblasts and keratinocytes with a collagen matrix: preliminary results of clinical trials. Ann Plast Surg. 1993;31(4):340-9; discussion 9-51.	6
537	Burns BF, McCauley RL, Murphy FL, Robson MC. Reconstructive management of patients with greater than 80 per cent TBSA burns. Burns. 1993;19(5):429-33.	3
538	Sarkisov DS, Alekseev AA, Tumanov VP, Glushchenko EV, Morozov SS, Paltsyn AA. [Cultured human skin cells in the treatment of burns]. Khirurgiia (Mosk). 1993(3):22-7.	7
539	Soeda J, Inokuchi S, Ueno S, Yokoyama S, Kidokoro M, Nakamura Y, et al. Use of cultured human epidermal allografts for the treatment of extensive partial thickness scald burn in children. Tokai J Exp Clin Med. 1993;18(1-2):65-70.	2

연번	서지정보	배제 사유
540	Trout RJ, Soltis MB. Collaborative atmosphere: rectal pouching in the burn center. J ET Nurs. 1992;19(1):24-7.	6
541	Arons JA, Wainwright DJ, Jordon RE. The surgical applications and implications of cultured human epidermis: a comprehensive review. Surgery. 1992;111(1):4-11.	6
542	Kumicakova E, Kumicak J. [Culture of skin grafts in vitro]. Cas Lek Cesk. 1992;131(10):301-3.	7
543	Nanchahal J, Ward CM. New grafts for old? A review of alternatives to autologous skin. Br J Plast Surg. 1992;45(5):354-63.	6
544	Mommaas AM, Teepe RG, Leigh IM, Mulder AA, Koebrugge EJ, Vermeer BJ. Ontogenesis of the basement membrane zone after grafting cultured human epithelium: a morphologic and immunoelectron microscopic study. J Invest Dermatol. 1992;99(1):71-7.	3
545	Odessey R. Addendum: multicenter experience with cultured epidermal autograft for treatment of burns. J Burn Care Rehabil. 1992;13(1):174-80.	5
546	Barillo DJ, Nangle ME, Farrell K. Preliminary experience with cultured epidermal autograft in a community hospital burn unit. J Burn Care Rehabil. 1992;13(1):158-65.	6
547	Herndon DN, Rutan RL. Comparison of cultured epidermal autograft and massive excision with serial autografting plus homograft overlay. J Burn Care Rehabil. 1992;13(1):154-7.	6
548	Nave M. Wound bed preparation: approaches to replacement of dermis. J Burn Care Rehabil. 1992;13(1):147-53.	6
549	Haith LR, Jr., Patton ML, Goldman WT. Cultured epidermal autograft and the treatment of the massive burn injury. J Burn Care Rehabil. 1992;13(1):142-6.	6
550	Weekley R, Klein R. Clinical nursing experience with cultured epidermal autografts. J Burn Care Rehabil. 1992;13(1):138-41.	6
551	Heimbach DM. A nonuser's questions about cultured epidermal autograft. J Burn Care Rehabil. 1992;13(1):127-9.	5
552	Munster AM. Use of cultured epidermal autograft in ten patients. J Burn Care Rehabil. 1992;13(1):124-6.	8
553	Zhao YB, Zhao XF, Li A, Lu SZ, Wang X, Huang SZ, et al. Clinical observations and methods for identifying the existence of cultured epidermal allografts. Burns. 1992;18(1):4-8.	2
554	Donati L, Magliacani G, Bormioli M, Signorini M, Preis FW. Clinical experiences with keratinocyte grafts. Burns. 1992;18 Suppl 1:S19-26.	5
555	De Luca M, Bondanza S, Cancedda R, Tamisani AM, Di Noto C, Muller L, et al. Permanent coverage of full skin thickness burns with autologous cultured epidermis and reepithelialization of partial skin thickness lesions induced by allogeneic cultured epidermis: a multicentre study in the treatment of children. Burns. 1992;18 Suppl 1:S16-9.	6
556	Coleman JJ, 3rd, Siwy BK. Cultured epidermal autografts: a life-saving and skin-saving technique in children. J Pediatr Surg. 1992;27(8):1029-32.	6
557	Zhao YB. [Primary observation of prolonged survival of cultured epidermal allografts]. Chung Hua Wai Ko Tsa Chih. 1992;30(2):104-6, 25-6.	7
558	Tamisani AM, Ferretti S, Sangiorgio L. Critical reflections on the use of human cultured keratinocytes in children with burns. Eur J Pediatr Surg. 1992;2(4):223-6.	2
559	Compton CC. Current concepts in pediatric burn care: the biology of cultured epithelial autografts: an eight-year study in pediatric burn patients. Eur J Pediatr Surg. 1992;2(4):216-22.	3
560	Krupp S, Benathan M, Meuli M, Deglise B, Holzer E, Wiesner L, et al. Current concepts in pediatric burn care: management of burn wounds with cultured epidermal autografts. Eur J Pediatr Surg. 1992;2(4):210-5.	6

연번	서지정보	배제 사유
561	Wilson JL, Dollard SC, Chow LT, Broker TR. Epithelial-specific gene expression during differentiation of stratified primary human keratinocyte cultures. <i>Cell Growth Differ.</i> 1992;3(8):471-83.	4
562	Bideaux JC, Echinard C, Damour O. [Experimental study on a skin substitute. Artificial dermis epidermised by human keratinocytes]. <i>Chirurgie.</i> 1992;118(6-7):411-5.	7
563	Phillips TJ, Gilchrest BA. Clinical applications of cultured epithelium. <i>Epithelial Cell Biol.</i> 1992;1(1):39-46.	6
564	Seah CS. Skin graft and skin equivalent in burns. <i>Ann Acad Med Singapore.</i> 1992;21(5):685-8.	6
565	Tompkins RG, Burke JF. Burn wound closure using permanent skin replacement materials. <i>World J Surg.</i> 1992;16(1):47-52.	5
566	Krejci NC, McGuire J. Treatment of burns with skin substitutes. <i>J Dermatol Sci.</i> 1992;4(3):149-55.	6
567	Sarkisov DS, Morozov SS, Tumanov VP. [The modern method of treating burn wounds]. <i>Voen Med Zh.</i> 1991(7):55-6.	7
568	Boyce ST, Foreman TJ, English KB, Stayner N, Cooper ML, Sakabu S, et al. Skin wound closure in athymic mice with cultured human cells, biopolymers, and growth factors. <i>Surgery.</i> 1991;110(5):866-76.	4
569	Carver N, Leigh IM. Keratinocyte grafts and skin equivalents. <i>Int J Dermatol.</i> 1991;30(8):540-51.	6
570	Prunieras M. To reconstitute skin: theme and variations. <i>Matrix.</i> 1991;11(4):302-5.	6
571	Fabre JW. Epidermal allografts. <i>Immunol Lett.</i> 1991;29(1-2):161-5.	6
572	Phillips TJ, Gilchrest BA. Cultured epidermal allografts as biological wound dressings. <i>Prog Clin Biol Res.</i> 1991;365:77-94.	5
573	Randall T. Cultured skin cells, artificial dermis join in race against time to cover burn wound. <i>Jama.</i> 1991;265(15):1918-9.	5
574	Leigh IM, Navsaria H, Purkis PE, McKay I. Clinical practice and biological effects of keratinocyte grafting. <i>Ann Acad Med Singapore.</i> 1991;20(4):549-55.	5
575	Schaller E, Meyer H, Mailander P, Berger A. [Acute and definitive management of burn wounds with cultured keratinocytes and free tissue transfer]. <i>Langenbecks Arch Chir Suppl Kongressbd.</i> 1991:623-7.	7
576	Klein P. [Skin culture on the burn wound as a final skin replacement]. <i>Langenbecks Arch Chir Suppl Kongressbd.</i> 1991:617-21.	7
577	Desai MH, Mlakar JM, McCauley RL, Abdullah KM, Rutan RL, Waymack JP, et al. Lack of long-term durability of cultured keratinocyte burn-wound coverage: a case report. <i>J Burn Care Rehabil.</i> 1991;12(6):540-5.	6
578	Clugston PA, Snelling CF, Macdonald IB, Maledy HL, Boyle JC, Germann E, et al. Cultured epithelial autografts: three years of clinical experience with eighteen patients. <i>J Burn Care Rehabil.</i> 1991;12(6):533-9.	6
579	Phipps AR, Clarke JA. The use of intermingled autograft and parental allograft skin in the treatment of major burns in children. <i>Br J Plast Surg.</i> 1991;44(8):608-11.	2
580	Angelini Rota M, Mezzetti E, Sabatini G, Monti M. [Human allogeneic cultured epidermis: medico-legal considerations]. <i>Ann Ital Chir.</i> 1991;62(4):369-72; discussion 72-3.	7
581	Greenleaf G, Cooper ML, Hansbrough JF. Microbial contamination in allografted wound beds in patients with burns. <i>J Burn Care Rehabil.</i> 1991;12(5):442-5.	2
582	Krejci NC, Cuono CB, Langdon RC, McGuire J. In vitro reconstitution of skin: fibroblasts facilitate keratinocyte growth and differentiation on acellular reticular dermis. <i>J Invest Dermatol.</i> 1991;97(5):843-8.	4



연번	서지정보	배제 사유
583	Oliver AM, Kaawach W, Mithoff EW, Watt A, Abramovich DR, Rayner CR. The differentiation and proliferation of newly formed epidermis on wounds treated with cultured epithelial allografts. <i>Br J Dermatol.</i> 1991;125(2):147-54.	2
584	de Luca M, Albanese E, Megna M, Cancedda R, Mangiante PE, Cadoni A, et al. Evidence that human oral epithelium reconstituted in vitro and transplanted onto patients with defects in the oral mucosa retains properties of the original donor site. <i>Transplantation.</i> 1990;50(3):454-9.	4
585	Yarbrough DR, 3rd. Improving survival in the burned patient. <i>J S C Med Assoc.</i> 1990;86(6):347-9.	5
586	Phillips TJ, Bhawan J, Gilchrest BA. Characterization of cellular elements in healed cultured keratinocyte autografts used to cover burn wounds. <i>Arch Dermatol.</i> 1990;126(8):1104-5.	5
587	Munster AM, Weiner SH, Spence RJ. Cultured epidermis for the coverage of massive burn wounds. A single center experience. <i>Ann Surg.</i> 1990;211(6):676-9; discussion 9-80.	8
588	Teepe RG, Kreis RW, Koebrugge EJ, Kempenaar JA, Vloemans AF, Hermans RP, et al. The use of cultured autologous epidermis in the treatment of extensive burn wounds. <i>J Trauma.</i> 1990;30(3):269-75.	2
589	Itin P, Pittelkow MR. [Autologous epithelium transplantation using cultured keratinocytes. History and current role in burn treatment]. <i>Helv Chir Acta.</i> 1990;57(3):519-28.	7
590	Elberg JJ, Siemssen SJ, Nielsen EH, Ronne M. Grafting of human autogenous keratinocytes cultured in vitro. <i>In Vivo.</i> 1990;4(6):367-9.	6
591	Dioguardi D, Barbuti S, Calvario A, Brienza E, Larocca AM. Assessment of disinfectant topical agents in keratinocyte cell culture grafting technique in severely burned patients. <i>Acta Chir Plast.</i> 1990;32(4):218-24.	3
592	Compton CC, Regauer S, Seiler GR, Landry DB. Human Merkel cell regeneration in skin derived from cultured keratinocyte grafts. <i>Lab Invest.</i> 1990;63(2):233-41.	4
593	Woodley DT, Briggaman RA, Herzog SR, Meyers AA, Peterson HD, O'Keefe EJ. Characterization of "neo-dermis" formation beneath cultured human epidermal autografts transplanted on muscle fascia. <i>J Invest Dermatol.</i> 1990;95(1):20-6.	4
594	Foyatier JL, Faure M, Hezez G, Masson C, Paulus C, Chomel P, et al. [Clinical application of grafts of cultured epidermis in burn patients. Apropos of 16 patients]. <i>Ann Chir Plast Esthet.</i> 1990;35(1):39-46.	7
595	Abbes M, Bourgeon Y, Regnier M, Gioanni J, Ettore F. [Our experience with the use of epidermal cell cultures in reconstructive surgery]. <i>Ann Chir Plast Esthet.</i> 1990;35(1):31-8.	7
596	Petersen MJ, Lessane B, Woodley DT. Characterization of cellular elements in healed cultured keratinocyte autografts used to cover burn wounds. <i>Arch Dermatol.</i> 1990;126(2):175-80.	3
597	Hilligweg UR, Close D. Absence makes the skin grow fonder. <i>Can Nurse.</i> 1989;85(10):20-1.	6
598	Woodley DT. Covering wounds with cultured keratinocytes. <i>Jama.</i> 1989;262(15):2140-1.	5
599	Hansbrough JF, Boyce ST, Cooper ML, Foreman TJ. Burn wound closure with cultured autologous keratinocytes and fibroblasts attached to a collagen-glycosaminoglycan substrate. <i>Jama.</i> 1989;262(15):2125-30.	6
600	Frenk E, Benathan M. [Epidermal cell cultures: does it have clinical applications?]. <i>Rev Med Suisse Romande.</i> 1989;109(6):423-5.	7
601	Khalfan HA, Hamza AA, Saeed T, Mohammed AW, Abalkhail A. Novel method of skin substitution in plastic surgery. <i>Burns.</i> 1989;15(5):335-7.	6

연번	서지정보	배제 사유
602	Hufnagel B, Ninnemann JL, Hettich R. Immunology of intermingled skin grafts in rats: preliminary results. <i>Burns Incl Therm Inj.</i> 1989;15(1):31-5.	4
603	Hancock K, Leigh IM. Cultured keratinocytes and keratinocyte grafts. <i>Bmj.</i> 1989;299(6709):1179-80.	5
604	Morioka N, Otsuka F, Kobayashi K, Tsuchida T, Matsuyama T, Seki Y, et al. [A trial of a burn wound treatment employing autografting of cultured keratinocytes]. <i>Nippon Hifuka Gakkai Zasshi.</i> 1989;99(1):73-6.	7
605	Compton CC, Gill JM, Bradford DA, Regauer S, Gallico GG, O'Connor NE. Skin regenerated from cultured epithelial autografts on full-thickness burn wounds from 6 days to 5 years after grafting. A light, electron microscopic and immunohistochemical study. <i>Lab Invest.</i> 1989;60(5):600-12.	3
606	Aihara M. Ultrastructural study of grafted autologous cultured human epithelium. <i>Br J Plast Surg.</i> 1989;42(1):35-42.	3
607	Herzog SR, Meyer A, Woodley D, Peterson HD. Wound coverage with cultured autologous keratinocytes: use after burn wound excision, including biopsy followup. <i>J Trauma.</i> 1988;28(2):195-8.	6
608	Phillips TJ. Cultured skin grafts. Past, present, future. <i>Arch Dermatol.</i> 1988;124(7):1035-8.	6
609	Woodley DT, Peterson HD, Herzog SR, Stricklin GP, Burgeson RE, Briggaman RA, et al. Burn wounds resurfaced by cultured epidermal autografts show abnormal reconstitution of anchoring fibrils. <i>Jama.</i> 1988;259(17):2566-71.	3
610	Aubock J, Irschick E, Romani N, Kompatscher P, Hopfl R, Herold M, et al. Rejection, after a slightly prolonged survival time, of Langerhans cell-free allogeneic cultured epidermis used for wound coverage in humans. <i>Transplantation.</i> 1988;45(4):730-7.	4
611	Soehnchen R, Braun-Falco O. [Epithelium transplantation with cultivated keratinocytes]. <i>Hautarzt.</i> 1988;39(11):701-7.	7
612	Fang CH, Yu GS, Fan YF, Wang K, Alexander JW. A preliminary report on transplantation of microskin autografts overlaid with sheet allograft in the treatment of large burns. <i>J Burn Care Rehabil.</i> 1988;9(6):629-33.	2
613	Lee JJ, Marvin JA, Heimbach DM, Grube BJ. Use of 5% sulfamylon (mafenide) solution after excision and grafting of burns. <i>J Burn Care Rehabil.</i> 1988;9(6):602-5.	2
614	Berger A, Burke JF. [Skin transplantation or artificial skin replacement]. <i>Zentralbl Chir.</i> 1988;113(12):751-7.	7
615	Aubock J. [Skin replacement with cultured keratinocytes]. <i>Z Hautkr.</i> 1988;63(7):565-7.	7
616	Fang RH, Yeh FL, Yinn JH, Shen HD, Chiang H, Lin YM, et al. [The clinical application of autologous human cultured epithelia]. <i>Chung Hua I Hsueh Tsa Chih (Taipei).</i> 1988;41(4):299-304.	7
617	Bettex-Galland M, Slongo T, Hunziker T, Wiesmann U, Bettex M. Use of cultured keratinocytes in the treatment of severe burns. <i>Z Kinderchir.</i> 1988;43(3):224-8.	6
618	Jedrzejczak WW. [Transplantation of epidermis grown in vitro]. <i>Pol Tyg Lek.</i> 1988;43(5):157-60.	7
619	Bagot M, Bertaux B, Heslan M, Coulomb B, Dubertret L. Reconstructed human epidermis: absence of Langerhans cells and failure to stimulate allogeneic lymphocytes in vitro. <i>Clin Exp Immunol.</i> 1988;71(1):138-43.	4
620	Pye RJ. Cultured keratinocytes as biological wound dressings. <i>Eye.</i> 1988;2(Pt 2):172-8.	6
621	Langdon RC, Cuono CB, Birchall N, Madri JA, Kuklinska E, McGuire J, et al. Reconstitution of structure and cell function in human skin grafts derived from cryopreserved allogeneic dermis and autologous cultured keratinocytes. <i>J Invest Dermatol.</i> 1988;91(5):478-85.	4
622	Deglise B, Benathan M, Frenk E, Krupp S. [Preliminary results of burn treatment using an autograft of cultured epidermis]. <i>Schweiz Med Wochenschr.</i> 1987;117(37):1380-3.	7

연번	서지정보	배제 사유
623	Teepe RG, Kreis RW, Hoekstra MJ, Ponec M. [Permanent wound cover using cultured autologous epidermis transplants: a new treatment method for patients with severe burns]. <i>Ned Tijdschr Geneeskd.</i> 1987;131(22):946-51.	7
624	Latarjet J, Gangolphe M, Hezez G, Masson C, Chomel PY, Cognet JB, et al. The grafting of burns with cultured epidermis as autografts in man. Two case reports. <i>Scand J Plast Reconstr Surg Hand Surg.</i> 1987;21(3):241-4.	6
625	Thompson CH, Streamer KJ, Baker P, Rose BR, Kennedy PJ, Cossart YE. Transplantation of cultured autologous epidermis to a patient with burns. <i>Med J Aust.</i> 1987;147(10):507-10.	6
626	Cuono CB, Langdon R, Birchall N, Barttelbort S, McGuire J. Composite autologous-allogeneic skin replacement: development and clinical application. <i>Plast Reconstr Surg.</i> 1987;80(4):626-37.	2
627	Eldad A, Burt A, Clarke JA, Gusterson B. Cultured epithelium as a skin substitute. <i>Burns Incl Therm Inj.</i> 1987;13(3):173-80.	6
628	Kuri-Harcuch W, Bolivar-Flores YJ, Marsch-Moreno M, Arkuch S. [Skin autografts obtained for in vitro culture of epidermal cells]. <i>Rev Invest Clin.</i> 1987;39(1):5-9.	7
629	Shiple GD, Pittelkow MR. Control of growth and differentiation in vitro of human keratinocytes cultured in serum-free medium. <i>Arch Dermatol.</i> 1987;123(11):1541a-4a.	4
630	Gangolphe M, Hezez G, Latarjet J, Masson CL, Foyatier JL, Faure M, et al. [Culture of the human epidermis. Use as an autograft in burn patients]. <i>Ann Chir Plast Esthet.</i> 1987;32(1):38-41.	7
631	Thivolet J, Faure M, Demidem A, Mauduit G. Long-term survival and immunological tolerance of human epidermal allografts produced in culture. <i>Transplantation.</i> 1986;42(3):274-80.	6
632	Madden MR, Finkelstein JL, Staiano-Coico L, Goodwin CW, Shires GT, Nolan EE, et al. Grafting of cultured allogeneic epidermis on second- and third-degree burn wounds on 26 patients. <i>J Trauma.</i> 1986;26(11):955-62.	2
633	Twomey JA. Cultured autologous skin for burns. <i>Mayo Clin Proc.</i> 1986;61(10):832-4.	6
634	Carney SA. Generation of autograft; the state of the art. <i>Burns Incl Therm Inj.</i> 1986;12(4):231-5.	6
635	Brown AS, Barot LR. Biologic dressings and skin substitutes. <i>Clin Plast Surg.</i> 1986;13(1):69-74.	5
636	Clarke JA, Burt AM, Eldad A, Gusterson BA. Cultured skin for burn injury. <i>Lancet.</i> 1986;2(8510):809.	5
637	Thomson AW, Sewell HF, Abramovich DR. Skin replacement after burn injury. <i>Lancet.</i> 1986;2(8499):163.	5
638	Teepe RG, Ponec M, Kreis RW, Hermans RP. Improved grafting method for treatment of burns with autologous cultured human epithelium. <i>Lancet.</i> 1986;1(8477):385.	5
639	Bonnekoh B, Thiele B, Mahrle G, Steigleder GK. [Epidermal cell cultures--significance for wound coverage in the human]. <i>Z Hautkr.</i> 1986;61(20):1433-42.	7
640	Thivolet J, Faure M, Gaucherand M, Reano A. [Rapid production of large surface epidermal autografts]. <i>Ann Chir Plast Esthet.</i> 1986;31(2):137-42.	7
641	Pittelkow MR, Scott RE. New techniques for the in vitro culture of human skin keratinocytes and perspectives on their use for grafting of patients with extensive burns. <i>Mayo Clin Proc.</i> 1986;61(10):771-7.	4
642	Cuono C, Langdon R, McGuire J. Use of cultured epidermal autografts and dermal allografts as skin replacement after burn injury. <i>Lancet.</i> 1986;1(8490):1123-4.	6
643	Gallico GG, 3rd, O'Connor NE. Cultured epithelium as a skin substitute. <i>Clin Plast Surg.</i> 1985;12(2):149-57.	5
644	Fisher JC. Skin--the ultimate solution for the burn wound. <i>N Engl J Med.</i> 1984;311(7):466-7.	5

연번	서지정보	배제 사유
645	Gallico GG, 3rd, O'Connor NE, Compton CC, Kehinde O, Green H. Permanent coverage of large burn wounds with autologous cultured human epithelium. N Engl J Med. 1984;311(7):448-51.	6
646	Hefton JM, Madden MR, Finkelstein JL, Shires GT. Grafting of burn patients with allografts of cultured epidermal cells. Lancet. 1983;2(8347):428-30.	2
647	Yannas IV, Burke JF, Orgill DP, Skrabut EM. Wound tissue can utilize a polymeric template to synthesize a functional extension of skin. Science. 1982;215(4529):174-6.	6
648	Cason JS. Some aspects on prevention and treatment of infection in burns. Prog Pediatr Surg. 1981;14:3-18.	6
649	Yarborough MF. Care of the burn wound. Cutis. 1978;22(4):447-52.	2
650	Szabo SE, Toomey JM, Linn BS. Does skin have antimicrobial properties? An in-vitro experiment and literature review. Am Surg. 1978;44(1):55-8.	4
651	Klein P, Wienkotter R. [Cultivation of the epithelium in burns]. Chirurg. 1976;47(7):400-5.	7
652	Igel HJ, Freeman AE, Boeckman CR, Kleinfeld KL. A new method for covering large surface area wounds with autografts. II. Surgical application of tissue culture expanded rabbit-skin autografts. Arch Surg. 1974;108(5):724-9.	6
653	Freeman AE, Igel HJ, Waldman NL, Losikoff AM. A new method for covering large surface area wounds with autografts. I. In vitro multiplication of rabbit-skin epithelial cells. Arch Surg. 1974;108(5):721-3.	6
654	Summerlin WT, Broutbar C, Foanes RB, Payne R, Stutman O, Hayflick L, et al. Acceptance of phenotypically differing cultured skin in man and mice. Transplant Proc. 1973;5(1):707-10.	6
655	Rubin LR, Bongiovi J, Jr. Human skin--antibacterial in vitro? J Surg Res. 1971;11(7):321-4.	6
656	Larsson AP, Briheim K, Hanna V, Gustafsson K, Starkenberg A, Vintertun HN, et al. Transplantation of autologous cells and porous gelatin microcarriers to promote wound healing. Burns. 2021;47(3):601-10.	4
657	Borzenok SA, Malyugin BE, Gerasimov MY, Ostrovsky DS. Cultivated autologous oral mucosal epithelial transplantation. [Russian]. Vestnik Transplantologii i Iskusstvennykh Organov. 2021;23(1):171-7.	7
658	Benchaprathanphorn K, Sakulaue P, Siriwatwechakul W, Muangman P, Chinaroonchai K, Viravaidya-Pasuwat K. Preparation and characterization of human keratinocyte-fibroblast cell sheets constructed using PNIAm-co-AM grafted surfaces for burn wound healing. Journal of Materials Science: Materials in Medicine. 2020;31 (12) (no pagination)(126).	6
659	Cortez Ghio S, Larouche D, Doucet EJ, Germain L. The role of cultured autologous bilayered skin substitutes as epithelial stem cell niches after grafting: A systematic review of clinical studies. Burns Open. 2021;5(2):56-66.	6
660	Supp DM, Hahn JM, Lloyd CM, Combs KA, Swope VB, Abdel-Malek Z, et al. Light or dark pigmentation of engineered skin substitutes containing melanocytes protects against ultraviolet light-induced DNA damage in vivo. Journal of Burn Care and Research. 2020;41(4):751-60.	2
661	Li Z, Maitz P. Cell therapy for severe burn wound healing. Burns and Trauma. 2018;6 (1) (no pagination)(13).	6
662	Shahin H, Elmasry M, Steinvall I, Soberg F, El-Serafi A. Vascularization is the next challenge for skin tissue engineering as a solution for burn management. Burns and Trauma. 2020;8 (no pagination)(TKAA022).	6
663	Guglielmetti A, Jahr C, Gompertz M. First experience of autologous fibroblasts for the treatment of cutaneous loxoscelism. Journal of the Dermatology Nurses' Association Conference: 24th World Congress of Dermatology Milan Italy. 2020;12(2).	5

연번	서지정보	배제 사유
664	Mitra D. A randomized control trial to study the efficacy of cultured autologous keratinocyte-melanocyte transplant versus conservative wound dressing in the management of post burns wound healing. Journal of the Dermatology Nurses' Association Conference: 24th World Congress of Dermatology Milan Italy. 2020;12(2).	5
665	Zahorec P, Sarkozyova N, Ferancikova N, Bukovcan P, Danisovic L, Bohac M, et al. Autologous mesenchymal stem cells application in post-burn scars treatment: a preliminary study. Cell and tissue banking. 2021;22(1):39-46.	2
666	Navsaria HA, Rugg EL. Telomere shortening: Significant for keratinocyte grafting? Lancet. 2003;361(9366):1316-7.	5
667	De Rosa L. Hologene projects. Acta Dermato-Venereologica. 2020;100 (SUPPL 220):12.	5
668	Moravvej H, Hormozi AK, Hosseini SN, Sorouri R, Mozafari N, Ghazisaidi MR, et al. Comparison of the application of allogeneic fibroblast and autologous mesh grafting with the conventional method in the treatment of third-degree burns. Journal of Burn Care and Research. 2012;07.	2
669	Wood FM, Giles N, Stevenson A, Rea S, Fear M. Characterisation of the cell suspension harvested from the dermal epidermal junction using a ReCell<sup></sup> kit. Burns.	2
670	Christensen JM, Shanbhag SS, Shih GC, Goverman J, Pomahac B, Chodosh J, et al. Multidisciplinary treatment to restore vision in ocular burns. Journal of Burn Care and Research. 2020;41(4):859-65.	1
671	Nguyen T, Ewe KYH, Wood F, Rea S, Bowen AC. Case report: Scald burn to the scalp complicated by fungal kerion. Burns Open. 2020;4(4):191-3.	6
672	Rechichi M, Ferrise M, Romano F, Gallelli L, Toschi V, Dominijanni A, et al. Autologous platelet-rich plasma in the treatment of refractory corneal ulcers: A case report. American Journal of Ophthalmology Case Reports. 2020;20 (no pagination)(100838).	1
673	Greenwood JE, Damkat-Thomas L, Schmitt B, Dearman B. Successful proof of the 'two-stage strategy' for major burn wound repair. Burns Open. 2020;4(3):121-31.	6
674	Abdo J, Ortman H. Biologic and Synthetic Cellular and/or Tissue-Based Products and Smart Wound Dressings/Coverings. Surgical Clinics of North America. 2020;100(4):741-56.	6
675	Abdo JM, Sopko NA, Milner SM. The applied anatomy of human skin: A model for regeneration. Wound Medicine. 2020;28 (no pagination)(100179).	6
676	Bajouri A, Orouji Z, Taghiabadi E, Nazari A, Shahbazi A, Fallah N, et al. Long-term follow-up of autologous fibroblast transplantation for facial contour deformities, a non-randomized phase IIa clinical trial. Cell j. 2020;22(1):75-84.	1
677	Arias-Santiago S, Fernandez-Gonzalez A, Pharm GGC, Gacto P, Lizana-Moreno AM, Ruiz-Garcia A, et al. 16846 Clinical evaluation, cutaneous homeostasis, and epidermal barrier analysis of a human tissue engineered skin substitute for severe burn patients. Journal of the American Academy of Dermatology. 2020;83 (6 Supplement):AB73.	5
678	Altinors DD, Asena L, Oto S. Limbal stem cell transplantation for restoration of the ocular surface health in bilateral limbal stem cell deficiency. Transplantation. 2020;104 (SUPPL 3):S212-S3.	5
679	Ghahary A, Baradar R, Jalili, Kilani RT. Challenges and solutions for treating complicated wounds: Result of a feasibility human pilot study. Wound Repair and Regeneration. 2020;28 (2):A20.	5
680	Trouillas M, Dedier M, Magne B, Nivet M, Peltzer J, Banzet S. Interest of interleukin-1beta priming on various sources of MSC for wound healing: role of interleukin-6, a MSC secreted factor. Cytotherapy. 2020;22 (5 Supplement):S109-S10.	5
681	Flamant S, Busson E, Ribault A, L'Homme B B, Massourides E, Pinset C, et al. Human induced pluripotent stem cells as a source of mesenchymal stromal cells for the treatment of cutaneous radiation injuries. Cytotherapy. 2020;22 (5 Supplement):S44.	5

연번	서지정보	배제 사유
682	Joethy J, Koh K, Wong AWJ, Cheng A, Sellam S, Chong SJ. Initial impression of the CelluTome™ Epidermal Harvesting System in burns management. <i>Burns Open</i> . 2019;3(2):68-74.	2
683	Samberg M, Stone R, Natesan S, Kowalczewski A, Becerra S, Wrice N, et al. Platelet rich plasma hydrogels promote in vitro and in vivo angiogenic potential of adipose-derived stem cells. <i>Acta Biomater</i> . 2019;87:76-87.	6
684	Kaur A, Midha S, Giri S, Mohanty S. Functional skin grafts: Where biomaterials meet stem cells. <i>Stem Cells Int</i> . 2019;2019 (no pagination)(1286054).	6
685	Abdel-Sayed P, Michetti M, Scaletta C, Flahaut M, Hirt-Burri N, De Buys Roessingh A, et al. Cell therapies for skin regeneration: An overview of 40 years of experience in burn units. <i>Swiss Medical Weekly</i> . 2019;149 (19-20) (no pagination)(w20079).	6
686	Ardailou R. Historical progresses in the knowledge and therapeutical use of stem cells in humans. <i>Archives of Hellenic Medicine</i> . 2019;36 (Supplement 2):11-2.	5
687	Merayo-Lloves J, Sanchez-Avila R, Chacon M, Vazquez N, Berisa S, Medina MP, et al. Plasma Rich in Growth Factors membrane in the treatment of Limbal Stem Cell Deficiency. <i>Investigative Ophthalmology and Visual Science Conference</i> . 2019;60(9).	5
688	Niimi Y, Suzan A, Fukuda S, Osada A, Hawkins H, Cox R, et al. Use of artificial dermis as a wound bed for cultured keratinocyte sheet in an ovine burn wound model. <i>Critical Care Medicine Conference: 48th Critical Care Congress of the Society of Critical Care Medicine, SCCM</i> . 2019;47(1 Supplement 1).	5
689	Glat PM, Nanassy AD, Burkey BA, Misra AC, Meyer LK, Gates L, et al. The use of an autologous cell harvesting and processing device in two burn patients at an urban pediatric burn center. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S146.	5
690	Niimi Y, Alharbi S, Fukuda S, Popov VL, Liu K, Osada A, et al. Use of artificial dermis as a wound bed for cultured keratinocyte sheet cultured using temperature response dishes in an ovine burn wound healing model. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S231-S2.	5
691	Foster KN, Richey KJ, Osborn SC, Burkey B, Carter J, Davis W, et al. Healing of donor sites with an autologous skin cell suspension for large tbsa burn injuries: A prospective evaluation. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S225.	5
692	Sangji NF, Levin JM, Friedstat JS, Goverman J, Chang KM, Schulz JT. Acute and reconstructive burn surgery with a bilaminar polyurethane skin substitute: A case series. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S211.	5
693	Fear M, Wood F. 10 years of clinical experience using point of care non cultured autologous skin cell suspension. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S71.	5
694	Molnar JA, Holmes J, Carter J, Lee J, Foster K, Kahn S, et al. Evaluation of autologous skin cell suspension for healing of burn injuries of the hand. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S70.	5
695	Malara MM, Blackstone BN, Baumann ME, DeBruler DM, Bailey JK, Supp DM, et al. Combinatorial use of CEAs with dermal substitutes containing dermal papilla-like structures. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S151.	5
696	Craig CK, Williams JW, Barone CG, Aitken MM, Carter JE, Molnar JA, et al. Post-operative wound management following the use of an autologous cell harvesting device in the treatment of patients with life-threatening injuries: A single center's experience. <i>Journal of Burn Care and Research</i> . 2019;40 (Supplement 1):S69.	5

연번	서지정보	배제 사유
697	Niimi Y, Suzan A, Fukuda S, Wiener DJ, Liu K, Popov VL, et al. Effect of artificial dermis as a wound bed for keratinocyte sheet grafting cultured by temperature responsive dish in clinically relevant ovine burn wound model. <i>Shock</i> . 2019;51 (6 Supplement 1):69.	5
698	Yeni B. Transgenic skin quality of an epidermolysis bullosa patient after autologous transplantation of transgenic epidermal stem cells. <i>European Surgical Research</i> . 2019;60 (Supplement 1):25-6.	5
699	Trouillas M, Magne B, Dedier M, Nivet M, Lataillade J, Banzet S. HGF, MMPs AND TGF-beta1 CONTRIBUTE TO INTERLEUKIN-1beta PRIMED MESENCHYMAL STROMAL CELLS EFFECT ON WOUND HEALING AND EPIDERMAL SUBSTITUTE ENGRAFTMENT. <i>Cytotherapy</i> . 2019;21 (5 Supplement):S51.	5
700	Mangum L, Stone R, Natesan S, Christy B, Clifford J, Kumar R, et al. Adipose stem cell tissue source is associated with differential expression profiles in cellular mRNA and exosomal mirna. <i>Wound Repair and Regeneration</i> . 2019;27 (3):A9-A10.	5
701	Alharbi S, Niimi Y, Wiener D, Hawkins H, Cox R, Popov V, et al. Non-enzymatic detachment for keratinocyte sheets using temperature responsive dishes for treating burn wound. <i>Wound Repair and Regeneration</i> . 2019;27 (3):A28.	5
702	Perin B, Addetia A, Qin X. Autologous skin microbiota transplantation: Development of a technique to transfer representative and viable cutaneous microbial communities. <i>Journal of the American Academy of Dermatology</i> . 2018;79 (3 Supplement 1):AB55.	5
703	Burmeister DM, Roy DC, Becerra SC, Natesan S, Christy RJ. In situ delivery of fibrin-based hydrogels prevents contraction and reduces inflammation. <i>Journal of Burn Care and Research</i> . 2018;39(1):40-53.	4
704	Liu J, Qu W, Li R, Zheng C, Zhang L. Efficacy of autologous platelet-rich gel in the treatment of deep grade II burn wounds. <i>International Journal of Clinical and Experimental Medicine</i> . 2018;11(3):2654-9.	2
705	Figueiredo GS, Salvador-Culla B, Baylis OJ, Mudhar HS, Lako M, Figueiredo FC. Outcomes of Penetrating Keratoplasty Following Autologous Cultivated Limbal Epithelial Stem Cell Transplantation. <i>Stem Cells</i> . 2018;36(6):925-31.	1
706	Aries P, Hoffmann C, Schaal JV, Leclerc T, Donat N, Cirodde A, et al. <i>Aspergillus tamarii</i> : An uncommon burn wound infection. <i>Journal of Clinical Pathology</i> . 2018;71(4):379-80.	5
707	Brockmann I, Ehrenpfordt J, Sturmheit T, Brandenburger M, Kruse C, Zille M, et al. Skin-derived stem cells for wound treatment using cultured epidermal autografts: Clinical applications and challenges. <i>Stem Cells Int</i> . 2018;2018 (no pagination)(4623615).	6
708	Germain L, Larouche D, Nedelec B, Perreault I, Duranceau L, Bortoluzzi P, et al. Autologous bilayered self-assembled skin substitutes (Sasss) as permanent grafts: A case series of 14 severely burned patients indicating clinical effectiveness. <i>European Cells and Materials</i> . 2018;36:128-41.	2
709	Seol YJ, Lee H, Copus JS, Kang HW, Cho DW, Atala A, et al. 3D bioprinted biomask for facial skin reconstruction. <i>Bioprinting</i> . 2018;10 (no pagination)(e00028).	4
710	Tashiro A, Nei T, Sugimoto R, Watanabe A, Hagiwara J, Takiguchi T, et al. <i>Kodamaea ohmeri</i> fungemia in severe burn: Case study and literature review. <i>Medical Mycology Case Reports</i> . 2018;22:21-3.	6
711	Ornelas-Flores MC, Garcia-Lopez J, Melgarejo-Ramirez Y, Sanchez-Sanchez R, Leyva-Gomez G, Zacula-Juarez N, et al. Implantation of a heterologous dermo-epidermal skin substitute in a patient with deep dermal burn that enhances biomechanical and functional recovery: Case report. <i>Burns Open</i> . 2018;2(3):144-53.	2

연번	서지정보	배제 사유
712	Gibbs S, Van Den Broek LJ, Thon M, Atac B, Monsuur H, Lindner G, et al. Introduction of hair papillae into tissueengineered skin. <i>Wound Repair and Regeneration</i> . 2018;26 (2):A22.	5
713	Akita S. Novel application of cultured epithelial autografts (CEA) with expanded mesh skin grafting over artificial dermis. <i>Wound Repair and Regeneration</i> . 2018;26 (4):A10.	5
714	Stysly B, Miles MJ, Dillard TA, Jamindar P. Findings that will make you squirm: A case of hypoxic respiratory failure secondary to pulmonary strongyloides. <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2018;197(MeetingAbstracts).	5
715	Alharbi S, Niimi Y, Hawkins H, Cox R, Popov V, Osada A, et al. Effects of autologous keratinocyte sheets on ovine burn wound healing. <i>Shock</i> . 2018;49 (6 Supplement 1):140.	5
716	Alharbi S, Niimi Y, Hawkins H, Cox R, Osada A, Ihara K, et al. Efficacy of keratinocyte sheet cultured in temperature responsive dish in ovine burn wound healing. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S155.	5
717	Niimi Y, Perez-Bello D, Fukuda S, Baljinyam T, Andersen CR, Kim J, et al. Omega-7 accelerates burn wound healing via increase telomerase activity. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S154-S5.	5
718	Allen A, Recht K, Litt JS. Successful posterior cultured epidermal autograft placement to a major burn victim: A university burn center experience and review of the literature. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S231.	5
719	O'Neil AM, Rush C, Griffard L, Roggy D, Sood R. Retrospective analysis of the utilization of a multidisciplinary algorithm for mobilization of the vented burn patient. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S46.	5
720	Walker NJ, Rebowe RE, Allred L, Craig CK, Williams J, Holmes JH, et al. Initial experience with autologous cell suspension for treatment of partial thickness facial burns. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S44.	5
721	Kueckelhaus M, Rothoef T, Teig N, Jacobsen F, Lehnhardt M, Pellegrini G, et al. Regeneration of the entire human epidermis by transgenic epidermal stem cell transplants and its implications for the treatment of burns. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S148.	5
722	Hernandez G, Smith C, Williamson S, Branski L. Autograft storage practices of ABA verified burn centers. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S120.	5
723	Magnusson S, McLntyre M, Burmeister DM, Christy RJ, Kjartansson H, Baldursson BT, et al. Fish skin grafts support ingrowth and colonization of adipose derived stem cells in vitro. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S63.	5
724	Holmes JH, Carter JE, Molnar JA, Cairns BA, Hwang J, King BT, et al. Demonstration of the safety and effectiveness of autologous skin cell suspension combined with meshed skin grafts for the reduction of donor area in the treatment of acute burns. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S60.	5
725	Foster K, Bilir P, Kruger E, Kowal S, Holmes JH, Hickerson W, et al. Cost-effectiveness (CE) of an autologous regenerative epithelial suspension (RES) versus standard of care (SOC) for treatment of severe burns in the United States. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S113.	5
726	Chong S, Chua A, Kok Y. Only micrograft/CEA vs Cuono's method: A review. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S244.	5
727	Hickerson WL, Molnar JA, Carter JE, Cairns BA, Hwang J, King BT, et al. A comparative study of autologous skin cell suspension to split-thickness autografting in the treatment of acute burns. <i>Journal of Burn Care and Research</i> . 2018;39 (Supplement 1):S1.	5



연번	서지정보	배제 사유
728	Haddad AG, Giatsidis G, Orgill DP, Halvorson EG. Skin Substitutes and Bioscaffolds: Temporary and Permanent Coverage. <i>Clin Plast Surg.</i> 2017;44(3):627-34.	6
729	Leonard DA, Mallard C, Albritton A, Torabi R, Mastroianni M, Sachs DH, et al. Skin grafts from genetically modified alpha-1,3-galactosyltransferase knockout miniature swine: A functional equivalent to allografts. <i>Burns.</i> 2017;43(8):1717-24.	4
730	Berthet M, Gauthier Y, Lacroix C, Verrier B, Monge C. Nanoparticle-Based Dressing: The Future of Wound Treatment? <i>Trends Biotechnol.</i> 2017;35(8):770-84.	6
731	Strid Orrhult L, Sunden J, Gatenholm P. 3D bioprinting of a human skin tissue model. <i>Tissue Engineering - Part A.</i> 2017;23 (Supplement 1):S76.	5
732	Alharbi S, Niimi Y, Hawkins H, Cox R, Osada A, Ihara K, et al. Efficacy of keratinocyte sheet cultured in temperature responsive dish in ovine burn wound healing. <i>Tissue Engineering - Part A.</i> 2017;23 (Supplement 1):S74.	5
733	Manoharan S. ReCell autologous cell harvesting device in the treatment of challenging indications and skin types. <i>Australas J Dermatol.</i> 2017;58 (4):334.	5
734	Waibel J, Badiavas E, Davis S, Rodriguez-Menocal L. Hypertrophic burn scars modulation through laser assisted delivery of adipose derived stem cells vs bone-marrow mesenchymal stem cells with the CO <sub>2</sub> vs ER: YAG laser. <i>Lasers in Surgery and Medicine.</i> 2017;49 (Supplement 28):4.	5
735	Ritore A, Garzon I, Arias S, Carriel V, Martin-Piedra MA, Cuende N, et al. A top-notch approach: Fibrin-agarose based artificial tissues. <i>Cytotherapy.</i> 2017;19 (5 Supplement 1):S235-S6.	5
736	Guerrero-Calvo J, Fernandez-Gonzalez A, Lizana-Moreno A, Montes-Ramos P, Fernandez N, Ruiz-Garcia A, et al. Optimization of human keratinocyte culture to develop a tissue bio-engineered human skin model. <i>Cytotherapy.</i> 2017;19 (5 Supplement 1):S233-S4.	5
737	Fernandez-Gonzalez A, Lizana-Moreno A, Fernandez N, Guerrero-Calvo J, Ruiz-Garcia A, Espinosa O, et al. Clinical, histological and homeostasis evaluation of an autologous tissue bio-engineered skin substitute in a patient with 70% of total body surface area (TBSA) burn. <i>Cytotherapy.</i> 2017;19 (5 Supplement 1):S233.	5
738	Fernandez-Gonzalez A, Lizana-Moreno A, Ruiz-Garcia A, Guerrero-Calvo J, Espinosa O, Fernandez N, et al. Optimization of an autologous human skin model under gmp guidelines. <i>Cytotherapy.</i> 2017;19 (5 Supplement 1):S232-S3.	5
739	Turner M, Bienek C, McGowan N, Dhillon B, Campbell J, Glover A, et al. A phase I/II clinical trial of allogeneic ex vivo expanded corneal epithelial stem cells in patients with severe ocular surface disorder arising from limbal stem cell deficiency. <i>Cytotherapy.</i> 2017;19 (5 Supplement 1):S231.	5
740	Akbarzadeh S, Paul M, Herson M, Lo C, Cleland H. CEA as an adjunct treatment for major burns: A phase I study. <i>Cytotherapy.</i> 2017;19 (5 Supplement 1):S26-S7.	5
741	Murakami D, Harada H, Suehiro S, Ohki T, Yamato M. Prevention of stenosis with endoscopic transplantation of cultured autologous oral mucosal epithelial cell sheets after esophageal endoscopic submucosal dissection. <i>United European Gastroenterology Journal.</i> 2017;5 (5 Supplement 1):A339-A40.	5
742	He K, Zhao L, Bu S, Liu L, Wang M, Wang X, et al. Endoscopic autologous transplantation of esophageal mucosa for treating the refractory caustic esophageal stricture. <i>United European Gastroenterology Journal.</i> 2017;5 (5 Supplement 1):A579-A80.	5
743	Jean-Jacques L, Marina T, Muriel N, Elodie B, Patrick D, Brachet M, et al. Cell therapy of burns: Present and future. <i>Wound Repair and Regeneration.</i> 2017;25 (4):A42.	5
744	Rodriguez P, Reyna L. Burn treatment with epifast (live cultured human keratinocytes): Our experience at least five years- protocol treatment with a biological dressing. <i>Wound Repair and Regeneration.</i> 2017;25 (4):A25.	5

연번	서지정보	배제 사유
745	Yu T, He K, Wang K, Liu L, Fan Z. Endoscopic autologous transplantation of esophageal mucosa for treating the refractory caustic esophageal stricture. <i>Gastrointestinal Endoscopy</i> . 2017;85 (5 Supplement 1):AB523.	5
746	He K, Zhao L, Liu L, Wang M, Wang X, Fan Z. Endoscopic autologous transplantation of esophageal mucosa for treating the refractory caustic esophageal stricture. <i>Journal of Gastroenterology and Hepatology</i> . 2017;32 (Supplement 3):273.	5
747	Nanba D, Toki F, Matsumura H, Toki H, Nishimura EK. Locomotive ability of human keratinocyte stem cells is an intrinsic property for stem cell expansion and epidermal reconstruction. <i>J Invest Dermatol</i> . 2017;137 (10 Supplement 2):S310.	5
748	Mangum LH, Wrice N, Larson D, Natesan S, Christy R. Comparison of growth supplements on the differentiation of adipose derived stem cells from burned and non-burned human skin. <i>FASEB Journal Conference: Experimental Biology</i> . 2017;31(1 Supplement 1).	5
749	Vig K, Chaudhari A, Tripathi S, Dixit S, Sahu R, Pillai S, et al. Advances in skin regeneration using tissue engineering. <i>Int</i> . 2017;18 (4) (no pagination)(789).	6
750	Gupta GJ, Karki K, Jain P, Saxena AK. Autologous bone marrow aspirate therapy for skin tissue engineering and tissue regeneration. <i>Adv Wound Care (New Rochelle)</i> . 2017;6(4):135-42.	1
751	Nan W, Xu Z, Chen H. Advances in epidermis engineering: Application of epidermal stem cells (ESCs) and tissue engineered skin (TES). <i>Biomedical Research (India)</i> . 2017;28(5):2290-5.	5
752	Gardenier JC, Chopra VK, Filicori F, Murphy J, Greenway A, Gallagher JJ, et al. Angioinvasive mold in the surgical and burn intensive care unit: A case series and review of the literature. <i>Surgical Infections Case Reports</i> . 2016;1(1):72-8.	6
753	Lorente JA, Amaya-Villar R. Update in the management of critically ill burned patients. <i>Medicina Intensiva</i> . 2016;40(1):46-8.	5
754	Huh MI, An SH, Kim HG, Song YJ, Choi EC, An SH, et al. Rapid expansion and auto-grafting efficiency of porcine full skin expanded by a skin bioreactor ex vivo. <i>Tissue Engineering and Regenerative Medicine</i> . 2016;13(1):31-8.	4
755	Michele De L. Biological parameters determining the clinical outcome of autologous cultures of limbal stem cells. <i>Cell j</i> . 2016;18 (Supplement 1):14-5.	5
756	Nanba D, Toki F, Tate S, Imai M, Matsushita N, Toki H, et al. Cell motion predicts human epidermal stemness. <i>J Dermatol Sci</i> . 2016;84 (1):e51.	5
757	Velasquillo C, Sanchez-Sanchez R, Brena A, Martinez V, Melgarejo Y, Rodriguez L, et al. Biological skin substitutes based on amnion and pig skin seeded with mesenchymal stem cells for the treatment of autologous burns. <i>Tissue Engineering - Part A</i> . 2016;22 (Supplement 1):S15.	5
758	Kitala D, Kawecki M, Klama-Baryla A, Labus W, Glik J, Kraut M, et al. Amniotic stem cells share clusters of differentiation of fibroblasts and keratinocytes, influencing their wound healing ability. <i>Archivum Immunologiae et Therapiae Experimentalis</i> . 2016;64 (1 Supplement 1):S29.	5
759	Bobba S, Di Girolamo N, Watson SL. 9. Mid-term outcomes of therapy for limbal stem cell deficiency utilising an autologous xeno-free transplantation technique. <i>Clinical and Experimental Ophthalmology</i> . 2016;44 (Supplement 1):64-5.	5
760	Mullins RF, Cartie RJ, Hassan S, Homsombath B, Brandigi C, Fagan SP, et al. Retrospective review of adult and pediatric cultured epidermal autograft (CEA) patients. <i>Journal of Burn Care and Research</i> . 2016;1):S281.	5
761	Gibson A, Greenhalgh D. Variations in burn excision and grafting. <i>Journal of Burn Care and Research</i> . 2016;1):S271.	5

연번	서지정보	배제 사유
762	Agraz AN, Schaumberg MD, Hartman BC, Sood R. Case study evaluating the effectiveness of serial casting in the adult population for ankle plantarflexion contractures post burn injury. <i>Journal of Burn Care and Research</i> . 2016;1):S268.	5
763	Greenwood JE. Bioreactor development for composite cultured skin production. <i>Journal of Burn Care and Research</i> . 2016;1):S266.	5
764	Boyce ST, Lloyd CM, Kleiner MC, Supp DM. Cultured human melanocytes in engineered skin substitutes provide rapid restoration of natural skin color after transplantation to full-thickness wounds. <i>Journal of Burn Care and Research</i> . 2016;1):S86.	5
765	Sivan U, Kalliyana Krishnan V, Krishnan LK. Anti microbial and non-toxic concentration of silver nanoparticles support differentiation of adipose derived mesenchymal stem cells to fibroblast for skin tissue engineering. <i>Current Tissue Engineering</i> . 2016;5(2).	4
766	Saidi S. Live skin allograft in the management of severe burns. <i>Annals of African Surgery</i> . 2016;13(2):77-80.	2
767	Maver T, Maver U, Kleinschek KS, Rascan IM, Smrke DM. Advanced therapies of skin injuries. <i>Wiener Klinische Wochenschrift</i> . 2015;127(Supplement 5):187-98.	6
768	Steinke M, Dally I, Friedel G, Walles H, Walles T. Host-integration of a tissue-engineered airway patch: Two-year follow-up in a single patient. <i>Tissue Engineering - Part A</i> . 2015;21(3-4):573-9.	1
769	Kaasik K, Tiido T. First in Europe: Stem-cell based medicinal product for treatment of limbal stem cell deficiency caused by burns to the eye. [Estonian]. <i>Eesti Arst</i> . 2015;94(2):111-2.	7
770	Sehic A, Utheim OA, Ommundsen K, Utheim TP. Pre-clinical cell-based therapy for limbal stem cell deficiency. <i>Journal of Functional Biomaterials</i> . 2015;6(3):863-88.	6
771	Varkey M, Ding J, Tredget EE. Advances in skin substitutes-potential of tissue engineered skin for facilitating anti-fibrotic healing. <i>Journal of Functional Biomaterials</i> . 2015;6(3):547-63.	6
772	Wu L, Wang C, Yang W, Yang X, Zhang L, Wang J. Inhibition of bone marrow mesenchymal stem cell autologous transplantation on inflammation following rabbit corneal alkali burn. [Chinese]. <i>Zhonghua Shiyen Yanke Zazhi/Chinese Journal of Experimental Ophthalmology</i> . 2015;33(9):798-804.	7
773	Moe M. Clinical results and in vitro characterization of cornea limbal epithelial stem cells cultured in autologous serum. <i>Acta Ophthalmologica Conference</i> . 2015;93(Supplement 255).	5
774	Huber B, Gehrke S, Kluger P, Jacobsen F, Nottrodt N. Development of a three layered skin model consisting of a fatty tissue layer with dermis and epidermis. <i>Int J Artif Organs</i> . 2015;38 (7):378.	5
775	Savkovic V, Schneider M, Sulflow K, Loth T, Hacker M, Schulz-Siegmund M, et al. Why only vitiligo? Pigmented epidermal graft composed of human hair follicle-derived melanocytes and keratinocytes for purposes of treating depigmentation disorders and wound healing. <i>Regen</i> . 2015;1):257.	5
776	Kremling H, Schutze K, Marino D, Meyer S. Raman spectroscopy: A novel tool for label-free and non-invasive cell analysis and quality assurance in 3D-skin grafts. <i>Regen</i> . 2015;1):78.	5
777	Carney B, Prinzeze N, Jo D, Moffatt L, Shupp J. Optimizing the preparation and subsequent culturing of cells obtained from an autologous cell harvesting device. <i>Journal of Burn Care and Research</i> . 2015;1):S274.	5
778	Rodriguez-Ferreya Sr P, Lule I. Burn treatment with live cultured human keratinocytes "biological dressing: Our experience at least five (5) years". <i>Journal of Burn Care and Research</i> . 2015;1):S268.	5

연번	서지정보	배제 사유
779	Shang Y, Shen C, Chai J, Ma L, Li D, Li L, et al. An experimental study on the repair of full skin loss wound in rats with graft of autologous keratinocyte suspension and microskin. <i>Journal of Burn Care and Research</i> . 2015;1):S201.	5
780	Beaudoin Cloutier C, Goyer B, Perron C, Gauvin R, Guignard R, Lacroix D, et al. Bilayered self-assembled skin substitute using a tissue-engineered dermal matrix grafted on athymic mice. <i>Journal of Burn Care and Research</i> . 2015;1):S200.	5
781	Benjamin NC, Williams F, Norbury W, Andersen CR, Suman O, Herndon DN. Effects of cultured substitutes on outcome measures in massively burned children. <i>Journal of Burn Care and Research</i> . 2015;1):S126.	5
782	Boyce ST, Simpson PS, Rieman MT, Warner P, Yakuboff KP, Bailey JK, et al. Autologous engineered skin substitutes reduce mortality and harvesting of skin graft donor sites for closure of extensive, full-thickness burns. <i>Journal of Burn Care and Research</i> . 2015;1):S107.	5
783	Boyce ST, Lloyd CM, Supp DM. Restoration of natural skin color by transplantation of cryopreserved human melanocytes in engineered skin substitutes. <i>Journal of Burn Care and Research</i> . 2015;1):S83.	5
784	Jerrom R, Jackson D, Patel A. From ancient Indian potters to WWII pilots: The elaborate history of skin grafting. <i>Journal of the American Academy of Dermatology</i> . 2015;1):AB26.	5
785	Shoae-Hassani A, Hamidieh AA, Mohseni R, Keyhanvar P, Azimi A, Mortazavi Tabatabaei SA, et al. Human adipocyte derived mesenchymal stem cell supports keratinocyte growth in a modified collagen-hyaluronic acid matrix. <i>Bone Marrow Transplantation</i> . 2015;1):S558.	5
786	Wood F, Fowler B, Tuckerman J, Rea S. Cultured epithelial autograft in the treatment of acute burn wounds. <i>Cochrane Database of Systematic Reviews</i> . 2014;2014 (7) (no pagination)(CD005988).	5
787	Zuber A, Borowczyk J, Zimolag E, Krok M, Madeja Z, Pamula E, et al. Poly(L-lactide-co-glycolide) thin films can act as autologous cell carriers for skin tissue engineering. <i>Cellular and Molecular Biology Letters</i> . 2014;19(2):297-314.	4
788	Blackstone BN, Drexler JW, Powell HM. Tunable engineered skin mechanics via coaxial electrospun fiber core diameter. <i>Tissue Engineering - Part A</i> . 2014;20(19-20):2746-55.	4
789	Kolli S, Ahmad S, Mudhar HS, Meeny A, Lako M, Figueiredo FC. Successful application of ex vivo expanded human autologous oral mucosal epithelium for the treatment of total bilateral limbal stem cell deficiency. <i>Stem Cells</i> . 2014;32(8):2135-46.	4
790	Wang X, Hu X, You C, Han C. Delivery of epithelial autografts and cutaneous wound healing in burn patients. <i>Burns</i> . 2014;40(1):166-8.	5
791	Gardien KLM, Middelkoop E, Ulrich MMW. Progress towards cell-based burn wound treatments. <i>Regen</i> . 2014;9(2):201-18.	6
792	Zhang DM, Hong WS, Fu LF, Wei XD, Xu AE. A randomized controlled study of the effects of different modalities of narrow-band ultraviolet B therapy on the outcome of cultured autologous melanocytes transplantation in treating vitiligo. <i>Dermatologic Surgery</i> . 2014;40(4):420-6.	1
793	Ai JX, He LX, Yang RH, Ya ZM. Constructing autologous composite skin around the expander for the repair of skin defects. [Chinese]. <i>Chinese Journal of Tissue Engineering Research</i> . 2014;18(20):3139-43.	7
794	Li XS, Du J, Song ZL, Li WP. Bone marrow mesenchymal stem cells for epidermal and skin appendage regeneration. [Chinese]. <i>Chinese Journal of Tissue Engineering Research</i> . 2014;18(6):853-9.	7
795	Kaviani M, Geramizadeh B, Rahsaz M, Marzban S. Consideration in the improvement of human epidermal keratinocytes culture in vitro. <i>Experimental and Clinical Transplantation</i> . 2014;2):295-6.	5

연번	서지정보	배제 사유
796	Figueiredo FC, Kolli S, Ahmad S, Lako M. Long-term outcomes of ex vivo expanded limbal stem cell transplantation in humans. <i>Cytotherapy</i> . 2014;4):S15.	5
797	Martin YH, Brown RA, Metcalfe AD. Microcarriers and compressed collagen to improve delivery of autologous keratinocytes in the treatment of severe burn injuries. <i>Wound Repair and Regeneration</i> . 2014;22 (5):A91.	5
798	Akershoek JJ, Vlig M, Talhout W, Boekema B, Richters CD, Ulrich MMW. Cell therapy for full-thickness wounds: Are fetal dermal cells a potential source? <i>Wound Repair and Regeneration</i> . 2014;22 (5):A73.	5
799	Moulin VJ, Duranceau L, Genest H, Bortoluzzi P, Auger FA, Germain K. Grafting of autologous skin (dermis and epidermis) substitutes reconstructed using a tissue engineering method: A success on burn patients. <i>Wound Repair and Regeneration</i> . 2014;22 (2):A53.	5
800	Chang PH. Treatment of foreign pediatric patients with chronic burn wounds with a delayed closure protocol. <i>J Am Coll Surg</i> . 2014;1):e42.	5
801	Crespo S, Malik G, Moro R, Martinez-Santamaria L, Gomez De Salazar M, Rodriguez L, et al. Assessment of the performance and fate of adipose-derived mesenchymal stem cells forming part of bioengineered skin equivalents in vitro and in vivo. <i>J Invest Dermatol</i> . 2014;2):S99.	5
802	Jerrom R, Jackson D, Patel A, Varma S. From ancient Indian potters to WWII pilots: The elaborate history of skin grafting. <i>Br J Dermatol</i> . 2014;1):141.	5
803	Gratz K, Comer A, Pirnstill J, Pirnstill S, LaDuc T, Vander Zanden M, et al. Advanced development of a human skin substitute for severe burns. <i>Journal of Burn Care and Research</i> . 2014;1):S168.	5
804	Sakamoto M, Morimoto N, Inoie M, Ogino S, Jinno C, Kawai K, et al. Cultured epidermis accelerates wound healing in combination with meshed skin grafting. <i>Journal of Burn Care and Research</i> . 2014;1):S157.	5
805	Baradar Jalili R, Curran T, Farrokhi A, Ghahary A. Ex vivo generated regulatory t cells have potent inhibitory function with a two-week life span in vivo. <i>Journal of Burn Care and Research</i> . 2014;1):S96.	5
806	Mochida K, Amano M, Iikawa M, Ishii Y, Setoyama M. Use of autologous cultured epidermal grafts as skin replacement after burn injury: Two cases report. [Japanese]. <i>Nishinohon Journal of Dermatology</i> . 2013;75(4):350-6.	7
807	Gibbs S. New developments in wound healing. <i>Nederlands Tijdschrift voor Dermatologie en Venereologie</i> . 2013;23(9):551-5.	5
808	O'Halloran E, Stewart N, Vetrichevel TP, Rea S, Wood F. Sweet's syndrome mimicking alkali burn: A clinical conundrum. <i>Journal of Plastic, Reconstructive and Aesthetic Surgery</i> . 2013;66(6):867-9.	6
809	Lootens L, Brusselaers N, Beele H, Monstrey S. Keratinocytes in the treatment of severe burn injury: An update. <i>Int Wound J</i> . 2013;10(1):6-12.	6
810	Catalano F, Malvasio V, Stella M. Chemical burns: A 5-years experience in the Turin Burn Center. <i>Journal of Plastic Dermatology</i> . 2013;9(2):145-8.	2
811	Wong M, Chua A, Tan BK. Cultured epithelial autografts for the coverage of large wounds: Minimizing skin graft donor sites in the sick patient. <i>European Journal of Plastic Surgery</i> . 2013;36(6):371-6.	6
812	Dearman BL, Stefani K, Li A, Greenwood JE. Take of a polymer-based autologous cultured composite skin on an integrated temporizing dermal matrix: Proof of concept. <i>Journal of Burn Care and Research</i> . 2013;34(1):151-60.	4
813	Medina A, Tredget EE. Strategies to increase flap survival in nasal reconstruction in patients with deep panfacial burns. <i>Journal of Burn Care and Research</i> . 2013;34(1):e42-e7.	5

연번	서지정보	배제 사유
814	Vashi NA, Kundu RV. Facial hyperpigmentation: Causes and treatment. Br J Dermatol. 2013;169(SUPPL. 3):41-56.	6
815	De Luca M. Biological parameters determining the clinical outcome of autologous cultures of limbal stem cells. Regen. 2013;1):2.	5
816	Anonymous. 23rd Annual Meeting of the European Tissue Repair Society. Wound Repair and Regeneration Conference: 23rd Annual Meeting of the European Tissue Repair Society Reims France Conference Publication:. 2013;21(6).	6
817	Verween G, Draye JP, Boone MALM, Aiti A, Verbeken G, De Vos D, et al. Cultivation and storage of several human skin cell types to repopulate acellular human dermal matrices. Wound Repair and Regeneration. 2013;21 (6):A87.	5
818	Ray S, Dheansa B, Metcalfe A, Sharpe J. The effect of microcarrier-delivered keratinocytes and single-stage matriderm on wound contraction and epithelialization. Wound Repair and Regeneration. 2013;21 (6):A79.	5
819	Gardien KLM, Marck RE, Bloemen MCT, Waaijman T, Gibbs S, Middelkoop E, et al. Cultured autologous keratinocytes in combination with a split skin autograft for burn wound treatment: Preliminary results of a prospective randomized controlled multicenter study. Wound Repair and Regeneration. 2013;21 (6):A65.	5
820	Alexaline M, Nivet M, Thepenier C, Leclerc T, Bey E, Saint-Blancard P, et al. Evaluation of cultured epithelial autografts for massive burn treatment. Wound Repair and Regeneration. 2013;21 (6):A56.	5
821	Alexaline M, Trouillas M, Nivet M, Thepenier C, Leclerc T, Bey E, et al. Development of a preclinical model of third-degree burn. Wound Repair and Regeneration. 2013;21 (6):A55.	5
822	English IJT, Ziembicki JA, Corcos A, Varcelotti J, Esteban-Vives R, Gerlach JC. Characterization of skin cell preparations for autologous single-cell spray-grafting of deep partial thickness burns. Journal of Burn Care and Research. 2013;1):S191.	6
823	Jalili RB, Curran TA, Kilani RT, Hartwell R, Ghahary A. A novel strategy for induction of immune tolerance in skin allotransplantation for burn patients. Journal of Burn Care and Research. 2013;1):S119.	5
824	Dickinson A, Lako M, Kolli S, Figueiredo FC. Outcomes of ex vivo expanded limbal stem cell transplantation in humans. Cytotherapy. 2013;1):S44-S5.	5
825	Then KY, Umi Kalthum MN, Wong CY, Ooi GC. Autologous cultivated oral mucosal epithelial cells transplantation (COMET) with amniotic membrane based sheets for severe ocular chemical injury. Cytotherapy. 2013;1):S11.	5
826	Mahjour SB, Ghaffarpasand F, Wang H. Hair follicle regeneration in skin grafts: Current concepts and future perspectives. Tissue Engineering - Part B: Reviews. 2012;18(1):15-23.	6
827	Tenenhaus M, Rennekampff HO. Surgical Advances in Burn and Reconstructive Plastic Surgery. New and Emerging Technologies. Clin Plast Surg. 2012;39(4):435-43.	6
828	Alexaline M, Thepenier C, Trouillas M, Leclerc T, Desmouliere A, Benderriter M. Cell therapy of burns. Regenerative Medicine and Cell Therapy. 2012;Biomedical and Health Research.:406-17.	5
829	Gomez CMA, Soto GJM, Munera LMR. Skin substitutes developed by tissue engineering. Iatreia. 2012;25(1):42-53.	5
830	Ching JA, Smith Jr DJ. Levamisole-induced necrosis of skin, soft tissue, and bone: Case report and review of literature. Journal of Burn Care and Research. 2012;33(1):e1-e5.	6
831	Orman JA, Chung KK, King BT, Jones JA, Cancio LC, Renz EM, et al. The hot 100: An analysis of the highest impact burn papers in the last 55 years. Journal of Burn Care and Research. 2012;1):S196.	5

연번	서지정보	배제 사유
832	Greenwood J, Dearman B, Li A. Wound closure over integrated polymer neo-dermes using cultured epithelial autograft and composite cultured 'skin equivalents'. Journal of Burn Care and Research. 2012;1):S173.	5
833	Rodriguez Sr P, Lule I, Carranza L. Wounds healing of partial thickness burns with biological dressing of cultured human keratinocytes live (epifast). Journal of Burn Care and Research. 2012;1):S172.	5
834	Gondo M, Matsumura H, Shibata D, Segawa M, Shimada K, Ida Y, et al. Chronological histological findings of cultured epidermal autograft over artificial dermis focusing on basement membrane proteins. Journal of Burn Care and Research. 2012;1):S172.	5
835	Coyne NL, Foster KN, Peck MD, Matthews MR, Pressman MA, Caruso DM. Are cultured epidermal autografts worth the cost? Journal of Burn Care and Research. 2012;1):S163.	5
836	Rawlins JM. Early surgery for all pediatric scalds: Developing a dermal preserving surgical algorithm for the benefit of patients and Health Care Providers. Journal of Burn Care and Research. 2012;1):S144.	5
837	Hawkins HK, Adler-Baugh K, Nguyen A, Huang T, Lee J, Herndon DN, et al. Preliminary report: Chronic ulcerations and the incidence of squamous cell carcinoma in burn patients. Journal of Burn Care and Research. 2012;1):S139.	5
838	Curran T, Jalili R, Ghahary A. Immuno-protective role of CD4+CD25-converted to regulatory t cells in allogenic skin graft. Journal of Burn Care and Research. 2012;1):S112.	5
839	Antoni FH, Paul MV, Kaur P, Cleland H, Akbarzadeh S. Investigation of human adult dermal pericytes and their possible role in skin tissue regeneration. Wound Repair and Regeneration. 2012;20 (5):A69.	5
840	Verween G, Draye JP, Colemonts-Vroninks H, Verbeken G, Rose T, Jennes S, et al. Culturing growth rate improvement of neonatal foreskin keratinocytes with a new defined animal origin-free cell culture medium supplement. Wound Repair and Regeneration. 2012;20 (5):A118.	5
841	Trouillas M, Alexaline M, Gourven M, Thepenier C, Prat M, Lataillade JJ. Preclinical animal model of third-degree burn for burn repair. Wound Repair and Regeneration. 2012;20 (5):A115.	5
842	Lataillade JJ, Thepenier C, Alexaline M, Leclerc T, Trouillas M. Interest of mesenchymal stromal cells in cell and tissue therapy for severe burns. Wound Repair and Regeneration. 2012;20 (5):A100.	5
843	Caviggioli F, Ibatci A, Vinci V, Klinger M. Scar treatment using autologous fat graft. Wound Repair and Regeneration. 2012;20 (5):A87.	5
844	Batista FRX, Rehder J, Puzzi MB. Epidermal substitutes: Advances in cell culture technology resulting in a success on clinical applications. New Biotechnology. 2012;1):S115.	5
845	Sobhani Z, Mahdavi-Mazdeh M, Ayaz M, Tavakoli SAH, Nozary Heshmati B, Khodadadi A, et al. The use of partial thickness skin allograft as a temporary biologic dressing of burns. Transplantation. 2012;10S):488.	5
846	Wolfe L. Production of cell therapies from clinical trial to commercial distribution. Journal of Tissue Engineering and Regenerative Medicine. 2012;1):426.	5
847	Marck RE, Bloemen MCT, Waaijman T, Gibbs S, Middelkoop E, Ulrich MMVV. Cultured keratinocytes for burn wound treatment, preliminary results of a prospective randomized controlled intra-patient comparative study. Journal of Tissue Engineering and Regenerative Medicine. 2012;1):101.	5
848	Meuli M. Keynote: Skingeneering for large full thickness skin defects. Journal of Tissue Engineering and Regenerative Medicine. 2012;1):101.	5

연번	서지정보	배제 사유
849	Finnerty C, McCauley R, Hawkins H, Herndon D. Genomic alterations in cultured epithelial autograft (CEA) associated with the development of squamous cell carcinoma. <i>Journal of Tissue Engineering and Regenerative Medicine</i> . 2012;1):94.	5
850	Pikula M, Langa P, Drukala J, Kondej K, Hedrych-Ozimina A, Graczyk-Aszkielowicz M, et al. Isolation and culture of human keratinocytes and epidermal stem cells for clinical purposes. <i>Journal of Tissue Engineering and Regenerative Medicine</i> . 2012;1):92.	5
851	Tiede S, Lehmann J, Nagel S, A EP, Vidali S, Siemers F, et al. Cell-based regenerative medicine by nestin+ progenitors: Stimulation of reepithelialization, angiogenesis, and granulation tissue formation in human skin organ culture. <i>Experimental Dermatology</i> . 2012;21 (3):e9.	5
852	Moravvej H, Kalantar Hormozi A, Hosseini S, Mahmoudi Rad M, Abdolzadeh M. Comparison of the application of allogeneic fibroblast and autologous mesh grafting with conventional method in the treatment of third-degree burns. <i>Wound Repair and Regeneration</i> . 2012;20 (2):A32.	5
853	Curran T, Jalili R, Ghahary A. Immunoprotective role of IDO microenvironment generated regulatory T cells in vivo. <i>Wound Repair and Regeneration</i> . 2012;20 (2):A20.	5
854	Anonymous. Delivery of autologous keratinocytes on gelatin microcarriers to reduce contraction of full-thickness wounds. <i>Regen</i> . 2012;7(1):31-2.	6
855	Greenwood JE, Dearman BL. Comparison of a sealed, polymer foam biodegradable temporizing matrix against integra dermal regeneration template in a porcine wound model. <i>Journal of Burn Care &amp; Research</i> . 2011;13.	4
856	Lee ST, Ma DR, Chua A, Ting YX, Song C. Translational journey of epidermal stem cells- from cultured epithelial autograft to induced pluripotent stem cells. <i>Proceedings of Singapore Healthcare</i> . 2011;1):219.	5
857	Ahrari I, Khosravi Maharlooe M, Mohamadi A, Monabati A. Comparison of two methods cell spray and injection of epithelial cell suspension for the treatment of postburn hypopigmentation. <i>Cell j</i> . 2011;1):37-8.	5
858	Eldardiri M, Martin Y, Sharpe J. Culture and delivery of autologous keratinocytes on microcarriers for cutaneous repair. <i>European Cells and Materials</i> . 2011;2):68.	5
859	Chang J, Davids LM. An investigation into the mechanisms of repigmentation in vitiligo and partial-thickness burn wounds. <i>Clinical Anatomy</i> . 2011;24 (7):933.	5
860	Esteban R, Young M, Over P, Chinnici C, Amico G, Gerlach J. Delivery of epidermal and dermal stem cells to a burn wound via a spray device - Exploring human fetal skin progenitor cells for regenerative medicine cell based therapy. <i>Molecular Biology of the Cell Conference: Annual Meeting of the American Society for Cell Biology, ASCB</i> . 2011;22(24).	5
861	Rennekampff H, Herold C, Vogt M. Keratinocyte suspension for the treatment of facial burns. <i>Burns</i> . 2011;1):S17.	5
862	Brucato V, Carfi Pavia F, La Carrubba V, Ghersi G, Rigogliuso S, Conte F. Pre-vascularized PLLA scaffolds: A new approach to develop deep tissue regeneration. <i>Connective Tissue Research</i> . 2011;52 (4):275-7.	5
863	Martinez FF, Barrera LA, Lomeli RCY, Estrada VE, Sandoval ZH, Villegas CH, et al. Gene therapy approaches based on Bcl-2 to increased resistance of human cultured keratinocytes for transplantation. <i>Molecular Therapy</i> . 2011;19 (7):1373.	5
864	Rama P, Matuska S, Paganoni G, Spinelli A, De Luca M, Pellegrini G. Limbal stem-cell therapy and long-term corneal regeneration. <i>Hum Gene Ther</i> . 2011;22 (6):A8.	5
865	Henderson M, Lim H, Hamzavi I, Huggins R. Case report: Autologous noncultured melanocyte-keratinocyte transplan-tation procedure in an African American male with postburn injury hypopigmentation. <i>Journal of the American Academy of Dermatology</i> . 2011;1):AB143.	5



연번	서지정보	배제 사유
866	Gomez C, Galan JM, Torrero V, Ferreiro I, Perez D, Palao R, et al. Use of an autologous bioengineered composite skin in extensive burns: Clinical and functional outcomes. A multicentric study. <i>Burns</i> . 2011;37(4):580-9.	2
867	Lepow BD, Downey M, Yurgelon J, Klassen L, Armstrong DG. Bioengineered tissues in wound healing: A progress report. <i>Expert Review of Dermatology</i> . 2011;6(3):255-62.	6
868	O'Neill TB, Rawlins JM, Rea S, Wood FM. Methamphetamine laboratory-related burns in Western Australia - Why the explosion? <i>Burns</i> . 2011;37(6):1044-8.	2
869	Supp DM. Skin substitutes for burn wound healing: Current and future approaches. <i>Expert Review of Dermatology</i> . 2011;6(2):217-27.	6
870	Chew HF. Limbal stem cell disease: Treatment and advances in technology. <i>Saudi Journal of Ophthalmology</i> . 2011;25(3):213-8.	6
871	Lavoie A, Fugere C, Fradette J, Larouche D, Paquet C, Beuparlant A, et al. Considerations in the choice of a skin donor site for harvesting keratinocytes containing a high proportion of stem cells for culture in vitro. <i>Burns</i> . 2011;37(3):440-7.	4
872	Singer NG, Caplan AI. Mesenchymal stem cells: Mechanisms of inflammation. <i>Annual Review of Pathology: Mechanisms of Disease</i> . 2011;6:457-78.	5
873	Bhatia VY, Singh M. Innovative technique using heterogenous collagen and intermingled autologous split-thickness skin graft for coverage of raw areas in deep burns with paucity of donor site as a life-saving measure. <i>European Journal of Plastic Surgery</i> . 2011;34(2):93-7.	2
874	Beier JP, Boos AM, Kamolz L, Vogt PM, Koller R, Horch RE. Skin tissue engineering - From split skin to engineered skin grafts?. [German]. <i>Handchirurgie Mikrochirurgie Plastische Chirurgie</i> . 2010;42(6):342-53.	7
875	Ha J, Heath CH, Suzanne R, Wood F. Ophthalmic manifestation of candida: Case report and review of the literature. <i>Eur J Pediatr Surg</i> . 2009;19(3):197-9.	1
876	Tan KKB, Chan JKY, Lane B. Characterisation of foetal keratinocytes for skin grafting. <i>Annals of the Academy of Medicine Singapore</i> . 2010;1):S94.	5
877	Pataia E, Arleo S, Somma F, Menichini G, D'Antonio A, Alfano C. Use of biotechnologies in cutaneous injuries limb repair. <i>European Surgical Research</i> . 2010;45 (3-4):228.	5
878	De Luca M, Rama P, Mavilio F, Pellegrini G. Human epithelial stem cells in corneal regeneration and epidermal gene therapy. <i>FASEB Journal Conference: Experimental Biology</i> . 2010;24(Meeting Abstracts).	5
879	Nacchiero E, Ranieri M, Debellis L, Colella M, Maggio G, Pascone M. Calcium signalling in keratinocytes and fibroblasts grown in low-calcium medium. <i>European Journal of Clinical Investigation</i> . 2010;1):66.	5
880	Khoo TL, Halim AS, Saad AZM, Dorai AA. The application of glycerol-preserved skin allograft in the treatment of burn injuries: An analysis based on indications. <i>Burns</i> . 2010;36(6):897-904.	2
881	Shortt AJ, Tuft SJ, Daniels JT. Ex vivo cultured limbal epithelial transplantation. A clinical perspective. <i>Ocular Surface</i> . 2010;8(2):80-90.	6
882	Danks RR, Lairet K. Innovations in caring for a large burn in the Iraq war zone. <i>Journal of Burn Care and Research</i> . 2010;31(4):665-9.	6
883	Koch N, Wiesner L, Benathan M, Rotman S, Darwiche S, Raffoul W. Chronic rejection of allogenic dermal grafts in 73% TBSA burn patient. <i>Burns</i> . 2009;1):S47.	5
884	De Angelis B, Lucarini L, Montone A, Cervelli V. ReCell: A new in device for skin autologous inplants in the treatment of burns and scars. <i>Burns</i> . 2009;1):S36.	5
885	Lindford AJ, Frey I, Vuola J, Koljonen V. Review of the helsinki skin bank 2001-2008. <i>Burns</i> . 2009;1):S35.	5

연번	서지정보	배제 사유
886	Cirodde A, Leclerc T, Jault P, Bargues L, Le Bever H, Carsin H. A retrospective study of the use of cultured epidermal autografts (CEA) in severe burn patients. <i>Burns</i> . 2009;1):S19.	5
887	Sood S, Roggy D, Zieger M, Mir H, Coleman J. Cultured epithelial autografts (CEA) for coverage of large burn wounds in 88 patients: The Indiana University experience. <i>Burns</i> . 2009;1):S17.	5
888	Salib GE, Duranceau L, Armour AD. Use of cultured epithelial autograft on donor sites of burn patients. <i>Burns</i> . 2009;1):S13.	5
889	Majd H, Pietramaggiore G, Pittet-Cuenod B, Quinn T, Hinz B. Culture on highly expandable and compliant surfaces accelerates the expansion of non-fibroblastic fibroblasts for cell therapy. <i>Burns</i> . 2009;1):S13.	5
890	Ums M, Derow E, Lohberg L, Schuler G, Hartmann A. Keratinocytes from different compartments of human skin-defining conditions for stem cell protection. <i>Hum Gene Ther</i> . 2009;20 (11):1448-9.	5
891	James SE. The clinical application of cultured cells - Are they permanent or just an effective temporary dressing? The FASEB Journal Conference: Experimental Biology. 2009;23(S1).	5
892	Vaughan MB, Andrews CM, Wright WE, Shay JW. Keratinocyte senescence effects on differentiation and migration in a skin equivalent. The FASEB Journal Conference: Experimental Biology. 2009;23(S1).	5
893	Auger FA, Gibot L. In vivo remodeling of a vascularized human skin substitute. <i>Microcirculation</i> . 2009;16 (8):750.	5
894	Blais M, Grenier M, Berthod F. Enrichment of tissue-engineered skin with Schwann cells improves cutaneous nerve regeneration. <i>J Invest Dermatol</i> . 2009;1):S2.	5
895	Isaev AA, Prihodko AV, Saburina IN, Kiselev SL, Melihova VS, Potapov IV, et al. Developing new immunological compatible transplants derived from umbilical cord. <i>Cellular Therapy and Transplantation</i> . 2009;2 (5):65.	5
896	Kita K, Finnerty CC, Herndon DN, Jeschke MG. Characterization and immunogenicity of human umbilical cord-derived mesenchymal stem cells. <i>Surgical Infections</i> . 2009;10 (2):192.	5
897	Pi YL, Tang WQ, Lu JY, Dong Y. Tissue-engineered corneal epithelium transplantation for treatment of corneal alkaline burn in rabbits: Morphological observation. <i>Journal of Clinical Rehabilitative Tissue Engineering Research</i> . 2009;13(41):8011-6.	4
898	Schrader S, Notara M, Beaconsfield M, Tuft SJ, Daniels JT, Geerling G. Tissue engineering for conjunctival reconstruction: Established methods and future outlooks. <i>Current Eye Research</i> . 2009;34(11):913-24.	6
899	Peral MC, Huaman Martinez MA, Valdez JC. Bacteriotherapy with <i>Lactobacillus plantarum</i> in burns. <i>Int Wound J</i> . 2009;6(1):73-81.	2
900	Negri S, Federici G, Farinato S, Fila C. Human plasma as a dermal scaffold for the generation of a completely autologous bioengineered skin. <i>Journal of Clinical Rehabilitative Tissue Engineering Research</i> . 2009;13(47):9211-6.	4
901	Khodadadi L, Shafieyan S, Aghdami N, Baharvand H. Cell therapy in burn repair. <i>Yakhteh</i> . 2008;10(3):167-78.	6
902	Chen JP, Liu DW, Mao YG. Structure and properties of tissue-engineered skin. [Chinese]. <i>Journal of Clinical Rehabilitative Tissue Engineering Research</i> . 2008;12(24):4731-4.	7
903	Constantinides J, Misra A, Nassab R, Wilson Y. <i>Absidia corymbifera</i> fungal infection in burns: A case report and review of the literature. <i>Journal of Burn Care and Research</i> . 2008;29(2):416-9.	2

연번	서지정보	배제 사유
904	Fear M. Reply letter to: "The use of non-cultured autologous cell suspension and integrin dermal regeneration template to repair full thickness skin wounds in a porcine skin model: A one-step process" [Burns 33(6) (2007) 693-700]. Burns. 2008;34(6):900-1.	5
905	Safari M, Ghorbani R, Emami M, Yousefi B, Ghahari L. The effects of GH and PDGF on the growth and proliferation of the epithelial cells. [Arabic]. Koomesh. 2008;9(3):229-35+33.	7
906	Li DQ, Huang XY, Yang XH. Morphology of human skin fibroblasts after heat injury in vitro. [Chinese]. Journal of Clinical Rehabilitative Tissue Engineering Research. 2008;12(11):2056-60.	7
907	Karchilaki I, Topakas G, Castana O, Sotiriou P, Michelakis D, Alexakis D, et al. The use of cultured autologous fibroblasts in burn wounds healing process. Burns. 2007;33(6):791-2.	6
908	Tenenhaus M, Bhavsar D, Rennekampff HO. Treatment of deep partial thickness and indeterminate depth facial burn wounds with water-jet debridement and a biosynthetic dressing. Injury. 2007;38(5 SUPPL.):S38-S44.	2
909	Wong T, McGrath JA, Navsaria H. The role of fibroblasts in tissue engineering and regeneration. Br J Dermatol. 2007;156(6):1149-55.	6
910	Rochat A, Claudinot S, Nicolas M, Barrandon Y. Stem cells and skin engineering. Swiss Medical Weekly. 2007;137(SUPPL. 155):49S-54S.	6
911	Tominac Trcin M. Tissue bank of the University Hospital of Traumatology in Zagreb, Croatia. [Croatian]. Acta Medica Croatica. 2007;61(5):497-503.	7
912	Arnez ZM, Toffanetti F, Papa G, Ramella V, Renzi N, Panizzo N. Homologous skin and tissue substitutes in burn patients. [Italian]. Rivista Italiana di Chirurgia Plastica. 2007;39(3-4):177-82.	7
913	Holland AJA. Pediatric burns: The forgotten trauma of childhood. Canadian Journal of Surgery. 2006;49(4):272-7.	6
914	Inatomi T, Nakamura T, Koizumi N, Sotozono C, Yokoi N, Kinoshita S. Midterm results on ocular surface reconstruction using cultivated autologous oral mucosal epithelial transplantation. Am J Ophthalmol. 2006;141(2):267-75.e1.	1
915	Fu X, Qu Z, Sheng Z. Potentiality of Mesenchymal Stem Cells in Regeneration of Sweat Glands. J Surg Res. 2006;136(2):204-8.	2
916	Zhu N, Warner RM, Simpson C, Glover M, Hernon CA, Kelly J, et al. Treatment of burns and chronic wounds using a new cell transfer dressing for delivery of autologous keratinocytes. European Journal of Plastic Surgery. 2005;28(5):319-30.	1
917	Atiyeh BS, Gunn SW, Hayek SN. State of the art in burn treatment. World J Surg. 2005;29(2):131-48.	6
918	Atiyeh BS, Hayek SN, Gunn SW. New technologies for burn wound closure and healing - Review of the literature. Burns. 2005;31(8):944-56.	6
919	Chandra A, Maurya OPS, Reddy B, Kumar G, Pandey K, Bhaduri G. Amniotic membrane transplantation in ocular surface disorders. Journal of the Indian Medical Association. 2005;103(7):364-8.	1
920	Fernandes M, Sridhar MS, Sangwan VS, Rao GN. Amniotic membrane transplantation for ocular surface reconstruction. Cornea. 2005;24(6):643-53.	6
921	Elsinghorst Th AM. New findings on animal diseases published since 2003. 1. Dogs. Veterinary Quarterly. 2004;26(1):19-24.	6
922	Chester DL, Papini RPG. Skin and skin substitutes in burn management. Trauma. 2004;6(2):87-99.	6
923	Lam PK, Chan ESY, Ho WS, Liew CT. In vitro cytotoxicity testing of a nanocrystalline silver dressing (Acticoat) on cultured keratinocytes. British Journal of Biomedical Science. 2004;61(3):125-7.	4

연번	서지정보	배제 사유
924	Boyce ST. Fabrication, quality assurance, and assessment of cultured skin substitutes for treatment of skin wounds. <i>Biochemical Engineering Journal</i> . 2004;20(2-3):107-12.	4
925	Takami Y, Shimazaki S, Yamaguchi R, Tanaka H, Harii K. Transplantation of tissue-engineered skin composed of autologous cells and acellular allogeneic dermal matrix. [Japanese]. <i>Japanese Journal of Plastic and Reconstructive Surgery</i> . 2004;47(8):867-73.	7
926	Wisser D, Steffes J. Skin replacement with a collagen based dermal substitute, autologous keratinocytes and fibroblasts in burn trauma. <i>Burns</i> . 2003;29(4):375-80.	6
927	Wang TW, Huang YC, Sun JS, Lin FH. Organotypic keratinocyte-fibroblast cocultures on a bilayer gelatin scaffold as a model of skin equivalent. <i>Biomed Sci Instrum</i> . 2003;39:523-8.	2
928	Barillo DJ, McManus AT, Cancio LC, Sofer A, Goodwin CW. Burn center management of necrotizing fasciitis. <i>Journal of Burn Care and Rehabilitation</i> . 2003;24(3):127-32.	2
929	Hutmacher DW, Vanscheidt W. Matrices for tissue-engineered skin. <i>Drugs of Today</i> . 2002;38(2):113-33.	6
930	Van den Bogaerd AJ, Van Zuijlen PPM, Van Galen M, Lamme EN, Middelkoop E. The suitability of cells from different tissues for use in tissue-engineered skin substitutes. <i>Archives of Dermatological Research</i> . 2002;294(3):135-42.	2
931	Kangesu T. Study of dermal grafts and cultured autologous keratinocytes in an experimental model. <i>Ann R Coll Surg Engl</i> . 2001;83(3):197-202.	1
932	Ho WS. Skin substitutes: An overview. <i>Annals of the College of Surgeons of Hong Kong</i> . 2002;6(4):102-8.	6
933	Burd A, Chan E. Keratinocyte-keloid interaction. <i>Plastic and Reconstructive Surgery</i> . 2002;110(1):197-202.	6
934	Matsuzaki K, Kumagai N, Inoue H, Oshima H, Tanabe M. The evolution of allogeneic cultured keratinocyte treatments; clinical experiences and future possibilities. [Japanese]. <i>Japanese Journal of Plastic and Reconstructive Surgery</i> . 2002;45(7):611-7.	7
935	Grant I, Warwick K, Marshall J, Green C, Martin R. The co-application of sprayed cultured autologous keratinocytes and autologous fibrin sealant in a porcine wound model. <i>Br J Plast Surg</i> . 2002;55(3):219-27.	4
936	Sawa M. Current clinical and research approach in keratoconjunctival disorders. [Japanese]. <i>Japanese Journal of Clinical Ophthalmology</i> . 2002;56(3):229-40.	7
937	Cochran A, Morris SE, Edelman LS, Saffle JR. Systemic Candida infection in burn patients: A case-control study of management patterns and outcomes. <i>Surgical Infections</i> . 2002;3(4):367-74.	2
938	Supp DM, Boyce ST. Overexpression of vascular endothelial growth factor accelerates early vascularization and improves healing of genetically modified cultured skin substitutes. <i>Journal of Burn Care and Rehabilitation</i> . 2002;23(1):10-20.	4
939	Chan ESY, Lam PK, Liew CT, Lau HCH, Yen RSC, King WWK. A new technique to resurface wounds with composite biocompatible epidermal graft and artificial skin. <i>Journal of Trauma - Injury, Infection and Critical Care</i> . 2001;50(2):358-62.	6
940	Bello YM, Falabella AF. Use of skin substitutes in dermatology. <i>Dermatologic Clinics</i> . 2001;19(3):555-61.	6
941	Henckel-Donnersmarck G, Schmiedl S, Horbrand F, Bohmer R, Muhlbauer W. The use of cultured epidermal grafts in treating deep dermal burn wounds. [German]. <i>Aktuelle Traumatologie</i> . 2001;31(5):216-20.	7
942	Balasubramani M, Kumar TR, Babu M. Skin substitutes: A review. <i>Burns</i> . 2001;27(5):534-44.	6
943	Carsin H. Letter to the Editor. <i>Burns</i> . 2001;27(4):418-9.	5
944	Inokuchi S, Houjou M, Kidokoro M, Tanino R. Treatment of full thickness burn wounds with cultured autologous composite skin grafts. [Japanese]. <i>Japanese Journal of Plastic and Reconstructive Surgery</i> . 2000;43(6):581-7.	7

연번	서지정보	배제 사유
945	Yanaga H, Tai Y. Clinical application of combined dermabrasion and autologous cultured epidermal grafting to the large scar and post skin grafting area. [Japanese]. Japanese Journal of Plastic and Reconstructive Surgery. 2000;43(6):541-6.	7
946	Toriyama K, Kazeto T, Hata K, Ueda M, Torii S. Clinical results of cultured epithelial grafts. [Japanese]. Japanese Journal of Plastic and Reconstructive Surgery. 2000;43(6):557-62.	7
947	Kuroyanagi Y. Advances in cultured skin substitutes. [Japanese]. Japanese Journal of Plastic and Reconstructive Surgery. 2000;43(6):571-80.	7
948	Matsuzaki K, Kumagai N. Autologous cultured keratinocytes: Significance in plastic surgery. [Japanese]. Japanese Journal of Plastic and Reconstructive Surgery. 2000;43(6):533-40.	7
949	Tanaka H, Wada T, Suzuki T, Sakaki S, Shimazaki S. The present status of the skin bank network. [Japanese]. Japanese Journal of Plastic and Reconstructive Surgery. 2000;43(11):1103-10.	7
950	Nagase T, Kumagai N, Matsuzaki K, Tanabe M. Treatment of burns by grafting of cultured epithelium. [Japanese]. Japanese Journal of Plastic and Reconstructive Surgery. 2000;43(11):1091-6.	7
951	Koller R, Bierochs B, Bayer GS, Meissl G, Frey M. The role of cultivated epidermal auto- and allografts in the treatment of burn injuries. [German]. Acta Chirurgica Austriaca. 1997;29(6):331-5.	7
952	Meneguzzi G, Ortonne JP. Gene therapy and keratinocytes. [French]. Nouvelles Dermatologiques. 1999;18(10):654-6.	7
953	Stoner ML, Wood FM. Cultured epithelial autograft 'take' confirmed by the presence of cytokeratin 9 [1]. J Invest Dermatol. 1999;112(3):391-2.	5
954	Raghunath M, Tschodrich-Rotter M, Sasaki T, Meuli M, Chu ML, Timpl R. Confocal laser scanning analysis of the association of fibulin-2 with fibrillin-1 and fibronectin define different stages of skin regeneration. J Invest Dermatol. 1999;112(1):97-101.	4
955	Raghunath M, Unsold C, Kubitscheck U, Bruckner-Tuderman L, Peters R, Meuli M. The cutaneous microfibrillar apparatus contains latent transforming growth factor-beta binding protein-1 (LTBP-1) and is a repository for latent TGF-beta1. J Invest Dermatol. 1998;111(4):559-64.	4
956	Duinslaeger L, Delaey B, Vanderkelen A. Short- and long-term results of application of allogeneic cultured keratinocytes on burn wounds and burn scar. European Journal of Plastic Surgery. 1998;21(1):14-8.	2
957	Ng RLH, Woodward B, Bevan S, Green C, Martin R. Retroviral marking identifies grafted autologous keratinocytes in porcine wounds receiving cultured epithelium. J Invest Dermatol. 1997;108(4):457-62.	4
958	Fei X, Lee ST, Seah CS, Lim SM. Combined autologous and allogeneic (chimeric) keratinocytes grown on bovine fibrin glue. Asian Journal of Surgery. 1997;20(1):30-4.	6
959	Compton CC, Kupper TS, Nadire KB. HIV-infected Langerhans cells constitute a significant proportion of the epidermal Langerhans cell population throughout the course of HIV disease. J Invest Dermatol. 1996;107(6):822-6.	2
960	Rouabhia M. Skin replacement using chimeric (allogeneic-syngeneic) cultured epidermal sheets. [French]. Medecine/Sciences. 1996;12(12):1370-7.	7
961	Stoner ML, Wood FM. Systemic factors influencing the growth of cultured epithelial autograft. Burns. 1996;22(3):197-9.	4
962	Tanz R, Gottschalk E, Leppin F. Cultured epidermal autografts (CEA) in treatment of severe thermic-lesions in children. [German]. Zentralblatt fur Kinderchirurgie. 1996;5(2):105-8.	7
963	Myers S, Navsaria H, Sanders R, Green C, Leigh I. Transplantation of keratinocytes in the treatment of wounds. Am J Surg. 1995;170(1):75-83.	6

연번	서지정보	배제 사유
964	Dougherty WR, Chalabian JR. Skin substitutes. <i>Western Journal of Medicine</i> . 1995;162(6):540-1.	5
965	Horch R, Stark GB, Kopp J, Andree C. Keratinocyte-fibrin-glue suspension (KFGS) and conventional cultured epithelial sheet grafting (CEA). [German]. <i>Transplantationsmedizin: Organ der Deutschen Transplantationsgesellschaft</i> . 1995;7(2):99-103.	7
966	Stark GB, Kaiser HW, Horch R, Kopp J, Spilker G. Cultured autologous keratinocytes suspended in fibrin glue (KFGS) with allogenic overgraft for definitive burn wound coverage. <i>European Journal of Plastic Surgery</i> . 1995;18(6):267-71.	6
967	Compton CC, Tong Y, Trookman N, Zhao H, Roy D, Press W. Transforming growth factor alpha gene expression in cultured human keratinocytes is unaffected by cellular aging. <i>Arch Dermatol</i> . 1995;131(6):683-90.	4
968	Compton CC, Press W, Gill JM, Bantick G, Nadire KB, Warland G, et al. The generation of anchoring fibrils by epidermal keratinocytes: A quantitative long-term study. <i>Epithelial Cell Biol</i> . 1995;4(3):93-103.	3
969	Matsuzaki K, Kumagai N, Fukushi S, Ohshima H, Tanabe M, Ishida H, et al. Autologous cultured epithelium on the allogeneic dermis for massive burns. [Japanese]. <i>Japanese Journal of Plastic and Reconstructive Surgery</i> . 1995;38(2):193-200.	7
970	Wassermann D, Schlotterer M, Lebreton F. Modern management of burn injuries. [French]. <i>Medecine et Hygiene</i> . 1994;52(2023):961-9.	7
971	Rouabhia M, Germain L, Bergeron J, Auger FA. Successful transplantation of chimeric allogeneic-autologous cultured epithelium. <i>Transplant Proc</i> . 1994;26(6):3361-2.	5
972	Hickerson WL, Love Jr RT, Bishop JF. Cultured epidermal autografts: The current status. <i>Problems in General Surgery</i> . 1994;11(4):747-52.	6
973	Wendt JR, Ulich TR, Ruzics EP, Hostetler JR. Indefinite survival of human skin allografts in patients with long-term immunosuppression. <i>Ann Plast Surg</i> . 1994;32(4):411-7.	2
974	Phillips TJ. Biologic skin substitutes. <i>Journal of Dermatologic Surgery and Oncology</i> . 1993;19(8):794-800.	2
975	Vindenes H, Bjerknes R. The frequency of bacteremia and fungemia following wound cleaning and excision in patients with large burns. <i>Journal of Trauma</i> . 1993;35(5):742-9.	2
976	Grimaldi M, Cervelli V, Giudiceandrea F, Maschio MCE. Autologous cultured epidermal autografts for skin repair after burn injuries. [Italian]. <i>European Review for Medical and Pharmacological Sciences</i> . 1992;14(3):199-201.	7
977	Lorda Barraguer E, Pardo Mateu L, Tafalla Pena M, Aranda Lopez I. Considerations on 'in vitro' cultured epidermis and its clinical application. [Spanish]. <i>Cirurgia Plastica Ibero-Latinoamericana</i> . 1992;18(3):269-75.	7
978	Blight A, Mountford EM, Cheshire IM, Clancy JMP, Levick PL. Treatment of full skin thickness burn injury using cultured epithelial grafts. <i>Burns</i> . 1991;17(6):495-8.	2
979	Bassetto F, Toffanin M, Mazzoleni F, Conconi MT, Parnigotto PP. Influence of different supports on production and clinical utilization of keratinocyte sheets. [Italian]. <i>Rivista Italiana di Chirurgia Plastica</i> . 1991;23(3):279-82.	7
980	Petersen MJ, Woodley DT. Characterization of cellular elements in healed cultured keratinocyte autografts used to cover burn wounds (I: Reply). <i>Arch Dermatol</i> . 1990;126(8):1105.	5
981	Hergueter CA, O'Connor NE. Skin substitutes in upper extremity burns. <i>Hand Clin</i> . 1990;6(2):239-41.	5
982	Deglise B, Benathan M, Frenk E, Krupp S. First experience with treatment of burns by autologous cultured epidermal transplant. [French]. <i>Schweizerische Medizinische Wochenschrift</i> . 1987;117(37):1380-3.	7

연번	서지정보	배제 사유
983	Doherty D, Austin EV. Effective management of cultured epithelial cells. Two case reports. <i>Journal of Burn Care and Rehabilitation</i> . 1986;7(1):33-4.	6
984	Thivolet J, Faure M, Demidem A, Mauduit G. Cultured human epidermal allografts are not rejected for a long period. <i>Archives of Dermatological Research</i> . 1986;278(3):252-4.	4
985	Curtsinger ILJ, Schultz GS, Brown GL. Progress in burn therapy: Skin banking and epidermal culture. <i>Journal of the Kentucky Medical Association</i> . 1985;83(5):251-3.	6
986	Yannas IV, Hansbrough JF. What criteria should be used for designing artificial skin replacements and how well do the current grafting materials meet these criteria? <i>Journal of Trauma</i> . 1984;24(9 SUPPL.):S29-S39.	6
987	Steigleder GK. Coverage of wounds with autologous cultured human epithelium. A new mile stone. [German]. <i>H+G Zeitschrift fur Hautkrankheiten</i> . 1984;59(23):1579-80.	7
988	Frank DH, Wachtel T, Frank HA, Sanders R. Comparison of biobrane, porcine, and human allograft as biologic dressings for burn wounds. <i>Journal of Burn Care and Rehabilitation</i> . 1983;4(3):186-90.	2
989	Petersen SR, Umphred E, Warden GD. The incidence of bacteremia following burn wound excision. <i>Journal of Trauma</i> . 1982;22(4):274-9.	2
990	O'Connor NE, Mulliken JB, Banks-Schlegel S. Grafting of burns with cultured epithelium prepared from autologous epidermal cells. <i>Lancet</i> . 1981;1(8211):75-8.	6
991	Grant CG, Warren R, Salter E. Expansion of lapine, porcine and human skin in vitro for grafting on to extensive burns. <i>Burns</i> . 1980;7(1):57-69.	6
992	Prunieras M, Regnier M, Schlotterer M. New method to culture human epidermal cells on allogenic or xenogenic dermis: Preparation of recombined grafts. [French]. <i>Annales de Chirurgie Plastique</i> . 1979;24(4):357-62.	7
993	Tavis MJ, Thornton J, Danet R, Bartlett RH. Current status of skin substitutes. <i>Surgical Clinics of North America</i> . 1978;58(6):1233-48.	6
994	Silvetti AN. A new approach to skin autografts: comparison of healing of micro square and pulverized skin grafts in rabbits. <i>Ircs MedSciConnTiss</i> . 1975;3(2):96-7.	4
995	Nct. Application of Cultured Autologous Keratinocytes for Burn Wound Healing. <a href="https://clinicaltrials.gov/show/NCT00832156">https://clinicaltrials.gov/show/NCT00832156</a> . 2009.	6
996	Nct. Autologous Keratinocyte Suspension Versus Adipose-Derived Stem Cell-Keratinocyte Suspension for Post-Burn Raw Area. <a href="https://clinicaltrials.gov/show/NCT03686449">https://clinicaltrials.gov/show/NCT03686449</a> . 2018.	6
997	Nct. Non-cultured Autologous Keratinocyte Suspension Versus Traditional Split Skin Graft for Burn Wounds Treatment. <a href="https://clinicaltrials.gov/show/NCT03675568">https://clinicaltrials.gov/show/NCT03675568</a> . 2018.	6
998	Nct. Comparative Assessment of Adjuvant Effect of Cultured Epidermal Autografts Versus Skin Allografts on Wound Healing of Burns Treated With Widely Expanded Skin Autograft Using Meek Micrografting Technique MEEKADEAU. <a href="https://clinicaltrials.gov/show/NCT01330407">https://clinicaltrials.gov/show/NCT01330407</a> . 2011.	6
999	Nct. Controlled and Double Blind Comparison of a Traditional Dressing Versus a Biologic Dressing Composed of Fetal Fibroblasts and Keratinocytes in Association With a Collagen Matrix on Skin Donor Sites. <a href="https://clinicaltrials.gov/show/NCT03334656">https://clinicaltrials.gov/show/NCT03334656</a> . 2017.	6
1000	Euctr NL. Application of cultured autologous keratinocytes in combination with a meshed split skin autograft for burn wound healing - Cultured keratinocytes for burn wound treatment. <a href="http://www.who.int/trialsearch/Trial2.aspx?TrialID=EUCTR2007-004296-19-NL">http://www.who.int/trialsearch/Trial2.aspx?TrialID=EUCTR2007-004296-19-NL</a> . 2007.	6
1001	Isrctn. Cultured keratinocytes in burn wound care. <a href="http://www.who.int/trialsearch/Trial2.aspx?TrialID=ISRCTN63305738">http://www.who.int/trialsearch/Trial2.aspx?TrialID=ISRCTN63305738</a> . 2012.	6
1002	Nct. A Randomized, Controlled Trial of Autologous Platelet Gel Treatment in Diabetic Foot Ulcers. <a href="https://clinicaltrials.gov/show/NCT00338702">https://clinicaltrials.gov/show/NCT00338702</a> . 2006.	6

연번	서지정보	배제 사유
1003	Beele H, Naeyaert JM, Kettler M, Monstrey S, Kint A. Treatment of burns with autologous meshed split thickness skin covered by allogenic cultured keratinocytes or by cadaver skin: a clinical and histological study. 4th annual meeting of the european tissue repair society; 1994, 25-28 august; oxford, england. 1994:195.	6
1004	Ctri. A study to Compare the use of pigment producing[Melanin] Cell Suspension[taken from the skin] With Skin Grafting On Adequate Re-pigmentation Of white patches seen post burns. <a href="http://www.who.int/trialssearch/Trial2.aspx?TrialID=CTRI/2019/04/018804">http://www.who.int/trialssearch/Trial2.aspx?TrialID=CTRI/2019/04/018804</a> . 2019.	6
1005	ChiCtr. The effect and mechanism of negative pressure wound therapy on the survival rate and scar formation of modified Meek skin graft: a prospective randomized controlled study. <a href="http://www.who.int/trialssearch/Trial2.aspx?TrialID=ChiCTR2000034266">http://www.who.int/trialssearch/Trial2.aspx?TrialID=ChiCTR2000034266</a> . 2020.	6
1006	Irct138808062587N. Assessing Carbon Selenium Complex healing effect on partial thickness burns. <a href="http://www.who.int/trialssearch/Trial2.aspx?TrialID=IRCT138808062587N2">http://www.who.int/trialssearch/Trial2.aspx?TrialID=IRCT138808062587N2</a> . 2010.	6
1007	Jprn U. Using of a combination of meshed dermis graft and cultured epithelial autograft for massive burned patients. <a href="http://www.who.int/trialssearch/Trial2.aspx?TrialID=JPRN-UMIN000031258">http://www.who.int/trialssearch/Trial2.aspx?TrialID=JPRN-UMIN000031258</a> . 2018.	6

## 2. 국내 DB

연번	서지정보	배제 사유
1	임다혜, 배슬기, 오태호. Anti-inflammatory Effect of Shea Butter Extracts in Canine Keratinocytes. 한국임상수의학회지. 2021;38(1):27-31.	2
2	Rattanaprichavej P, Laoruengthana A, Galassi M, Weerakul S, Rasamimongkol S. Contamination Rate of Burnt Necrotic Tissue after Electrocoagulation in Total Knee Arthroplasty. Clinics in Orthopedic Surgery. 2020;12(1):43-8.	2
3	박지현, 오준상, 송지선, 김자영, 성기호. Bisfusarium Delphinoides, an Emerging Opportunistic Pathogen in a Burn Patient with Diabetes Mellitus. Mycobiology. 2019;47(3):340-5.	2
4	모영웅, 이동락, 신혜경, 정규용. Deep Second Degree Chemical Burn due to Topical Application of Clematis terniflora: A Case Report. Journal of Wound Management and Research. 2019;15(1):39-42.	2
5	주홍실, 이승제, 성건용. Early Debridement and Cultured Allogenic Keratinocyte Dressing Prevent Hypertrophic Scarring in Infants with Deep Dermal Burns. Archives of Aesthetic Plastic Surgery. 2018;24(3):111-5.	2
6	Seo G-Y, 임용호, 고통수, 허정식, 현창림, 김영미, et al. TMF and glycerin act synergistically on keratinocytes and fibroblasts to promote wound healing and anti-scarring activity. Experimental & Molecular Medicine. 2017;49(3):12-.	2
7	박지환, 박서화, 강은규, 경규천, 안효동, 안소연. 당뇨병 환자에서 발생한 피부 털곰팡이증. The Ewha Medical Journal. 2016;39(1):10-3.	2
8	김준현, 최성훈, 김영준. Alcaligenes Xylosoxidans 감염에 합병된 피부 궤양 증례. 대한화상학회지. 2015;18(1):35-7.	2
9	김대현, 김경식, 최준, 김승홍. 화상치료에서 동종유래배양세포이식과 젖산 용합체 합성상 치료제 적용의 비교. 대한화상학회지. 2014;17(2):73-80.	2
10	송병한, 윤종현, 박은주, 권인호, 김광호, 김광중. 편평 태선으로 오인한 만성 피부점막칸디다증. 대한의진균학회지. 2013;18(1):11-5.	2
11	정성필, 황태식, 송근정, 신명하, 이성준. 돼지의 화상 및 피부결손 모델에서 상처 치유과정의 비교. 대한화상학회지. 2013;16(1):58-63.	2



연번	서지정보	배제 사유
12	이방석, 임수아, 윤희일. 조기 박피술과 동종유래 표피세포를 이용한 깊은 2도 화상 치료의 효용성. 대한화상학회지. 2013;16(1):5-11.	2
13	김은지, 문구현. 동종유래각질세포(Cultured Allogenic Keratinocyte)의 다양한 임상적 적용. 대한창상학회지. 2013;9(2):122-5.	2
14	허시현, 한규범, 이영준, 김지현, 윤광호, 한만덕, et al. Recombinant human granulocyte macrophage colony-stimulating factor (rhGM-CSF) could accelerate burn wound healing in hamster skin. Animal Cells and Systems. 2012;16(3):207-14.	2
15	유경한, 이동락, 윤희일. 거자에 의해 발생한 화학적 화상의 증례보고. 대한화상학회지. 2012;15(2):134-6.	2
16	은석찬. 공여항원에 감작된 수지상세포를 이용한 피부 동종이식. 대한화상학회지. 2012;15(2):127-30.	2
17	정효선, 윤천재, 이정숙. 열 손상에서의 디펜신의 발현. 대한화상학회지. 2012;15(1):39-44.	2
18	장영호, 손용훈, 김상규, 박준모, 이미영, 김진모. 중화상 환자에서 중심정맥 카테터 감염: 쇄골하정맥과 대퇴정맥의 비교. 대한중환자의학회지. 2011;26(4):245-9.	2
19	최우영, 김규보, 양정열. 2도 화상 환자의 치료에서 동종 유래 배양피부(칼로덤)의 유용성. 대한화상학회지. 2011;14(2):111-7.	2
20	주지훈, 조현환, 이경희. 자초 뿌리의 함유성분 및 아토피피부염 관련 연구현황. 생약학회지. 2010;41(2):73-88.	2
21	장영호, 손용훈, 김상규, 박준모, 이미영. 중증 화상 환자에서 카테터 균체형성 감소를 위한 항균 중심정맥 카테터의 효과. 대한화상학회지. 2010;13(2):111-5.	2
22	허시현, 김지현, 윤광호, 이영준, 한규범, 김완중. 햄스터의 심재성 2도 화상치유에 미치는 rrhGM-CSF의 효과. 대한창상학회지. 2010;6(2):69-76.	2
23	여현정, 김준형, 정영진, 손대구, 한기환. 부분층 화상에 적용한 건조소양막과 동종배양표피 세포의 치료효과 비교. 대한성형외과학회지. 2009;36(4):385-92.	2
24	김재현, 탁경석, 서인석, 정찬민. 마늘에 의해 발생한 화학적 화상의 증례보고. 대한화상학회지. 2009;12(1):76-8.	2
25	권준성, 조용석, 김도현, 임해준, 허준, 전옥, et al. 중증 화상환자에서 칸디다혈증의 발생률 및 연관된 사망률. 대한화상학회지. 2009;12(1):64-7.	2
26	정철수, 최동휘. 자기 유래 배양피부(시트형태와 현탁액형태). 대한화상학회지. 2009;12(1):12-5.	6
27	허상택, 최영실, 박미연, 백경란, 송재훈, 오원섭, et al. Q열로 진단된 불명열 3예. 대한내과학회지. 2008;74(1):100-5.	2
28	윤신혁, 송철홍, 정재민, 심정수, 박대한. 화상치료에 있어서 동종유래표피세포의 유용성. 대한성형외과학회지. 2008;35(4):413-8.	2
29	서영권, 송계용, 박정극. 손상된 피부 재건을 위한 바이오인공피부의 개발 동향. 한국생물공학회지. 2008;23(1):8-17.	2
30	강교신, 하경임, 서무규, 이종임, 고우태. 와상백선의 양상을 보인 Trichophyton verrucosum 에 의한 몸백선 1예. 대한의진균학회지. 2008;13(4):181-6.	2
31	송종훈, 김도현, 허준, 전옥, 김종현. 동종피부를 이용한 광범위 화상의 치료. 대한외과학회지. 2007;72(1):11-7.	8
32	하태순, 조용석, 김도현, 허준, 전옥, 김종현, et al. 중증화상환자에서 액체부유 형태 자기유래배양피부의 임상적 유용성. 대한화상학회지. 2007;10(2):112-7.	6
33	최중호, 고장후, 서동국, 이종욱, 오석준, 장영철, et al. 동종 배양표피를 이용한 부분층 2도 화상의 치료. 대한성형외과학회지. 2006;33(5):587-91.	2
34	전옥. 광범위화상에서 인체유래 무세포성 진피대체물(AlloDerm)의 임상적 유용성. 대한창상학회지. 2006;2(2):87-92.	2
35	Cho Lee A-R. Enhancing Dermal Matrix Regeneration and Biomechanical Properties of Degree-Burn Wounds by EGF-Impregnated Collagen Sponge Dressing. Archives of Pharmacal Research. 2005;28(11):1311-6.	2
36	정국현/Jung K김허전김장전KDHHJCWKJHJYCSK. 동종피부(사체피부)이식과 자기유래 배양피부를 이용한 광범위 화상의 치료. 대한화상학회지. 2005;8(2):171-8.	6

연번	서지정보	배제 사유
37	전옥. 동종피부를 이용한 광범위 화상의 치료. 대한창상학회지. 2005;2(1):8-13.	2
38	김도완/Kim D봉정이박이BJGJJHLYSPJHLBK. 화상 창상 감염 : 위험 인자 및 예후에 미치는 영향. 침례병원학술지. 2002;18(0):26-37.	2
39	조애리. 설파디아진은의 피부세포 증식 및 화상모델에 있어서의 상처치유과정에 미치는 영향. 약학논문지-덕성여자대학교. 2002;13(1):135-9.	2
40	김도완, 봉진구, 정재한, 이윤식, 박진현, 이병철. 화상 창상 감염 위험 인자 및 예후에 미치는 영향. 대한외과학회지. 2001;61(2):195-202.	2
41	정재호, 이성길, 이승헌, 박명철. 화상 환자의 피부 이식편 괴사 창상에서 동정된 미생물. 대한수부재건외과학회지. 2001;6(2):97-101.	2
42	김영걸/Kim Y박PKK. 자외선 B조사에 의한 멜라닌세포의 아포프토시스. 대한피부연구학회지. 1999;6(2):92-6.	2
43	조광현/Cho K정이CJHLDK. 구강점막 상피세포의 삼차원적 배양. 대한피부연구학회지. 1999;6(2):65-70.	2
44	송인환/In Hwan Song AUBNDK. 시험관내 인공피부 제작. 체질인류학회지. 1998;11(2):247-60.	2
45	Clinical Experience of Gentian Violet Dressing for Local Treatment of Wound Infection with MRSA. 병원감염관리. 1998;3(1):23-31.	2
46	이지범, 정민영, 류경연, 기우천. 당뇨병 환자의 화상 창상에서 발생한 원발성 피부 아스페르길루스 1예. 대한피부과학회지. 1996;34(2):350-8.	2
47	이두익, 엄재광, 나영선, 박대희. Atelocollagen을 이용한 인공진피 제작에 관한 연구. 대한미용성형외과학회지. 1995;1(1):92-104.	2
48	문수만, 김태혁, 서기식. 화상환자 274예에 대한 임상적 고찰. 대한외과학회지. 1994;46(2):163-73.	2
49	KMbase 이심서강. 화염병화상의 특성. 대한외과학회지. 1993;45(6):1008-16.	2
50	김종혁, 이성렬, 김수남. Amorolfine 크림 0.25%를 이용한 피부사상균증 환자의 치료효과에 대한 연구. 감염. 1993;25(4):375-81.	2
51	박재경, 박재경, 박재경. 수축된 콜라겐 격자와 배양된 각질형성세포를 이용한 피부 대용물질의 제조에 관한 연구. 의공학회지. 1993;14(1):51-62.	2
52	KMbase 박이이유손김김이. Q Fever 1예 : Pancytopenia, Hepatitis, Myocarditis가 동반된 1예. 감염. 1992;24(1):45-54.	2
53	김동철, 이영호. 인체 상피세포의 시험관내 증식에 관한 연구. 대한성형외과학회지. 1991;18(3):390-406.	2
54	최원진, 정재현, 윤대원. 화상환자에 대한 임상적 고찰. 대한구급의학회지. 1991;6(1):27-37.	2
55	최삼섭, 朴贊, 洪元植. 溫病學 辯證論治의 發展에 關한 研究. 경희한의대논문집. 1987;10(0):415-66.	2
56	이경렬, 장일성, 최은숙. 화상환자에 대한 임상적 연구. 대한외과학회지. 1984;27(4):430-8.	2
57	주영화, 홍기웅. 고압전류화상 환자 46예에 대한 임상적 고찰. 대한외과학회지. 1984;26(5):673-82.	2
58	한경자/Kyung Ja Han AU정심이ISCSISCMMLK. Fusarium sp.에 의한 체부백선 1예. 대한병리학회지. 1983;17(4):442-6.	2
59	정해광, 이택호, 이종원, 이원영, 김주덕. 인체 상피세포의 시험관내 배양에 관한 연구. 대한성형외과학회지. 1983;10(3):269-78.	2
60	최용만. 火傷患者 282例에 關한 臨床的 考察. 한국생활과학연구원논문총. 1978;21(0):179-94.	2
61	김윤정, 최성희, 김선구. 임상검사물에서 분리된 혐기성세균 균종과 그 환자들의 임상상태. 대한병리학회지. 1978;12(4):427-32.	2
62	김년태, 朴容喆, 尹忠. 火傷患者 246例에 對한 臨床的 考察. 대한외과학회지. 1977;19(8):53-61.	2
63	玄鍾鳴/Hyun JM李朴金LNHPYJKJTK. 數種 皮膚疾患에 對한 Frason^®의 臨床的 經驗. 대한피부과학회지. 1977;15(2):191-5.	2
64	이열희, 서창환, 변기정, 김효순, 이상일. 안면화상으로 인한 골수염의 치험예. 대한치과의사협회지. 1977;15(2):121-4.	2

연번	서지정보	배제 사유
65	李永吉, 姜振聲. 羊膜을 이용한 火傷의 生化學的 處置. 대한성형외과학회지. 1977;4(2):43-54.	2
66	백근수, 양문석, 이방제. 화상환자의 피부이식에 대한 임상적 고찰. 대한외과학회지. 1976;18(3):57-61.	2
67	강중신. 0.5% Silver Nitrate Cream을 이용한 火傷의 處置. 대한외과학회지. 1974;16(12):17-20.	2
68	朱權源/Choo KW李李LCHLMSK. 尿道炎에 對한 Minocycline(Minocin)의 臨床效果. 최신의학. 1972;15(6):75-9.	2
69	安興仁, 愼鏞燦, 李華永. 電氣火傷患者 144例에 對한 臨床的 觀察. 대한외과학회지. 1972;14(2):25-37.	2
70	배수동. 화상의 치료. 한국의과학. 1972;4(7):438-43.	2
71	박영집. 지수의 간접계산법. 한일병원임상잡지. 1971;2(1):137-47.	2
72	이상훈, 이동훈, 장중동, 전병혁. 화상 후 발생한 심부 화농성 관절염. 대한정형외과학회지. 2005;40(7):992-1000.	2
73	이지연, 김성범, 박대원, 신성희, 김용현, 윤소영, et al. 백혈병환자에서 Liposomal Amphotericin B로 치료된 파종성 Fusarium 감염 1예. 감염과화학요법. 2003;35(5):321-4.	2
74	홍성표. 인공 피부의 개발과 이용 - 개발 상태와 연구 방향. 대한피부연구학회지. 1999;6(2):71-6.	2
75	박성주/Sung Ju Park AU민이김이강HGMHGLJMKESLHJKK. 비후성 반흔/켈로이드에 대한 냉동 외과술의 효과. 대한피부과학회지. 1998;36(5):765-71.	2
76	최성원/Choi SDBK. 알로에 베라 성분의 상피세포 증식 촉진효과. 대한피부연구학회지. 1998;5(2):179-92.	2
77	KMbase 박오주박최. 배양된 상피세포를 이용한 광범위 화상의 치료와 피부재건에 관한 연구. 대한외과학회지. 1989;37(2):135-48.	5
78	Jie F, Tong-da L, Meng-Yan Y, Yong-Heng W, Fu-Yuan C, Qingqi Y, et al. Keratin/PEO/hydroxyapatite Nanofiber Membrane with Improved Mechanical Property for Potential Burn Dressing Application. Fibers and polymers. 2020;21(2):366-75.	4
79	서영권, 박정극, 송계용, 서성준, 김영진. 포스터발표3 : 생물화공 ; P-417 : Manufacturing of collagen membrane for bioartificial skin. 화학공학의이론과응용. 2007;10(1):468-.	2
80	In vitro human 3-D culture model for bioscreening system. 한국동물실험대체법학회 학술대회집. 2008;2008(1):157-73.	2
81	김권천. 화상환자 675예에 대한 임상적 고찰. 最新醫學. 1994;37(3):19-28.	2
82	윤재천, 강성래, 이상호, 현태일, 노상현. 화상 환자에 대한 임상적 고찰. 大韓外傷學會誌. 1991;4(2):78-85.	2
83	서영권, 안재일, 이두훈, 권순용, 정도현, 박용순, et al. 각질형성세포가 배양된 인체 상피제 거양막의 상처치유효과. 조직공학과 재생의학. 2004;1(2):178-83.	2
84	이주영. <호살랑곳>과 <호탈곳>의 주술과 연행주체의 세계관 비교. 한국무속학. 2010;0(21):311-38.	2
85	황춘섭, 정현주. 마오리族 傳統 服飾과 文身 考察. 服飾文化研究. 1995;3(2):241-60.	2
86	P317 : Cutaneous Infection by serratia plymuthica in Social Community. 대한피부과학회 학술발표대회집. 2016;68(1):413-4.	2
87	하동주, 이무형. 백반증 환자의 표피이식 부위에 발생한 표재성 칸디다 감염. 대한의진균학회지. 2000;5(4):179-82.	2
88	P293 : A case of cutaneous mucormycosis after systemic steroid therapy. 대한피부과학회 학술발표대회집. 2015;67(1):408-.	2
89	Lee HT, Jang HS, Kim CG, Kwon KS, Chung TA. Clinical Efficacy and Tolerability of Terbinafine 1 % Cream in Patients with Pityriasis Versicolor. Annals of Dermatology. 1996;8(1):169-76.	2
90	손동수. 소아의 치조골 돌출도 및 전치상경사도에 관한 두부방사선계측학적 연구. 대한치과의사학회지. 2001;20(12):1049-52.	2
91	이부영. '재생' 의 상징적 의미. 心性研究. 1997;12(2):89-114.	2

연번	서지정보	배제 사유
92	서현호, 방난석, 권영대, 하지운, 김창식. 매치실린 저항성 포도당 구균 감염창에 대한 Gentian Violet Dressing. Arch. 1997;24(6):1334-42.	2
93	강희경, 장두희, 구한모. 농업인의 농약사용 실태에 관한 조사 연구. 資源科學研究論文集. 1999;7(-):1-18.	2
94	李永澤. 近世朝鮮의 法醫學的裁判과 無冤錄에 關한 研究. 서울대학교 論文集. 1956;4(-):177-279.	2
95	Yong Jin P, Jisu J, Jae Seok K, Dong Soo C, Goang-Won C, Jin Seong P, et al. Fabrication of Antimicrobial Wound Dressings Using Silver-Citrate Nanorods and Analysis of Their Wound-Healing Efficacy. 조선자연과학논문집. 2019;12(2):47-57.	6
96	박덕미. 고대 한국의 미용문화 연구. 韓服文化. 2005;8(1):43-54.	2
97	P113: A case of pemphigus vulgaris localized to the oral mucosa and esophagus. 대한피부과학회 학술발표대회집. 2018;70(1):352-3.	2
98	P252: Cutaneous mucormycosis arising in an old patient with pityriasis lichenoides et varioliformis acuta (PLEVA). 대한피부과학회 학술발표대회집. 2018;70(1):405-6.	2
99	박호철, 오수명, 주흥재. 234. 배양된 상피세포를 이용한 광범위화상의 치료와 피부재건에 관한 연구. 대한외과학회 학술대회 초록집. 1988;1988(11):110-1.	2
100	이영철, 조홍래, 김동건, 김성, 최원진, 이규만. 10. 광범위 화상환자의 피부 재건을 위한 in Vitro 상피세포 배양에 관한 연구. 대한외과학회 학술대회 초록집. 1992;1992(5):68-.	2
101	김종민, 이규만, 전진곤, 민형근. 화상환자의 세균 배양 양상과 항생제 감수성에 관한 연구. 大韓皮膚科學會誌. 2001;39(12):1391-6.	2
102	김영근, 은희철, 황규왕, 표형찬, 박영립. 수증 국소항균제의 배양 인체 각질형성세포 및 섬유아세포에 대한 세포독성도의 비교연구. 大韓皮膚科學會誌. 1995;33(5):895-906.	2
103	김수영, 김민정, 정호정, 이유나, 이양원, 최용범, et al. 정상 면역인에서 발생한 Stenotrophomonas maltophilia에 의한 국소 피부감염증. 大韓皮膚科學會誌. 2015;53(1):80-2.	2
104	전진곤, 민형근, 김종민, 이규만. 제53차 춘계학술대회 : 일반 포스터 ; 화상후 세균배양 양상과 항생제 감수성에 관한 연구. 초록집. 2001;39(10):113-.	6
105	張錫洛, 金弘泰, 金恒在, 金能守. 有熱患者의 血液培養에서 分離된 病原菌에 關하여. 慶北醫大誌. 1981;22(2):384-90.	2
106	구자성, 차재령, 오세행, 공명선. 셀포베타인 키토산의 실크 블렌드 필름의 제조 및 그들의 성질. 폴리머. 2014;38(1):54-61.	2
107	강상진. 3D 피부세포 배양계를 이용한 피부광노화 연구. 대한화장품학회지(J Soc Cosmet Scientists Korea). 1999;25(2):59-75.	2
108	김진화, 박성민, 심관섭, 이범천, 표형배. 약용식물추출물의 광보호 효과와 항염증 효과 연구. 대한화장품학회지(J Soc Cosmet Scientists Korea). 2004;30(2):227-33.	2
109	강정대, 김소진, 유동오, 박석돈. 증례 : 화상 후 시행한 피부 이식 부위에 발생한 원발성 피부 아스페르길루스증 1예. 대한피부과학회지. 2002;40(12):1547-51.	2
110	이지범, 이승철, 원영호, 전인기. 증례 : 당뇨병 환자의 화상 창상에서 발생한 원발성 피부 아스페르길루스증 1예. 대한피부과학회지. 1996;34(2):350-4.	2
111	조애리. 원보 : 설파디아진은의 피부세포 증식 및 화상모델에 있어서의 상처치유과정에 미치는 영향. 약제학회지. 2002;32(2):113-7.	2
112	金赫. 甲骨卜辭所見의 商代求雨活動. 중국학논총. 2020;70(0):1-26.	2
113	조봉근. 한국복음주의조직신학회 20년사 회고록. 조직신학연구. 2017;27(0):46-73.	2
114	유영삼, 허건. 구강 수술에 사용가능한 LED 헤드라이트의 안전성 및 실용성. 대한두경부종양학회지 (대한두경부종양학회). 2010;26(2):187-92.	2
115	Lee, Ae Ri C, Moon, Hee K. Effect of Topically Applied Silver Sulfadiazine on Fibroblast Cell Proliferation and Biomechanical Properties of the Wound. Archives of pharmacal research (대한약학회). 2003;26(10):855-60.	2
116	이현우, 홍승욱. 가미대황고(加味大黃膏)의 멜라닌세포 활성화제가 자외선 조사로 인한 피부 손상 완화에 미치는 영향. 한방안이비인후피부과학회지 (대한한방안이비인후피부과학회). 2007;20(2):47-67.	2

연번	서지정보	배제 사유
117	노성원, 최원진, 변두진. 화학 화상. 대한외과학회지. 1995;48(4):461-8.	2
118	송경원, 홍기웅, 이용창, 박용철. 화상환자 443예에 대한 임상적 고찰. 대한외과학회지. 1980;22(8):35-44.	2
119	정재현, 최원진, 윤대원. 원저 : 화상환자에 대한 임상적 고찰. 대한중환자의학회지. 1991;6(1):27-37.	2
120	정재현, 최원진, 윤대원. 화상환자에 대한 임상적 고찰. 대한외과학회지. 1991;40(6):771-81.	8
121	박동수, 이영철, 박천규. 화상환자 952예에 대한 임상적 고찰. 대한외과학회지. 1986;31(2):225-35.	2
122	이선호, 이원기. 전기염상환자 171 예에 대한 관찰. 대한정형외과학회지. 1973;8(1):19-28.	2
123	임태진, 장수일, 황선희. 화상의 각종 국소요법에 관한 임상적 관찰. 대한외과학회지. 1978;20(1):37-44.	2
124	조형석, 김승철, 김익성. 급성 충수염에 대한 임상적 비교 고찰. 대한외과학회지. 1983;25(1):59-66.	2
125	임종명. 논문 : 여순사건의 재현과 공간(空間). 한국사학보. 2005;19(0):151-85.	2
126	전진곤, 민형근, 김종민, 이규만. 원저 : 화상환자의 세균 배양 양상과 항생제 감수성에 관한 연구. 대한피부과학회지. 2001;39(12):1391-6.	2
127	표형찬, 김영근, 황규왕, 박영립, 은희철. 원저 : 수종 국소항균제의 배양 인체 각질형성세포 및 섬유아세포에 대한 세포독성도의 비교연구. 대한피부과학회지. 1995;33(5):895-906.	2
128	조용훈. 다문화 사회에서 기독교의 윤리적 과제. 기독교사회윤리. 2011;22(0):303-30.	2
129	김현실, 김남희, 김진, 차인호. 배양된 구강점막 각화상피세포가 누드마우스 피부 창상 치유에 미치는 효과. 大韓口腔顎顔面外科學會誌 = Journal of the Korean Association of Oral and Maxillofacial Surgeons (대). 2004;30(4):308-15.	2
130	이종헌. 구강 상피성 종양연구에 필요한 배양된 정상인 구강 및 피부 각화상피세포에서 분화 단백질 표지자. 대한치과의사협회지 (대한치과의사협회). 2002;40(7):500-2.	2
131	Shin I-S, Kwon O-H, Kim S-N, Hong C-M, Lee K-H, Oh H-J, et al. In vitro culture of skin cells on a crosslinked gelatin based scaffold for artificial skin. 대한약학회 2003년도 Proceedings of the Convention of the Pharmaceutical Society of Korea Vol2-2. 2003;2003(10):100.2-2.	2
132	Song HJ, Lim HY, Chun W, Choi KC, Lee T-y, Sung JH, et al. Development of 3D skin-equivalent in a pump-less microfluidic chip. Journal of industrial and engineering chemistry : JIEC. 2018;60:355-9.	2
133	곽소정, 김상수, 전현욱, 이영무, 박문향, 송강원, et al. 각질세포와 섬유모세포가 배양된 고분자 미립구를 이용한 피부재생. 한국생물공학회 2003년도 생물공학의 동향(XII). 2003;2003(4):92-4.	2
134	Gwak S-J, Kim S-S, Choi C-Y, Kim B-S. Skin regeneration by spray of fibrin gel containing keratinocytes and epidermal growth factor. 한국생물공학회 2003년도 생물공학의 동향(XIII). 2003;2003(10):356-8.	2