

별첨 3

배제문헌

문현배제사유

1. 동물연구 또는 전임상시험
2. 원저가 아닌 연구(종설, letter, comment 등)
3. 중재검사를 수행하지 않은 연구
4. 적절한 의료결과를 하나 이상 보고하지 않은 연구
5. 논문 철회된 연구
6. 증례보고
7. 사전에 정의된 대상자를 대상으로 수행되지 않은 연구
8. 원문 확보가 불가한 경우
9. 중복출판된 연구

연번	서지정보	배제 사유
1	Aboelatta Y, Abdelsalam A. Volume overload of fluid resuscitation in acutely burned patients using transpulmonary thermodilution technique. <i>Journal of Burn Care & Research.</i> 2013;34(3):349–54.	3
2	Adler C, Reuter H, Seck C, Hellmich M, Zobel C. Fluid therapy and acute kidney injury in cardiogenic shock after cardiac arrest. <i>Resuscitation.</i> 2013;84(2):194–9.	3
3	Alavi M, Hadipourzadeh F, Azarfarin R, Babaee T, Ziayefard M. Effects of advanced hemodynamic monitoring on the postoperative intubation time in patients with moderate-to-severe left ventricular dysfunction undergoing cardiac surgery. <i>Iranian Heart Journal.</i> 2020;21(4):25–32.	4
4	Ali A, Abdullah T, Orhan-Sungur M, Orhun G, Aygun E, Aygun E, et al. Transpulmonary thermodilution monitoring-guided hemodynamic management improves cognitive function in patients with aneurysmal subarachnoid hemorrhage: a prospective cohort comparison. <i>Acta Neurochirurgica.</i> 2019;161(7):1317–24.	3
5	Alonso-Inigo JM, Escriba FJ, Carrasco JI, Fas MJ, Argente P, Galvis JM, et al. Measuring cardiac output in children undergoing cardiac catheterization: comparison between the Fick method and PRAM (pressure recording analytical method). <i>Paediatric Anaesthesia.</i> 2016;26(11):1097–105.	3
6	Ameloot K, Van De Vijver K, Broch O, Van Regenmortel N, De Laet I, Schoonheydt K, et al. Nexfin noninvasive continuous hemodynamic monitoring: Validation against continuous pulse contour and intermittent transpulmonary thermodilution derived cardiac output in critically ill patients. <i>The Scientific World Journal.</i> 2013;2013 (no pagination)(519080).	4
7	Bartha E, Davidson T, Berg HE, Kalman S. A 1-year perspective on goal-directed therapy in elderly with hip fracture: Secondary outcomes. <i>Acta Anaesthesiologica Scandinavica.</i> 2019;63(5):610–4.	3
8	Bartha E, Davidson T, Brodkorb TH, Carlsson P, Kalman S. Value of information: interim analysis of a randomized, controlled trial of goal-directed hemodynamic treatment for aged patients. <i>Trials.</i> 2013;14:205.	3

연번	서지정보	배제 사유
9	Baulig W, Bernhard EO, Bettex D, Schmidlin D, Schmid ER. Cardiac output measurement by pulse dye densitometry in cardiac surgery. <i>Anaesthesia</i> . 2005;60(10):968–73.	3
10	Baulig W, Schuett P, Goedje O, Schmid ER. Accuracy of a novel approach to measuring arterial thermodilution cardiac output during intra-aortic counterpulsation. <i>Journal of Clinical Monitoring & Computing</i> . 2007;21(3):147–53.	3
11	Bein B, Worthmann F, Tonner PH, Paris A, Steinfath M, Hedderich J, et al. Comparison of esophageal Doppler, pulse contour analysis, and real-time pulmonary artery thermodilution for the continuous measurement of cardiac output. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2004;18(2):185–9.	3
12	Belda FJ, Aguilar G, Teboul JL, Pestana D, Redondo FJ, Malbrain M, et al. Complications related to less-invasive haemodynamic monitoring. <i>British Journal of Anaesthesia</i> . 2011;106(4):482–6.	4
13	Bergstrom B, de la Cruz JS, Sally M, Louis S, Friedman M, Petersen F, et al. The Use of Stroke Volume Variation to Guide Donor Management Is Associated With Increased Organs Transplanted per Donor. <i>Progress in Transplantation</i> . 2017;27(2):200–6.	7
14	Boisson M, Pontier B, Kerforne T, Frasca D. Comparison of noninvasive and minimally invasive pulse contour analysis to measure stroke volume during major surgery: A prospective observational study. <i>European Journal of Anaesthesiology</i> . 2018;35(11):895–6.	4
15	Botsch A, Firstenberg MS. Comment on the Edwards FloTrac TM /Vigileo versus pulmonary artery catheter study: What is really going on with this patient? <i>International Journal of Critical Illness and Injury Science</i> . 2017;7(3):183–4.	2
16	Boyle M, Lawrence J, Belessis A, Murgo M, Shehabi Y. Comparison of dynamic measurements of pulse contour with pulsed heat continuous cardiac output in postoperative cardiac surgical patients. <i>Australian Critical Care</i> . 2007;20(1):27–32.	3
17	Bremer F, Schiele A, Tschaikowsky K. Cardiac output measurement by pulse dye densitometry: a comparison with the Fick's principle and thermodilution method. <i>Intensive Care Medicine</i> . 2002;28(4):399–405.	3
18	Breukers RM, Groeneveld AB, de Wilde RB, Jansen JR. Transpulmonary versus continuous thermodilution cardiac output after valvular and coronary artery surgery. <i>Interactive Cardiovascular & Thoracic Surgery</i> . 2009;9(1):4–8.	3
19	Broch O, Carstens A, Gruenewald M, Nischelsky E, Vellmer L, Bein B, et al. Non-invasive hemodynamic optimization in major abdominal surgery: a feasibility study. <i>Minerva anestesiologica</i> . 2016;82(11):1158–69.	3
20	Brock-Utne JG, Blake GT, Bosenberg AT, Gaffin SL, Humphrey D, Downing JW. An evaluation of the pulse-contour method of measuring cardiac output. <i>South African Medical Journal</i> Suid-Afrikaanse Tydskrif Vir Geneeskunde. 1984;66(12):451–3.	3
21	Buhre W, Weyland A, Kazmaier S, Hanekop GG, Baryalei MM, Sydow M, et al. Comparison of cardiac output assessed by pulse-contour analysis and thermodilution in patients undergoing minimally invasive direct coronary artery bypass grafting. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 1999;13(4):437–40.	4
22	Button D, Weibel L, Reuthebuch O, Genoni M, Zollinger A, Hofer CK. Clinical evaluation of the FloTrac/Vigileo system and two established continuous cardiac output monitoring devices in patients undergoing cardiac surgery. <i>British Journal of Anaesthesia</i> . 2007;99(3):329–36.	4
23	Cannesson M, Musard H, Desebbe O, Boucau C, Simon R, Henaine R, et al. The ability of stroke volume variations obtained with Vigileo/FloTrac system to monitor fluid responsiveness in mechanically ventilated patients. <i>Anesthesia & Analgesia</i> . 2009;108(2):513–7.	4
24	Chen ZH, Jin CD, Chen S, Chen XS, Wang ZE, Liu W, et al. The application of early goal directed therapy in patients during burn shock stage. <i>International Journal of Burns & Trauma</i> . 2017;7(3):27–33.	3
25	Chiao HY, Chou CY, Tzeng YS, Wang CH, Chen SG, Dai NT. Goal-Directed Fluid Resuscitation Protocol Based on Arterial Waveform Analysis of Major Burn Patients in a Mass Burn Casualty. <i>Annals of Plastic Surgery</i> . 2018;80(2S Suppl 1):S21–S5.	4
26	Compton F, Wittrock M, Schaefer JH, Zidek W, Tepel M, Scholze A. Noninvasive cardiac output determination using applanation tonometry-derived radial artery pulse contour	3

연번	서지정보	배제 사유
	analysis in critically ill patients. <i>Anesthesia & Analgesia</i> . 2008;106(1):171-4, table of contents.	
27	Cottis R, Magee N, Higgins DJ. Haemodynamic monitoring with pulse-induced contour cardiac output (PiCCO) in critical care. <i>Intensive & Critical Care Nursing</i> . 2003;19(5):301-7.	2
28	Csontos C, Foldi V, Fischer T, Bogar L. Arterial thermodilution in burn patients suggests a more rapid fluid administration during early resuscitation. <i>Acta Anaesthesiologica Scandinavica</i> . 2008;52(6):742-9.	3
29	de Wilde RB, Geerts BF, van den Berg PC, Jansen JR. A comparison of stroke volume variation measured by the LiDCOplus and FloTrac-Vigileo system. <i>Anaesthesia</i> . 2009;64(9):1004-9.	4
30	de Wilde RB, Schreuder JJ, van den Berg PC, Jansen JR. An evaluation of cardiac output by five arterial pulse contour techniques during cardiac surgery. <i>Anaesthesia</i> . 2007;62(8):760-8.	4
31	Della Rocca G, Costa GM, Coccia C, Pompei L, Di Marco P, Pietropaoli P. Preload index: pulmonary artery occlusion pressure versus intrathoracic blood volume monitoring during lung transplantation. <i>Anesthesia & Analgesia</i> . 2002;95(4):835-43, table of contents.	3
32	Della Rocca G, Costa MG, Coccia C, Pompei L, Di Marco P, Vilardi V, et al. Cardiac output monitoring: aortic transpulmonary thermodilution and pulse contour analysis agree with standard thermodilution methods in patients undergoing lung transplantation. <i>Canadian Journal of Anaesthesia</i> . 2003;50(7):707-11.	3
33	Della Rocca G, Costa MG, Coccia C, Pompei L, Pietropaoli P. Preload and haemodynamic assessment during liver transplantation: a comparison between the pulmonary artery catheter and transpulmonary indicator dilution techniques. <i>European Journal of Anaesthesiology</i> . 2002;19(12):868-75.	3
34	Della Rocca G, Costa MG, Pompei L, Coccia C, Pietropaoli P. Continuous and intermittent cardiac output measurement: pulmonary artery catheter versus aortic transpulmonary technique. <i>British Journal of Anaesthesia</i> . 2002;88(3):350-6.	3
35	Diaz S, Perez-Pena J, Sanz J, Olmedilla L, Garutti I, Barrio JM. Haemodynamic monitoring and liver function evaluation by pulsion cold system Z-201 (PCS) during orthotopic liver transplantation. <i>Clinical Transplantation</i> . 2003;17(1):47-55.	3
36	Donati A, Carsetti A, Tondi S, Scorcetta C, Domizi R, Damiani E, et al. Thermodilution vs pressure recording analytical method in hemodynamic stabilized patients. <i>Journal of Critical Care</i> . 2014;29(2):260-4.	4
37	Ebm C, Cecconi M, Sutton L, Rhodes A. A cost-effectiveness analysis of postoperative goal-directed therapy for high-risk surgical patients. <i>Critical Care Medicine</i> . 2014;42(5):1194-203.	3
38	Edanaga M, Hayashi H, Yamakage M. Usefulness of Flotrac/VigileoTM combined with transesophageal echocardiography under abdominal hysterectomy for a patient complicated with moyamoya disease and hypertrophic cardiomyopathy. <i>Anesthesia and Resuscitation</i> . 2013;49(4):101-3.	6
39	Felbinger TW, Reuter DA, Eltzschig HK, Bayerlein J, Goetz AE. Cardiac index measurements during rapid preload changes: a comparison of pulmonary artery thermodilution with arterial pulse contour analysis. <i>Journal of Clinical Anesthesia</i> . 2005;17(4):241-8.	4
40	Felbinger TW, Reuter DA, Eltzschig HK, Moerstedt K, Goedje O, Goetz AE. Comparison of pulmonary arterial thermodilution and arterial pulse contour analysis: evaluation of a new algorithm. <i>Journal of Clinical Anesthesia</i> . 2002;14(4):296-301.	3
41	Fischer MO, Diouf M, Wilde RBP, Dupont H, Hanouz JL, Lorne E. Evaluation of cardiac output by 5 arterial pulse contour techniques using trend interchangeability method. <i>Medicine</i> . 2016;95(25):e3530.	4
42	Fletcher AM, Andrews J, Frampton AE. Individualizing hemodynamic optimization during the management of circulatory collapse. <i>Expert Review of Cardiovascular Therapy</i> . 2012;10(10):1217-20.	4
43	Franchi F, Silvestri R, Cubattoli L, Taccone FS, Donadello K, Romano SM, et al. Comparison between an uncalibrated pulse contour method and thermodilution technique for cardiac output estimation in septic patients. <i>British Journal of Anaesthesia</i> . 2011;107(2):202-8.	3

연번	서지정보	배제 사유
44	Friescke S, Heinrich A, Abel P, Felix SB. Comparison of pulmonary artery and aortic transpulmonary thermodilution for monitoring of cardiac output in patients with severe heart failure: validation of a novel method. <i>Critical Care Medicine</i> . 2009;37(1):119–23.	3
45	Froghi F, Soggiu F, Ricciardi F, Vindrola-Padros C, Floros L, Martin D, et al. Ward based goal directed fluid therapy (GDFT) in acute pancreatitis (GAP) trial: a feasibility randomised controlled trial. <i>International journal of surgery (London, England)</i> . 2022;104:106737.	3
46	Führer H, Gunther A, Zinke J, Niesen WD. Optimizing cardiac out-put to increase cerebral penumbral perfusion in large middle cerebral artery ischemic lesion—OPTIMAL Study. <i>Frontiers in Neurology</i> . 2017;8(AUG) (no pagination)(402).	4
47	Garcia-Lorenzo B, Fernandez-Barcelo C, Maduell F, Sampietro-Colom L. Health Technology Assessment of a new water quality monitoring technology: Impact of automation, digitalization and remoteness in dialysis units. <i>PLoS ONE [Electronic Resource]</i> . 2021;16(2):e0247450.	3
48	Giomarelli P, Biagioli B, Scolletta S. Cardiac output monitoring by pressure recording analytical method in cardiac surgery. <i>European Journal of Cardio-Thoracic Surgery</i> . 2004;26(3):515–20.	3
49	Godje O, Hoke K, Lamm P, Schmitz C, Thiel C, Weinert M, et al. Continuous, less invasive, hemodynamic monitoring in intensive care after cardiac surgery. <i>Thoracic & Cardiovascular Surgeon</i> . 1998;46(4):242–9.	3
50	Godje O, Thiel C, Lamm P, Reichenspurner H, Schmitz C, Schutz A, et al. Less invasive, continuous hemodynamic monitoring during minimally invasive coronary surgery. <i>Annals of Thoracic Surgery</i> . 1999;68(4):1532–6.	4
51	Goedje O, Hoeke K, Lichtwarck-Aschoff M, Faltchauser A, Lamm P, Reichart B. Continuous cardiac output by femoral arterial thermodilution calibrated pulse contour analysis: comparison with pulmonary arterial thermodilution. <i>Critical Care Medicine</i> . 1999;27(11):2407–12.	3
52	Gratz I, Kraidin J, Jacobi AG, deCastro NG, Spagna P, Larjani GE. Continuous noninvasive cardiac output as estimated from the pulse contour curve. <i>Journal of Clinical Monitoring</i> . 1992;8(1):20–7.	3
53	Greiwe G, Luehsen K, Hapfelmeier A, Rogge D, Kubik M, Schulte-Uentrop L, et al. Cardiac output estimation by pulse wave analysis using the pressure recording analytical method and intermittent pulmonary artery thermodilution: A method comparison study after off-pump coronary artery bypass surgery. <i>European Journal of Anaesthesiology</i> . 2020;37(10):920–5.	3
54	Grigorov Tzenkov I, Arnal Velasco D, Perez Pena JM, Olmedilla Arnal L, Garutti Martinez I, Sanz Fernandez J. Cardiac output by femoral arterial thermodilution—calibrated pulse contour analysis during liver transplantation: comparison with pulmonary artery thermodilution. <i>Transplantation Proceedings</i> . 2003;35(5):1920–2.	3
55	Gruenewald M, Renner J, Meybohm P, Hocker J, Scholz J, Bein B. Reliability of continuous cardiac output measurement during intra-abdominal hypertension relies on repeated calibrations: An experimental animal study. <i>Critical Care</i> . 2008;12(5) (no pagination)(R132).	1
56	Grundmann CD, Wischermann JM, Fassbender P, Bischoff P, Frey UH. Hemodynamic monitoring with Hypotension Prediction Index versus arterial waveform analysis alone and incidence of perioperative hypotension. <i>Acta Anaesthesiologica Scandinavica</i> . 2021;65(10):1404–12.	4
57	Gust R, Gottschalk A, Bauer H, Bottiger BW, Bohrer H, Martin E. Cardiac output measurement by transpulmonary versus conventional thermodilution technique in intensive care patients after coronary artery bypass grafting. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 1998;12(5):519–22.	4
58	Halvorsen PS, Espinoza A, Lundblad R, Cvancarova M, Hol PK, Fosse E, et al. Agreement between PiCCO pulse-contour analysis, pulmonary artery thermodilution and transthoracic thermodilution during off-pump coronary artery by-pass surgery. <i>Acta Anaesthesiologica Scandinavica</i> . 2006;50(9):1050–7.	4
59	Halvorsen PS, Sokolov A, Cvancarova M, Hol PK, Lundblad R, Tonnessen TI. Continuous cardiac output during off-pump coronary artery bypass surgery: pulse-contour analyses vs pulmonary artery thermodilution. <i>British Journal of Anaesthesia</i> . 2007;99(4):484–92.	3

연번	서지정보	배제 사유
60	Hamilton TT, Huber LM, Jessen ME. PulseCO: a less-invasive method to monitor cardiac output from arterial pressure after cardiac surgery. <i>Annals of Thoracic Surgery</i> . 2002;74(4):S1408–12.	3
61	Heijne A, Krijtenburg P, Bremers A, Scheffer GJ, Malagon I, Slagt C. Four different methods of measuring cardiac index during cytoreductive surgery and hyperthermic intraperitoneal chemotherapy. <i>Korean J Anesthesiol</i> . 2021 Apr;74(2):120–133. doi: 10.4097/kja.20202. Epub 2020 Aug 21. PMID: 32819047; PMCID: PMC8024204.	4
62	Heise D, Faulstich M, Morer O, Brauer A, Quintel M. Influence of continuous renal replacement therapy on cardiac output measurement using thermodilution techniques. <i>Minerva Anestesiologica</i> . 2012;78(3):315–21.	3
63	Herold IH, Soliman Hamad MA, van Assen HC, Bouwman RA, Korsten HH, Mischi M. Pulmonary blood volume measured by contrast enhanced ultrasound: a comparison with transpulmonary thermodilution. <i>British Journal of Anaesthesia</i> . 2015;115(1):53–60.	4
64	Hilty MP, Franzen DP, Wyss C, Biaggi P, Maggiorini M. Validation of transpulmonary thermodilution variables in hemodynamically stable patients with heart diseases. <i>Annals of Intensive Care</i> . 2017;7(1):86.	3
65	Hofer CK, Furrer L, Matter-Ensner S, Maloigne M, Klaghofer R, Genoni M, et al. Volumetric preload measurement by thermodilution: a comparison with transoesophageal echocardiography. <i>British Journal of Anaesthesia</i> . 2005;94(6):748–55.	3
66	Holm C, Melcer B, Horbrand F, Henckel von Donnersmarck G, Mühlbauer W. Arterial thermodilution: an alternative to pulmonary artery catheter for cardiac output assessment in burn patients. <i>Burns</i> . 2001;27(2):161–6.	4
67	Huang CC, Chen NH, Li LF, Yang CT, Hsiao HF, Chen YH, et al. Effects of cardiac output levels on the measurement of transpulmonary thermodilution cardiac output in patients with acute respiratory distress syndrome. <i>The Journal of Trauma and Acute Care Surgery</i> . 2012;73(5):1236–41.	3
68	Huang CC, Kao KC, Fu JY, Hsieh MJ. Effects of extravascular lung water on the measurement of transpulmonary thermodilution cardiac output in acute respiratory distress syndrome patients. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2011;25(3):481–5.	3
69	Huang DT, Clermont G, Dremszov TT, Angus DC, Pro Cl. Implementation of early goal-directed therapy for severe sepsis and septic shock: A decision analysis. <i>Critical Care Medicine</i> . 2007;35(9):2090–100.	3
70	Huang J, Zhu P, Zhong F, Yu G, Ye B, Fang L. Clinical significance of pulse index contour continuous cardiac output monitoring in patients with constrictive pericarditis undergoing pericardectomy. <i>Interactive Cardiovascular & Thoracic Surgery</i> . 2020;31(3):364–8.	3
71	Huber W, Koenig J, Mair S, Schuster T, Saugel B, Eyer F, et al. Predictors of the accuracy of pulse-contour cardiac index and suggestion of a calibration-index: a prospective evaluation and validation study. <i>BMC Anesthesiology</i> . 2015;15:45.	4
72	Ishihara H, Okawa H, Tanabe K, Tsubo T, Sugo Y, Akiyama T, et al. A new non-invasive continuous cardiac output trend solely utilizing routine cardiovascular monitors. <i>Journal of Clinical Monitoring & Computing</i> . 2004;18(5–6):313–20.	3
73	Ishihara H, Sugo Y, Tsutsui M, Yamada T, Sato T, Akazawa T, et al. The ability of a new continuous cardiac output monitor to measure trends in cardiac output following implementation of a patient information calibration and an automated exclusion algorithm. <i>Journal of Clinical Monitoring and Computing</i> . 2012;26(6):465–71.	3
74	Isosu T, Obara S, Ohashi S, Hosono A, Nakano Y, Imaizumi T, et al. Examination of the usefulness of non-invasive stroke volume variation monitoring for adjusting fluid supplementation during laparoscopic adrenalectomy in patients with pheochromocytoma. <i>Fukushima Journal of Medical Science</i> . 2012;58(1):78–81.	4
75	Jammer I, Tuovila M, Ulvik A. Stroke volume variation to guide fluid therapy: is it suitable for high-risk surgical patients? A terminated randomized controlled trial. <i>Perioperative Medicine</i> . 2015;4:6.	3
76	Jiang Z, Chen J, Gao C, Tan M, Zhang W, Xie Y. Effects of PICCO in the guidance of goal-directed fluid therapy for gastrointestinal function after cytoreductive surgery for ovarian cancer. <i>American Journal Of Translational Research</i> . 2021;13(5):4852–9.	3
77	Kim DW, Lee WK, Kim JY, Kim JL, Park SJ, Tak SJ. The comparison of cardiac outputs	3

연번	서지정보	배제 사유
	between impedance cardiography and thermodilution technique. <i>J of KOSOMBE</i> . 9(2):165–9.	
78	Kirton OC, Calabrese RC, Staff I. Increasing use of less-invasive hemodynamic monitoring in 3 specialty surgical intensive care units: a 5-year experience at a tertiary medical center. <i>Journal of Intensive Care Medicine</i> . 2015;30(1):30–6.	4
79	Kitaguchi K, Gotohda N, Yamamoto H, Kato Y, Takahashi S, Konishi M, et al. Intraoperative circulatory management using the FloTrac TM system in laparoscopic liver resection. <i>Asian Journal of Endoscopic Surgery</i> . 2015;8(2):164–70.	4
80	Kovacs E, Gyarmathy VA, Pilecky D, Fekete-Gyor A, Szakal-Toth Z, Geller L, et al. An Interaction Effect Analysis of Thermodilution-Guided Hemodynamic Optimization, Patient Condition, and Mortality after Successful Cardiopulmonary Resuscitation. <i>International Journal of Environmental Research & Public Health [Electronic Resource]</i> . 2021;18(10):14.	3
81	Kraft R, Herndon DN, Branski LK, Finnerty CC, Leonard KR, Jeschke MG. Optimized fluid management improves outcomes of pediatric burn patients. <i>Journal of Surgical Research</i> . 2013;181(1):121–8.	3
82	Kratz T, Hinterobermaier J, Timmesfeld N, Kratz C, Wulf H, Steinfeldt T, et al. Pre-operative fluid bolus for improved haemodynamic stability during minor surgery: A prospectively randomized clinical trial. <i>Acta Anaesthesiologica Scandinavica</i> . 2018;62(9):1215–22.	4
83	Kroon M, Groeneveld AB, Smulders YM. Cardiac output measurement by pulse dye densitometry: comparison with pulmonary artery thermodilution in post-cardiac surgery patients. <i>Journal of Clinical Monitoring & Computing</i> . 2005;19(6):395–9.	3
84	Kuntscher MV, Blome-Eberwein S, Pelzer M, Erdmann D, Germann G. Transcardiopulmonary vs pulmonary arterial thermodilution methods for hemodynamic monitoring of burned patients. <i>Journal of Burn Care & Rehabilitation</i> . 2002;23(1):21–6.	3
85	Landais A, Morel M, Goldstein J, Loriau J, Fresnel A, Chevalier C, et al. Evaluation of financial burden following complications after major surgery in France: Potential return after perioperative goal-directed therapy. <i>Anaesthesia Critical Care & Pain Medicine</i> . 2017;36(3):151–5.	4
86	Lavi R, Cheng D. Pro: Continuous cardiac output and SvO_2 monitoring should be routine during off-pump coronary artery bypass graft surgery. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> . 2012;26(6):1131–5.	2
87	Lee HJ, Cho HS. Cardiac output measurement by thermodilution. <i>대한구급의학회지</i> . 1988;3(2):125–32.	3
88	Li T, Cai H, Pan H, Pu Q. Cardiac output measurement using a modified carbon dioxide Fick method: comparison analysis with pulmonary artery catheter method and pulse induced contour cardiac output method. <i>International journal of clinical and experimental medicine</i> . 2015;8(3):3530–7.	4
89	Li-Ping Q, Hong-Wei L, Chang-Ming H, Yong-Yi B, Ang L. Safety and efficacy of pulse-induced contour cardiac output monitoring in elderly patients with coronary artery disease and severe heart failure at coronary care units. <i>Frontiers in Cardiovascular Medicine</i> . 2022;9:910898.	3
90	Liu X, Ji W, Wang J, Pan T. Application strategy of PiCCO in septic shock patients. <i>Experimental & Therapeutic Medicine</i> . 2016;11(4):1335–9.	4
91	Lu X, Zhai H, Dong Y, Su F, Xie Y, Wang Y, et al. Therapeutic Effect and Prognosis of PiCCO in the Treatment of Myocardial Injury Complicated with Septic Shock. <i>Computational & Mathematical Methods in Medicine</i> . 2022;2910849.	3
92	Luo Y, Zhan S, Zhu L, Xiong M, Liu G, Wang C. The role of pulse indicator continuous cardiac output (PiCCO) and critical care ultrasound in volume status assessment during fluid resuscitation for and prognosis of septic shock patients. <i>Pakistan Journal of Medical Sciences</i> . 2023;39(1):214–8.	3
93	Maass SW, Roekaerts PM, Lance MD. Cardiac output measurement by bioimpedance and noninvasive pulse contour analysis compared with the continuous pulmonary artery thermodilution technique. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2014;28(3):534–9.	3
94	Manecke GR, Jr. Cardiac output from the arterial catheter: deceptively simple. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2007;21(5):629–31.	2

연번	서지정보	배제 사유
95	Mayer J, Boldt J, Schollhorn T, Rohm KD, Mengistu AM, Suttner S. Semi-invasive monitoring of cardiac output by a new device using arterial pressure waveform analysis: a comparison with intermittent pulmonary artery thermodilution in patients undergoing cardiac surgery. <i>British Journal of Anaesthesia</i> . 2007;98(2):176–82.	5
96	McGee WT, Raghunathan K. Physiologic goal-directed therapy in the perioperative period: The volume prescription for high-risk patients. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> . 2013;27(6):1079–86.	2
97	Mielck F, Buhre W, Hanekop G, Tirilomis T, Hilgers R, Sonntag H. Comparison of continuous cardiac output measurements in patients after cardiac surgery. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2003;17(2):211–6.	3
98	Missant C, Rex S, Wouters PF. Accuracy of cardiac output measurements with pulse contour analysis (PulseCO) and Doppler echocardiography during off-pump coronary artery bypass grafting. <i>European Journal of Anaesthesiology</i> . 2008;25(3):243–8.	3
99	Moppett IK, Rowlands M, Mannings A, Moran CG, Wiles MD, Investigators N. LiDCO-based fluid management in patients undergoing hip fracture surgery under spinal anaesthesia: a randomized trial and systematic review. <i>British Journal of Anaesthesia</i> . 2015;114(3):444–59.	3
100	Morisawa K, Fujitani S, Homma Y, Shigemitsu K, Saito N, Hayakawa K, et al. Can the global end-diastolic volume index guide fluid management in septic patients? A multicenter randomized controlled trial. <i>Acute Medicine & Surgery</i> . 2020;7(1):e468.	3
101	Mouncey PR, Osborn TM, Power GS, Harrison DA, Sadique MZ, Grieve RD, et al. Protocolised Management In Sepsis (ProMISE): a multicentre randomised controlled trial of the clinical effectiveness and cost-effectiveness of early, goal-directed, protocolised resuscitation for emerging septic shock. <i>Health Technology Assessment (Winchester, England)</i> . 2015;19(97):i–xxv, 1–150.	3
102	Mukkamala R, Kohl BA, Mahajan A. Comparison of accuracy of two uncalibrated pulse contour cardiac output monitors in off-pump coronary artery bypass surgery patients using pulmonary artery catheter-thermodilution as a reference. <i>BMC Anesthesiology</i> . 2021;21(1):189.	4
103	Mutoh T, Kazumata K, Ishikawa T, Terasaka S. Performance of bedside transpulmonary thermodilution monitoring for goal-directed hemodynamic management after subarachnoid hemorrhage. <i>Stroke</i> . 2009;40(7):2368–74.	3
104	Mutoh T, Kazumata K, Terasaka S, Taki Y, Suzuki A, Ishikawa T. Impact of transpulmonary thermodilution-based cardiac contractility and extravascular lung water measurements on clinical outcome of patients with Takotsubo cardiomyopathy after subarachnoid hemorrhage: a retrospective observational study. <i>Critical Care (London, England)</i> . 2014;18(4):482.	4
105	Ni X, Liu XJ, Ding TT. The Application of PiCCO-guided Fluid Resuscitation in Patients With Traumatic Shock. <i>American Surgeon</i> . 2022;31348221087898.	3
106	Nicklas JY, Saugel B. Non-Invasive Hemodynamic Monitoring for Hemodynamic Management in Perioperative Medicine. <i>Frontiers in Medicine</i> . 2017;4:209.	2
107	Orme RM, Pigott DW, Mihm FG. Measurement of cardiac output by transpulmonary arterial thermodilution using a long radial artery catheter. A comparison with intermittent pulmonary artery thermodilution. <i>Anaesthesia</i> . 2004;59(6):590–4.	3
108	Ostergaard M, Nielsen J, Rasmussen JP, Berthelsen PG. Cardiac output--pulse contour analysis vs. pulmonary artery thermodilution. <i>Acta Anaesthesiologica Scandinavica</i> . 2006;50(9):1044–9.	3
109	Paarmann H, Groesdonk HV, Sedemund-Adib B, Hanke T, Heinze H, Heringlake M, et al. Lack of agreement between pulmonary arterial thermodilution cardiac output and the pressure recording analytical method in postoperative cardiac surgery patients. <i>British Journal of Anaesthesia</i> . 2011;106(4):475–81.	3
110	Pala S, Aletti F, Toschi N, Guerrisi M, Coniglione F, Dauri M, et al. Comparisons of predictors of fluid responsiveness in major surgery. Annual International Conference Of The IEEE Engineering In Medicine And Biology Society. 2012;2012:3128–30.	4
111	Pan G, Fan X, Bian J, He Z, Yue J, Sun H, et al. Application and significance of PiCCO monitoring technique combined with troponin I detection in fluid resuscitation of elderly patients with septic myocardial dysfunction. <i>American Journal Of Translational Research</i> .	3

연번	서지정보	배제 사유
	2021;13(6):6846–54.	
112	Pauli C, Fakler U, Genz T, Hennig M, Lorenz HP, Hess J. Cardiac output determination in children: equivalence of the transpulmonary thermodilution method to the direct Fick principle. <i>Intensive Care Medicine</i> . 2002;28(7):947–52.	4
113	Pavlovic G, Diaper J, Ellenberger C, Frei A, Bendjelid K, Bonhomme F, et al. Impact of early haemodynamic goal-directed therapy in patients undergoing emergency surgery: an open prospective, randomised trial. <i>Journal of Clinical Monitoring & Computing</i> . 2016;30(1):87–99.	3
114	Pearson KS, Gomez MN, Moyers JR, Carter JG, Tinker JH. A cost/benefit analysis of randomized invasive monitoring for patients undergoing cardiac surgery. <i>Anesthesia & Analgesia</i> . 1989;69(3):336–41.	3
115	Penttila J, Snapir A, Kentala E, Koskenvuo J, Posti J, Scheinin M, et al. Estimation of cardiac output in a pharmacological trial using a simple method based on arterial blood pressure signal waveform: a comparison with pulmonary thermodilution and echocardiographic methods. <i>European Journal of Clinical Pharmacology</i> . 2006;62(6):401–7.	3
116	Perel A, Saugel B, Teboul JL, Malbrain ML, Belda FJ, Fernandez-Mondejar E, et al. The effects of advanced monitoring on hemodynamic management in critically ill patients: a pre and post questionnaire study. <i>Journal of Clinical Monitoring & Computing</i> . 2016;30(5):511–8.	4
117	Petzoldt M, Riedel C, Braeunig J, Haas S, Goepfert MS, Treede H, et al. Dynamic device properties of pulse contour cardiac output during transcatheter aortic valve implantation. <i>Journal of Clinical Monitoring & Computing</i> . 2015;29(3):323–31.	4
118	Petzoldt M, Riedel C, Braeunig J, Haas S, Goepfert MS, Treede H, et al. Stroke volume determination using transcardiopulmonary thermodilution and arterial pulse contour analysis in severe aortic valve disease. <i>Intensive Care Medicine</i> . 2013;39(4):601–11.	4
119	Pohl T, Kozieras J, Sakka SG. Influence of extravascular lung water on transpulmonary thermodilution-derived cardiac output measurement. <i>Intensive Care Medicine</i> . 2008;34(3):533–7.	4
120	Power P, Bone A, Simpson N, Yap CH, Gower S, Bailey M. Comparison of pulmonary artery catheter, echocardiography, and arterial waveform analysis monitoring in predicting the hemodynamic state during and after cardiac surgery. <i>International Journal of Critical Illness and Injury Science</i> . 2017;7(3):156–62.	4
121	Rauch H, Muller M, Fleischer F, Bauer H, Martin E, Bottiger BW. Pulse contour analysis versus thermodilution in cardiac surgery patients. <i>Acta Anaesthesiologica Scandinavica</i> . 2002;46(4):424–9.	3
122	Ritter S, Rudiger A, Maggiorini M. Transpulmonary thermodilution-derived cardiac function index identifies cardiac dysfunction in acute heart failure and septic patients: an observational study. <i>Critical Care (London, England)</i> . 2009;13(4):R133.	3
123	Rocca GD, Chiarandini P. Hemodynamic Monitoring during Liver Transplantation. <i>International Anesthesiology Clinics</i> . 2017;55(2):121–34.	2
124	Rodig G, Prasser C, Keyl C, Liebold A, Hobbahn J. Continuous cardiac output measurement: pulse contour analysis vs thermodilution technique in cardiac surgical patients. <i>British Journal of Anaesthesia</i> . 1999;82(4):525–30.	3
125	Romano SM, Pistolesi M. Assessment of cardiac output from systemic arterial pressure in humans. <i>Critical Care Medicine</i> . 2002;30(8):1834–41.	3
126	Rowan KM, Angus DC, Bailey M, Barnato AE, Bellomo R, Canter RR, et al. Early, goal-directed therapy for septic shock – A patient-level meta-analysis. <i>New England Journal of Medicine</i> . 2017;376(23):2223–34.	2
127	Sakka SG, Reinhart K, Meier-Hellmann A. Comparison of pulmonary artery and arterial thermodilution cardiac output in critically ill patients. <i>Intensive Care Medicine</i> . 1999;25(8):843–6.	4
128	Sakka SG, Reinhart K, Wegscheider K, Meier-Hellmann A. Is the placement of a pulmonary artery catheter still justified solely for the measurement of cardiac output? <i>Journal of Cardiothoracic and Vascular Anesthesia</i> . 2000;14(2):119–24.	4
129	Samra T, Arya VK. Comparison of cardiac output estimation by FloTrac/Vigileo TM and	6

연번	서지정보	배제 사유
	intermittent pulmonary artery thermodilution in patient with Takayasu arteritis. Annals of Cardiac Anaesthesia. 2011;14(2):163-4.	
130	Sander M, von Heymann C, Foer A, von Dossow V, Grosse J, Dushe S, et al. Pulse contour analysis after normothermic cardiopulmonary bypass in cardiac surgery patients. Critical Care (London, England). 2005;9(6):R729-34.	3
131	Schmid S, Kapfer B, Heim M, Bogdanski R, Anetsberger A, Blobner M, et al. Algorithm-guided goal-directed haemodynamic therapy does not improve renal function after major abdominal surgery compared to good standard clinical care: a prospective randomised trial. Critical Care (London, England). 2016;20:50.	3
132	Scolletta S, Miraldi F, Romano SM, Muzzi L. Continuous cardiac output monitoring with an uncalibrated pulse contour method in patients supported with mechanical pulsatile assist device. Interactive Cardiovascular & Thoracic Surgery. 2011;13(1):52-6.	3
133	Scully CG, Gomatam S, Forrest S, Strauss DG. Importance of re-calibration time on pulse contour analysis agreement with thermodilution measurements of cardiac output: a retrospective analysis of intensive care unit patients. Journal of Clinical Monitoring & Computing. 2016;30(5):577-86.	3
134	Scully TG, Grealy R, McLean AS, Orde SR. Calibrated cardiac output monitoring versus standard care for fluid management in the shocked ICU patient: a pilot randomised controlled trial. Journal of intensive care. 2019;7:1.	3
135	Segal E, Katzenelson R, Berkenstadt H, Perel A. Transpulmonary thermodilution cardiac output measurement using the axillary artery in critically ill patients. Journal of Clinical Anesthesia. 2002;14(3):210-3.	3
136	Shah S, Bhargava A, Hariharan U, Jain C, Kulkarni A, Gupta N. Goal-directed fluid therapy using transoesophageal echocardiographic inferior venacaval index in patients with low left ventricular ejection fraction undergoing major cytoreductive surgery: A clinical trial. Saudi Journal of Anaesthesia. 2020;14(1):7-14.	4
137	Shoemaker WC, Wo CC, Yu S, Farjam F, Thangathurai D. Invasive and noninvasive haemodynamic monitoring of acutely ill sepsis and septic shock patients in the emergency department. European Journal of Emergency Medicine. 2000;7(3):169-75.	3
138	Shoemaker WC, Wo CCJ, Demetriades D, Beez M. Noninvasive haemodynamic monitoring to predict outcome and guide therapy in acute critical illness. International Journal of Intensive Care. 2007;14(1):15-25.	3
139	Sinha AC, Singh PM, Grewal N, Aman M, Dubowitz G. Comparison between continuous non-invasive estimated cardiac output by pulse wave transit time and thermodilution method. Annals of Cardiac Anaesthesia. 2014;17(4):273-7.	3
140	Siranovic M, Kovac J, Gopcevic A, Kelecić M, Vučić M, Kovac N, et al. Constant cardiac output monitoring using the PiCCO and LiDCO methods versus PAK in septic patients: when to do calibration? Acta Clinica Croatica. 2011;50(2):267-72.	3
141	Smetkin AA, Kirov MY, Kuzkov WV, Lenkin AI, Eremeev AV, Slastilin VY, et al. Single transpulmonary thermodilution and continuous monitoring of central venous oxygen saturation during off-pump coronary surgery. Acta Anaesthesiologica Scandinavica. 2009;53(4):505-14.	3
142	Staier K, Wilhelm M, Wiesenack C, Thoma M, Keyl C. Pulmonary artery vs. transpulmonary thermodilution for the assessment of cardiac output in mitral regurgitation: a prospective observational study. European Journal of Anaesthesiology. 2012;29(9):431-7.	3
143	Sujatha P, Mehta Y, Dhar A, Sarkar D, Meharwal ZS, Datt V. Comparison of cardiac output in OPCAB: bolus thermodilution technique versus pulse contour analysis. Annals of Cardiac Anaesthesia. 2006;9(1):44-8.	3
144	Sun Y, Lu ZH, Zhang XS, Geng XP, Cao LJ, Yin L. The effects of fluid resuscitation according to PiCCO on the early stage of severe acute pancreatitis. Pancreatology. 2015;15(5):497-502.	3
145	Takeda C, Takeuchi M, Mizota T, Yonekura H, Nahara I, Joo WJ, et al. The association between arterial pulse waveform analysis device and in-hospital mortality in high-risk non-cardiac surgeries. Acta Anaesthesiologica Scandinavica. 2020;64(7):928-35.	3
146	Tang W, Qiu Y, Lu H, Xu M, Wu J. Stroke Volume Variation-Guided Goal-Directed Fluid Therapy Did Not Significantly Reduce the Incidence of Early Postoperative Complications	3

연번	서지정보	배제 사유
	in Elderly Patients Undergoing Minimally Invasive Esophagectomy: A Randomized Controlled Trial. <i>Frontiers in Surgery</i> . 2021;8:794272.	
147	Tannenbaum GA, Mathews D, Weissman C. Pulse contour cardiac output in surgical intensive care unit patients. <i>Journal of Clinical Anesthesia</i> . 1993;5(6):471–8.	4
148	Terada T, Ochiai R. Comparison of the ability of two continuous cardiac output monitors to detect stroke volume index: Estimated continuous cardiac output estimated by modified pulse wave transit time and measured by an arterial pulse contour-based cardiac output device. <i>Technology & Health Care</i> . 2021;29(3):499–504.	4
149	Thomas B. Monitoring of cardiac output by pulse contour method. <i>Acta Anaesthesiologica Belgica</i> . 1978;29(3):259–70.	3
150	Tran-Dinh A, Augustin P, Dufour G, Lasocki S, Allou N, Thabut G, et al. Evaluation of Cardiac Index and Extravascular Lung Water After Single-Lung Transplantation Using the Transpulmonary Thermodilution Technique by the PiCCO2 Device. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2018;32(4):1731–5.	3
151	Tribuddharat S, Sathitkarnmanee T, Ngamsaengsirisup K, Sornpirom S. Efficacy of early goal-directed therapy using FloTrac/EV1000 to improve postoperative outcomes in patients undergoing off-pump coronary artery bypass surgery: a randomized controlled trial. <i>Journal Of Cardiothoracic Surgery</i> . 2022;17(1):196.	9
152	Tsutsui M, Araki Y, Masui K, Kazama T, Sugo Y, Archer TL, et al. Pulse wave transit time measurements of cardiac output in patients undergoing partial hepatectomy: a comparison of the esCCO system with thermodilution. <i>Anesthesia & Analgesia</i> . 2013;117(6):1307–12.	3
153	Tusman G, Acosta CM, Wallin M, Hallbeck M, Esperatti M, Peralta G, et al. Perioperative Continuous Noninvasive Cardiac Output Monitoring in Cardiac Surgery Patients by a Novel Capnodynamic Method. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2022;36(8 Pt B):2900–7.	4
154	Uchino S, Bellomo R, Morimitsu H, Sugihara M, French C, Stephens D, et al. Pulmonary artery catheter versus pulse contour analysis: a prospective epidemiological study. <i>Critical Care (London, England)</i> . 2006;10(6):R174.	4
155	Vannucci A, Krejci V, Kangrga I. Performance of Vigileo and LiDCOplus cardiac output monitors during a prolonged cardiac arrest and resuscitation. <i>European Journal of Anaesthesiology</i> . 2009;26(10):885–7.	2
156	Velmahos GC, Wo CC, Demetriades D, Murray JA, Cornwell EE, 3rd, Asensio JA, et al. Invasive and non-invasive physiological monitoring of blunt trauma patients in the early period after emergency admission. <i>International Surgery</i> . 1999;84(4):354–60.	3
157	Vetrugno L, Bignami E, Barbariol F, Langiano N, De Lorenzo F, Matellon C, et al. Cardiac output measurement in liver transplantation patients using pulmonary and transpulmonary thermodilution: a comparative study. <i>Journal of Clinical Monitoring & Computing</i> . 2019;33(2):223–31.	3
158	Vilchez Monge AL, Alvarez-Cagigas IT, Perez-Pena J, Olmedilla L, Jimeno C, Sanz J, et al. Cardiac output monitoring with pulmonary versus transpulmonary thermodilution during liver transplantation: Interchangeable methods? <i>Minerva Anestesiologica</i> . 2014;80(11):1178–87.	3
159	Voet M, Overduin CG, Stille EL, Futterer JJ, Lemson J. Safety aspects of the PiCCO thermodilution–cardiac output catheter during magnetic resonance imaging at 3 Tesla. <i>Journal of Clinical Monitoring & Computing</i> . 2022;36(1):141–5.	1
160	Wang F, Qiang X, Jiang S, Shao J, Fang B, Zhou L. The fluid management and hemodynamic characteristics of PiCCO employed on young children with severe hand, foot, and mouth disease—a retrospective study. <i>BMC Infectious Diseases</i> . 2021;21(1):208.	3
161	Weissman C, Ornstein EJ, Young WL. Arterial pulse contour analysis trending of cardiac output: hemodynamic manipulations during cerebral arteriovenous malformation resection. <i>Journal of Clinical Monitoring</i> . 1993;9(5):347–53.	8
162	Werawatganon T, Punyatavorn S, Chatkaew P, Bunburaphong P. Validity and reliability of cardiac output by arterial thermodilution and arterial pulse contour analysis compared with pulmonary artery thermodilution in intensive care unit. <i>Journal of the Medical Association of Thailand</i> . 2003;86 Suppl 2:S323–30.	3
163	Wernly B, Lichtenauer M, Franz M, Fritzenwanger M, Kabisch B, Figulla HR, et al. Pulse contour cardiac output monitoring in acute heart failure patients : Assessment of	3

연번	서지정보	배제 사유
164	hemodynamic measurements. <i>Wiener Klinische Wochenschrift</i> . 2016;128(23–24):864–9. Wiesenack C, Prasser C, Keyl C, Rodig G. Assessment of intrathoracic blood volume as an indicator of cardiac preload: single transpulmonary thermodilution technique versus assessment of pressure preload parameters derived from a pulmonary artery catheter. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2001;15(5):584–8.	4
165	Wouters PF, Quaghebeur B, Sergeant P, Van Hemelrijck J, Vandermeersch E. Cardiac output monitoring using a brachial arterial catheter during off-pump coronary artery bypass grafting. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2005;19(2):160–4.	3
166	Yamada T, Tsutsui M, Sugo Y, Sato T, Akazawa T, Sato N, et al. Multicenter study verifying a method of noninvasive continuous cardiac output measurement using pulse wave transit time: a comparison with intermittent bolus thermodilution cardiac output. <i>Anesthesia & Analgesia</i> . 2012;115(1):82–7.	3
167	Yamashita K, Nishiyama T, Yokoyama T, Abe H, Manabe M. Effects of vasodilation on cardiac output measured by PulseCOTM. <i>Journal of Clinical Monitoring and Computing</i> . 2007;21(6):335–9.	3
168	Yamashita K, Nishiyama T, Yokoyama T, Abe H, Manabe M. The effects of vasodilation on cardiac output measured by PiCCO. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2008;22(5):688–92.	3
169	Yao Y, Guan Y, Zhuang Y, Liu W, Cui J, Sun Y. Effects of fluid resuscitation under the guidance of PICCO on the immune function and inflammatory mediator in patients with septic shock. <i>Biomedical Research (India)</i> . 2017;28(22):9722–5.	3
170	Yao Y, Su M, Guan Y, Gao X, Liu W, Sun Y. Clinical Application of Transabdominal Ultrasound Combined with PICCO in Septic Shock Fluid Resuscitation and Its Predictive Value for Survival Outcome. <i>Ultrasound in Medicine & Biology</i> . 2021;47(11):3196–201.	4
171	Yeo LL, Seow SC, Loh JP, Phua J. Comparison of cardiac output measurement by arterial waveform analysis and pulmonary artery catheter in mitral stenosis. <i>Annals of the Academy of Medicine, Singapore</i> . 2010;39(8):655–7.	6
172	Yuanbo Z, Jin W, Fei S, Liangong L, Xunfa L, Shihai X, et al. ICU management based on PiCCO parameters reduces duration of mechanical ventilation and ICU length of stay in patients with severe thoracic trauma and acute respiratory distress syndrome. <i>Annals of Intensive Care</i> . 2016;6(1):113.	3
173	Zangrillo A, Maj G, Monaco F, Scandroglio AM, Nuzzi M, Plumari V, et al. Cardiac Index Validation Using the Pressure Recording Analytic Method in Unstable Patients. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> . 2010;24(2):265–9.	3
174	Zhang G, Mukkamala R. Continuous and minimally invasive cardiac output monitoring by long time interval analysis of a radial arterial pressure waveform: assessment using a large, public intensive care unit patient database. <i>British Journal of Anaesthesia</i> . 2012;109(3):339–44.	3
175	Zhang G, Mukkamala R. Evaluation of monitoring cardiac output by long time interval analysis of a radial arterial blood pressure waveform using the MIMIC II database. <i>Annual International Conference Of The IEEE Engineering In Medicine And Biology Society</i> . 2010;2010:5217.	3
176	Zhang YB, Zhang ZZ, Li JX, Wang YH, Zhang WL, Tian XL, et al. Application of pulse index continuous cardiac output system in elderly patients with acute myocardial infarction complicated by cardiogenic shock: A prospective randomized study. <i>World Journal of Clinical Cases</i> . 2019;7(11):1291–301.	3
177	Zhu Y, Yang M, Ding L, Chu G, Cheng J, Lv G. Fluid resuscitation based on pulse contour cardiac output monitoring is associated with improved prognosis in adult severe burn patients: a retrospective cohort study. <i>Annals of Palliative Medicine</i> . 2021;10(10):10904–12.	3
178	Zollner C, Haller M, Weis M, Morstedt K, Lamm P, Kilger E, et al. Beat-to-beat measurement of cardiac output by intravascular pulse contour analysis: a prospective criterion standard study in patients after cardiac surgery. <i>Journal of Cardiothoracic & Vascular Anesthesia</i> . 2000;14(2):125–9.	3
179	Zoremba N, Bickenbach J, Krauss B, Rossaint R, Kuhlen R, Schalte G. Comparison of electrical velocimetry and thermodilution techniques for the measurement of cardiac output. <i>Acta Anaesthesiologica Scandinavica</i> . 2007;51(10):1314–9.	4

연번	서지정보	배제 사유
180	Zorrilla-Vaca A, Cata JP, Brown JK, Mehran RJ, Rice D, Mena GE. Goal-Directed Fluid Therapy Does Not Have an Impact on Renal Outcomes in an Enhanced Recovery Program. <i>Annals of Thoracic Surgery</i> . 2022;114(6):2059–65.	4